



Available online at www.sciencedirect.com



Environment International 33 (2007) 714–715

ENVIRONMENT
INTERNATIONAL

www.elsevier.com/locate/envint

Erratum

Erratum to “Evaluation of the usefulness of bird feathers as a non-destructive biomonitoring tool for organic pollutants: A comparative and meta-analytical approach” [Environ. Int. 33 (2007) 328–337]

V.L.B. Jaspers ^{a,*}, S. Voorspoels ^b, A. Covaci ^b, G. Lepoint ^c, M. Eens ^a

^a Department of Biology, University of Antwerp, Campus Drie Eiken, Universiteitsplein 1, 2610 Wilrijk, Belgium

^b Toxicological Center, University of Antwerp, Campus Drie Eiken, Universiteitsplein 1, 2610 Wilrijk, Belgium

^c Laboratory of Oceanology, University of Liège, Allée de la Chimie 3, 4000 Liège, Belgium

Available online 17 April 2007

The Publisher regrets that Fig. 2 of this article was printed with errors. The correct Fig. 2 has been printed on the next page. The publisher apologises for any inconvenience this may have caused.

DOI of original article: [10.1016/j.envint.2006.11.011](https://doi.org/10.1016/j.envint.2006.11.011).

* Corresponding author. Tel.: +32 3 820 22 85; fax: +32 3 820 22 71.
E-mail address: veerle.jaspers@ua.ac.be (V.L.B. Jaspers).



a) PCBs profile in feathers**b) PCBs profile in muscle**

Fig. 2. Percentage contribution of selected PCB congeners to the total PCB load in feathers (a) and in muscle tissue (b) of different aquatic and terrestrial birds from Belgium. Error bars represent 2 SE.