

2944

In: STEEMANNS P. & JAVVAUX E. (eds.), Pre-Cambrian to Palaeozoic Palaeopalynology and Palaeobotany.- Carnets de Géologie / Notebooks on Geology, Brest, Memoir 2005/02, Abstract 10 (CG2005\_M02/10)

LATE FAMENNIAN CORRELATION BY MIOSPORES BETWEEN THE REFRATH 1 BOREHOLE (BERGISCH GLADBACH-PAFFRATH SYNCLINE, GERMANY) AND THE REFERENCE SECTION OF CHANXHE (DINANT SYNCLINE, BELGIUM)

Maurice Stree<sup>1</sup> & Christoph Hartkopf-Froeder<sup>2</sup>

<sup>1</sup> Université de Liège, Département de Géologie, Campus du Sart-Tilman B18, B-4000 Liège 1, Belgium

<sup>2</sup> Geologisches Landesamt Nordrhein-Westfalen, De-Greif-Str. 195, Postfach 1080, D-47710 Krefeld, Germany.

**Keywords:** Famennian, miospores, Germany, Belgium, biostratigraphy.

The Bergisch Gladbach-Paffrath Syncline and the Dinant Syncline are both part of the Ardennes-Rhenish Massif but are separated by some 100 km ([Figure 1](#))




Fig. 1. The Ardennes-Rhenish Massif and the location of the two reference section and borehole, respectively in neritic (Chanxhe) and pelagic (Refrath 1) facies.

In the Upper Devonian succession of the Bergisch Gladbach-Paffrath Syncline (The Bergisches Land), the Knoppenbissen Formation is assigned the “Dasbergian” ([Hartkopf-Froeder et alii](#), 2004). In this locality the thermal alteration index (TAI of 2 to 2+) is remarkably low compared to that of other Upper Devonian sediments in the Ardennes-Rhenish area. Due to this low thermal maturity the preservation of miospores is very good to excellent ([Hartkopf-Froeder &](#)



<author>Streel</author>, 1994; <author>Hartkopf-Fröder</author>, 2004). The sedimentary rocks of the The Refrath 1 Borehole are dark grey, partly calcareous mudstones, and rare marlstones considered to be fine-grained turbidites, Both contain many fossils. Aside from the very well preserved miospores of the <i>Retispora lepidophyta-Knoxisporites literatus</i> (LL) miospore Zone, numerous other microfossils are known <i>i.e.</i> <i>conodonts</i> (<author>Piecha</author>, 2004) and entomozocean ostracodes (<author>Groos-Uffenorde</author>, 2004). Some characteristic ammonoid species are present in the lowermost part of the borehole (<author>Korn</author>, 2004).

In the Dinant Syncline, the Comblain-au-Pont Formation is assigned the “Strunian”. Thermal alteration is rather high therefore miospores in the formation are dark brown. Farther east, the Comblain-au-Pont Formation, the time-equivalent in Belgium of the French Etroeuungt Formation, is made up of some dm- to several m-thick beds of green shales and siltstones interbedded with micaceous, argillaceous or calcareous sandstones. From bottom to top three lithologic units may be distinguished: a shale dominant unit, a sandstone dominant unit and a carbonate dominant unit (<author>Conil</author>, 1964). The latter is the equivalent of the Etroeuungt Limestone <i>s.s.</i> (<author>Bultynck</author> & <author>Dejonghe</author>, 2001). The Chanxhe road section (Ourthe valley, eastern Belgium) displays the three lithologic units of the Comblain-au-Pont Formation. It is a site where miospores, foraminifers and conodonts have been described and occur in a sequence permitting intercorrelation of faunal components. Above the base of the <i>Retispora lepidophyta</i>-<i>Knoxisporites literatus</i> (LL) miospore Zone occurs, between Df3 $\delta$  and Df3 $\epsilon$  foraminifer faunas, a Late <i>expansa</i> conodont fauna. Above the Df3 $\epsilon$  foraminifer fauna occurs the next <i>Retispora lepidophyta</i>-<i>Indotrivates explanatus</i> (LE) miospore Zone. Benthic ostracodes belong to the “Eifelian Mega-Assemblage” (<author>Casier</author> <i>et alii</i>, 2005) and allow ecological interpretation. In the Chanxhe section changes in the diameter of <i>Retispora lepidophyta</i> has been examined in great detail by <author>Maziane</author> <i>et alii</i> (2002). They reports a significant shift of mean size (more than or less than 60  $\mu$ m) near the base of the carbonate dominant unit (around bed 115 on fig. 1). This shift allows the recognition of two varieties of <i>Retispora lepidophyta</i>, <i>i.e. R. l. var. lepidophyta</i> dominant below and <i>R. l. var minor</i> dominant above. Comparison with the range of miospores sizes supports the conclusion that this is not the result of a process of sorting in the carbonate facies but rather a consequence of some evolutionary process.

The Chanxhe road section is a candidate reference section, in neritic facies, for the transition between late Famennian and latest Famennian. Although an international agreement has yet not been reached to define lower limits of substages within the Famennian Stage, a subdivision into 4 substages has been adopted internationally. A proposal for a conodont-based definition of the lower limit of the uppermost Famennian at the base of the Late <i>expansa</i> Zone (<author>Streel</author>, 2005) is still under consideration by the International Subcommittee on Devonian Stratigraphy and, in this context, a search for a good reference section in a neritic facies is more important than ever.






Fig. 2. Late and latest Famennian biostratigraphic correlation between the Comblain-au-Pont Formation at Chanxhe and the Knoppenbissen Formation in the Refrath borehole

The base of the Late *expansa* Zone is absent in the Chanxhe section but new data from the Refrath 1 borehole allow this gap in our knowledge to be filled. In the Refrath 1 borehole, the lower part of the LL miospore Zone (with presumed ancestor *Retispora macroreticulata* and large specimens of *R. lepidophyta*) corresponds to a Middle *expansa* conodont Zone and to the Early *hemisphaerica*–*dichotoma* entomozoacean Zone. The conclusion is that the base of the Late *expansa* conodont Zone and the base of the *Q. kob. kobetusana* (DF3ε) foraminifer Zone are obviously much closer together than was believed previously (Fig. 2).

The few characteristic ammonoid species in the lowermost part of the borehole Refrath 1 imply an age slightly older than the age indicated by the 3 groups of abundant microfossils (Groos-Uffenorde, 2004; Hartkopf-Fröder, 2004; Piecha, 2004). This can be explained only if we accept that during their transport down the slope (Fig. 3) the turbiditic flows removed by erosion the upper levels of some limestone shoals or else that at present only a portion of the ammonoid fauna of the Knoppenbissen Formation is known.




Fig. 3. Location of the Refrath 1 Borehole in the basal facies of the Bergisch Gladbach-Paffath Syncline and easternmost area of the Condroz Facies displaying the Comblain-au-Pont Formation equivalent in Western Germany.

The facies relationships shown here indicates that the turbiditic flows on the slope started when the transgression on the shelf (<author>Maziane</author> <i>et alii</i>, 2002, p. 222) had reached its maximum and was succeeded by cyclic regressive phases.

### References

- <author>Bulyrnck</author> P., <author>Dejonghe</author> L. (2001).-- Devonian lithostratigraphic units (Belgium).-- *Geologica Belgica*, 4/1-2: 39-69.
- <author>Casier</author> J-G, <author>Lebon</author> A., <author>Mamet</author> B., <author>Préat</author> A. (2005).-- Ostracods and lithofacies close to the Devonian-Carboniferous boundary in the Chanxhe and Rivage sections, northeastern part of the Dinant Basin, Belgium.-- *Bull. Inst. Roy. Sc. Nat. Belgique, Sciences de la Terre* 75: 95-126.
- <author>Conil</author> R. (1964).-- Localités et coupes types pour l'étude du Tournaisien inférieur. (Révision des limites sous l'aspect micropaléontologique).-- *Mém. Acad. Roy. Belgique, Classe des Sciences*, 15, 4: 87 p.
- <author>Groos-Uffenorde</author> H. (2004).-- Late Famennian ostracodes from the Refrath 1 Borehole (Bergisch Gladbach-Paffath Syncline; Ardennes-Rhenish Massif, Germany).-- *Cour. Forsch.-Inst. Senckenberg*, 251: 185-235.



- <author>Hartkopf-Fröder</author> C., <author>Stree</author> M. (1994).- Late Famennian miospore assemblages from the Bergisch Gladbach-Paffrath Syncline, Rhenish Slate Mountains, Germany.- Ann. Soc. Geol. Belgique, 116: 333-357.
- <author>Hartkopf-Fröder</author> C. (2004).- Palynostratigraphy of upper Famennian sediments from the Refrath 1 Borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany) .- Cour. Forsch.-Inst. Senckenberg, 251: 77-87
- <author>Hartkopf-Fröder</author> C., <author>Jux</author> U., <author>Knapp</author> G., <author>Piecha</author> M. (2004).- The Late Devonian of the Bergisch Gladbach-Paffrath Syncline (Ardennes-Rhenish Massif, Germany): an overview.- Cour. Forsch.-Inst. Senckenberg, 251: 7-18.
- <author>Korn</author> D. (2004).- Late Famennian cephalopods from the Refrath 1 Borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany) .- Cour. Forsch.-Inst. Senckenberg, 251: 175-183.
- <author>Maziane</author> N., <author>Higgs</author> K.T., <author>Stree</author> M. (2002).- Biometry and paleoenvironment of <i>Retispora lepidophyta</i> (<author>Kedo</author>) <author>Playford</author> 1976 and associated miospores in the latest Famennian nearshore marine facies, eastern Ardenne (Belgium) .- Rev. Palaeobot. Palynol., 118: 211-226.
- <author>Piecha</author> M. (2004).- Late Famennian conodonts from the Refrath 1 Borehole (Bergisch Gladbach-Paffrath Syncline; Ardennes-Rhenish Massif, Germany) .- Cour. Forsch.-Inst. Senckenberg, 251: 253-265.
- <author>Stree</author> M. (2005).- Subdivision of the Famennian Stage into four Substages and correlation with the neritic and continental miospore zonation (SDS Business Meeting Florence, August 23, 2004) .- SDS Newsletter 21 (online version): 14,16-17.