

# Modeling the Distribution of Marketable Timber Products of Private Teak (*Tectona grandis* L.f.) Plantations

Noël H. Fonton<sup>1\*</sup>, Gilbert Atindogbé<sup>1</sup>, Arcadius Y. Akossou<sup>2</sup>, Brice T. Missanon<sup>1</sup>,  
Belarmain Fadohan<sup>1</sup>, Philippe Lejeune<sup>3</sup>

<sup>1</sup>Laboratory of Study and Research in Applied Statistics and Biometry,  
University of Abomey-Calavi, Abomey-Calavi, Benin

<sup>2</sup>Faculty of Agronomy, University of Parakou, Parakou, Benin

<sup>3</sup>Unit of Forest and Nature Management, Gembloux Agro-Bio Tech, University of Liege, Gembloux, Belgium  
Email: [hnfonton@gmail.com](mailto:hnfonton@gmail.com)

Received May 16<sup>th</sup>, 2013; revised July 2<sup>nd</sup>, 2013; accepted July 22<sup>nd</sup>, 2013

Copyright © 2013 Noël H. Fonton et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Management of marketable products of private plantations will not be sustainable without class girth being identifiable readily. Modeling marketable products is a key to obtain good fitness between observed and theoretical girth distribution. We determine the best parameter recovery method with the Weibull function for two silvicultural regimes (coppice and high forest). Data on stand variables were collected from 1101 sample plots. The three Weibull function parameters were estimated with three parameters recovery methods: the maximum likelihood method, the method of moments and the method of percentiles. Stepwise regression and the simultaneously re-estimated parameter using the Seemingly Unrelated Regression Estimation were applied to model each parameter. The results indicated that the three methods successfully predicted girth size distributions within the sample stands. The method of moments was the best one with lowest values of Reynolds error index and Kolmogorov-Smirnov statistic however the silvicultural regimes. The Weibull parameter distribution model developed for each of the two silvicultural regimes was quite reliable.

**Keywords:** Weibull; Parameter Recovery Method; Reynolds Index; Silvicultural Regime; Poles; Logs

## Introduction

The multipurpose management of small woodlots by smallholder forestry has been gaining more importance (Harrison et al., 2002). The growing of the demand for forest products (Scheer, 2004) explained the importance of their management mainly for the smallholder farmer to generate substantial income (Aoudji et al., 2012). Teak (*Tectona grandis* L.f.) is the most important reforestation and commercial plant species in coastal West Africa due to its fast growing potential (Niskane, 1998), good-quality timber (Louppe, 2008). Reforestation with this specie has increased the above ground biomass and carbon stock at 10-year-old about 45% higher than a nearby degraded secondary forest (Odiwe et al., 2012). In Benin, the success of state-owned plantations has encouraged farmers to invest in teak silviculture, establishing plantations on small plots ranging from 0.05 ha to 28.10 ha (Atindogbé, 2012). Various problems constrain both traders and smallholder farmers (Aoudji et al., 2012). These include the lack of market information, high transaction costs, difficulties for traders to get timber supplies (Anyonge and Roshetko, 2003; Nawir et al., 2007), and the low return to smallholder farmers (Maldonado & Louppe, 1999; Nawir et al., 2007). According to the above problems, efforts were needed to have information on the different classes of the merchants products on stand before harvesting. Forest owners

and managers have no reliable tools to provide them with a comprehensive scheme of resources available and monitoring, harvesting and sales operations. Therefore, one challenge is to determine the minimum level of information required to characterize harvests (Lafond et al., 2012).

Stand tables for total or marketable volume are based on the distribution of tree diameters using traditionally probability density functions (PDFs) (Parresol et al., 2010). Many functions have been suggested for establishing tree diameters size class distribution (e.g., normal, exponential, beta, Johnson's  $S_B$ , Gamma, Weibull, logit-logistic). However, the Weibull function appears the most often used (Little, 1983; Rennolls et al., 1985; Rondeux et al., 1992; Lindsay et al., 1996; Liu et al., 2004; Newton & Amponsah, 2005; Lei, 2008) owing to its flexibility (Hafley & Schreuder, 1977; Kilkki et al., 1989) and the best description of diameter structure. This function can also model many types of failure rate behaviors when appropriate parameters are included.

Many techniques for estimating Weibull function parameters have been developed: the graphical methods and the analytical methods. The accuracy of the estimate depends on the size of the sample and the method used. Graphical methods tend to provide crude estimates, while analytical methods provide better estimates that include confidence limits (Murthy et al., 2004) and are reported to be more accurate (Razali et al., 2009). The common analytical methods are the method of moments (MOM),

\*Corresponding author.

the maximum likelihood method (*MLM*), the method of percentiles (*MOP*) and the method of least squares (*MLS*). However, the most suitable method depends on the stands characteristics (Liu et al., 2004; Lei, 2008).

The aim of this study was to determine the best estimator method for Weibull function parameters for two different silvicultural regimes: the coppice and the high forest. The Modeled parameters were then used to predict the distribution of marketable products of the private teak plantations as a useful management tool.

### Methods

#### Study Site and Data

This study was carried out in the Guinea-Congo zone of Benin (West Africa) located between 6°17' and 6°58'N, and 1°56' and 2°31'E. The region has a bimodal rainfall regime, with a mean annual precipitation of 1100 mm and a daily mean temperature of 29.9°C over the period 1971-2009 (www.Worldclim.org, 2005). Clayey-sand and vertisol are the dominant soil types. The original native vegetation, a semi-evergreen dry forest, was strongly influenced by human activities and is now reduced to a few relict forests and forest reserves.

Data were collected using a snowball sampling method which yielded 1101 private teak plantations: 844 coppices and 257 high forests. The size of each plantation (area) was measured. Then five (for plantations <0.5 ha) or ten (for plantations ≥0.5 ha) replicates strips of five trees were randomly sampled. On each strip, the planting space between trees (*e*) and between lines (*l*), the survival rate (*t*), and the girth at breast height (cbh) for all trees over 10 cm (lowest girth size of the marketable products) were measured. Timber merchants use height classes of marketable products based on girth classes: small poles (10 - 19 cm), medium poles (20 - 39 cm), large poles (40 - 49 cm), small posts (50 - 64 cm), large posts (65 - 79 cm), small logs (80 - 109 cm) and large logs (≥110 cm).

#### Statistical Parameter Modeling

The complete three-parameter Weibull probability density function of trees girth *x* is given by (Bailey & Dell, 1973)

$$f(x; \theta) = \frac{\gamma}{\beta} \left( \frac{x - \alpha}{\beta} \right)^{\gamma-1} \exp \left[ - \left( \frac{x - \alpha}{\beta} \right)^\gamma \right] \quad (1)$$

for  $x \geq \alpha$ ,  $\alpha \geq 0$ ,  $\beta > 0$ ,  $\gamma > 0$

where  $\theta = (\alpha, \beta, \gamma)'$ ,  $\alpha$  is a location parameter,  $\beta$  is a scale parameter, and  $\gamma$  is a shape parameter. The recovery methods based on maximum likelihood, the maximum likelihood method (*MLM*), on moments, the method of moment (*MOM*) and on percentiles, the method of percentiles (*MOP*) were compared.

In relation to the unknown parameters  $\alpha$ ,  $\beta$ ,  $\gamma$  and  $n$  the number of trees, the logarithm of the likelihood function,  $\log(L(\theta))$  of Equation (1) is given by:

$$\log L(\theta) = \sum_{i=1}^n \log \left[ \frac{\gamma}{\beta} \left( \frac{x_i - \alpha}{\beta} \right)^{\gamma-1} \exp \left[ - \left( \frac{x_i - \alpha}{\beta} \right)^\gamma \right] \right] \quad (2)$$

For estimating these parameters with the *MLM*, the Equation (2) was maximized with a three-equation system as follows:

$$\begin{cases} \hat{\beta} = \left[ (n-1)(x_i - \hat{\alpha})^{\hat{\gamma}-1} \right] \\ \hat{\gamma} = \left[ \left( \sum_{i=1}^n (x_i - \hat{\alpha})^\gamma \log(x_i - \hat{\alpha}) \right) \left( \sum_{i=1}^n (x_i - \hat{\alpha})^\gamma \right)^{-1} \right. \\ \quad \left. - n^{-1} \sum_{i=1}^n \log(x_i - \hat{\alpha})^{-1} \right] \\ (\hat{\gamma} - 1) \sum_{i=1}^n (x_i - \hat{\alpha}) - \hat{\gamma} \hat{\beta}^{-\hat{\gamma}} \sum_{i=1}^n (x_i - \hat{\alpha})^{\hat{\gamma}} = 0 \end{cases} \quad (3)$$

where  $n$  is the number of trees in the plantation and  $x_i$  the girth of tree  $i$ . The SAS software (SAS 9.2) was used to solve iteratively the equation system (3).

The moment order  $k$  ( $\mu_k$ ) of the Weibull function is given by:

$$\mu_k = \beta^k \sum_{j=0}^k (-1)^j \binom{k}{j} \Gamma \left( \frac{k-1}{\gamma} + 1 \right) \left[ \Gamma \left( \frac{1}{\gamma} + 1 \right) \right]^j \quad (4)$$

with  $\Gamma$  the gamma function written for a real value  $s$  as:

$\Gamma(s) = \int_0^\infty x^{s-1} e^{-x} dx$ , ( $s > 0$ ). The parameters  $\alpha$ ,  $\beta$  and  $\gamma$  were estimated by *MOM* with two processes. The moments of order 1 ( $\mu$ ), order 2 ( $\sigma^2$ ) and order 3 ( $\mu_3$ ) (Razali et al., 2009) of the equation 4 were computed as follows:

$$\begin{cases} \mu = \hat{\alpha} \hat{\beta} + \Gamma \left( 1 + \frac{1}{\hat{\gamma}} \right) \\ \sigma^2 = \hat{\beta}^2 \Gamma \left( 1 + \frac{2}{\hat{\gamma}} \right) - \left[ \Gamma \left( 1 + \frac{1}{\hat{\gamma}} \right) \right]^2 \\ \mu_3 = \hat{\beta}^3 \left\{ \Gamma \left( 1 + \frac{3}{\hat{\gamma}} \right) - 3 \Gamma \left( 1 + \frac{1}{\hat{\gamma}} \right) \Gamma \left( 1 + \frac{2}{\hat{\gamma}} \right) + 2 \left[ \Gamma \left( 1 + \frac{1}{\hat{\gamma}} \right) \right]^3 \right\} \end{cases} \quad (5)$$

The system (5) was solved using the *R* package *rootSolve* (R2.14.1).

The parameter recovery method based on percentiles (*MOP*) requires computation of the 0<sup>th</sup> (minimum girth), 25<sup>th</sup>, 50<sup>th</sup>, and 95<sup>th</sup> percentiles of the distribution of the girth as  $x_0$ ,  $x_{25}$ ,  $x_{50}$ , and  $x_{95}$ , respectively. The three parameters were estimated by solving the following three equations simultaneously (6) (Borders et al., 1987):

$$\begin{cases} \hat{\alpha} = \frac{n^{1/3} x_0 - x_{50}}{n^{1/3} - 1} \\ \hat{\gamma} = \frac{\ln \left[ \frac{\ln(1-0.95)}{\ln(1-0.25)} \right]}{\ln(x_{95} - \hat{\alpha}) - \ln(x_{25} - \hat{\alpha})} \\ \hat{\beta} = - \frac{\hat{\alpha} \Gamma \left( 1 + \frac{1}{\hat{\gamma}} \right)}{\Gamma \left( 1 + \frac{2}{\hat{\gamma}} \right)} \\ \quad + \frac{\alpha^2}{\sqrt{\Gamma^2 \left( 1 + \frac{2}{\hat{\gamma}} \right) \left[ \Gamma^2 \left( 1 + \frac{1}{\hat{\gamma}} \right) - \Gamma \left( 1 + \frac{2}{\hat{\gamma}} \right) \right] + \frac{\bar{x}_g^2}{\Gamma \left( 1 + \frac{2}{\hat{\gamma}} \right)}} \end{cases} \quad (6)$$

where  $n$  is the number of trees in the plantation,  $\Gamma$  is the gamma function,  $\bar{x}_g$  is the quadratic mean girth of the plantation, and  $\ln$  is the natural logarithm.

**Comparison Criteria**

Two statistics were used to assess the goodness of fit of the three methods: the Kolmogorov–Smirnov statistic (KS) and the prediction index error of Reynolds (*e*) (Reynolds et al., 1988). The optimal recovery parameter method is the one with low value for the two criteria. The prediction index error of Reynolds (Equations (7) and (8)) was computed (Pauwels, 2003) as:

$$e(\%) = \frac{\sum_{j=1}^m |N_j - \hat{N}_j|}{N} \times 100 \quad (7)$$

where  $N_j$  is the observed and  $\hat{N}_j$  is the estimated frequency of trees in girth size class  $j$ ,  $N$  is the total number of trees, and  $m$  is the number of classes.  $\hat{N}_j$  is estimated as follows:

$$\hat{N}_j = N \int_{l_j}^{u_j} f(x) dx \quad (8)$$

where  $u_j$  and  $l_j$  are the upper and lower limits of class  $j$ .

**Modeling Weibull Parameters**

Multiple regression method was used to establish the relationship between the estimated parameters  $\alpha$ ,  $\beta$ , and  $\gamma$  (dependent variables) and dendrometric characteristics of stands. The predictor variables were density ( $N \cdot ha^{-1}$ ), surface, and basal area of stand ( $G \cdot m^2 \cdot ha^{-1}$ ); the girth of the tree of mean basal area ( $x_g$ , cm); the mean, maximum, and minimum of girth; and the 25<sup>th</sup> ( $P_{25}$ ), 50<sup>th</sup> ( $P_{50}$ ), and 75<sup>th</sup> ( $P_{75}$ ) percentiles of girth distribution. Models were established and tested for each set of dendrometric characteristics. A stepwise regression was then used to select the best subset of two variables. The estimated parameters were simultaneously re-estimated using the seemingly unrelated regression estimation (SURE). This can account for correlation errors between equations and is asymptotically efficient in the absence of specific errors (Liu et al., 2004).

**Results**

**Data Summary**

Overall, values for girth, density and basal area of trees av-

eraged 21.3 cm, 5170 stems/ha and 18.4  $m^2 \cdot ha^{-1}$ , respectively (**Table 1**). While the mean girth was larger in high forests than in coppices, the reverse trend was obtained for the density and basal area of trees.

**Optimal Method**

Descriptive statistics of the estimated parameters for coppices are presented (**Table 2**). For *MLM*, the parameters  $\hat{\alpha}$ ,  $\hat{\beta}$  and  $\hat{\gamma}$  averaged 8.14, 12.78, and 2.34, respectively. For *MOM*, mean values were 8.97, 11.79, and 2.20, respectively. For *MOP*, the three parameters averaged 9.20, 12.17, and 2.41, respectively. Statistics on the estimated parameters for high forests were presented (**Table 2**). For *MLM*, mean values for the parameters  $\hat{\alpha}$ ,  $\hat{\beta}$  and  $\hat{\gamma}$  varied 9.89, 15.97, and 3.07, respectively. For *MOM*, these values were 11.28, 13.94, 2.52 respectively. For *MOP*, the three parameters means were 10.14, 18.50, 3.72, respectively.

For coppices, the percentages of plantations that fitted the Weibull distribution were 0.95% for *MLM*, 0.97% for *MOM*, and 0.91% for *MOP*. For high forests, these percentages were 0.94% for *MLM*, 0.99% for *MOM*, and 0.96% for *MOP*. The *MOM* method showed the lowest values for the error index of Reynolds (*e*%) and the Kolmogorov-Smirnov statistic (KS) for both coppices and high forests (**Table 3**).

**Table 1.**

Density,  $N$  (/ha), quadratic mean of girth,  $x_g$  (cm) and basal area ( $G \cdot m^2 \cdot ha^{-1}$ ) of the study plantations.

Sylvicultural regimes	Variables	Mean	Min.	Max.	SE
Coppices ( $n = 844$ )	$x_g$ (cm)	20.2	10.7	51.6	0.2
	$N \cdot ha^{-1}$	5952	289	22,458	104
	$G \cdot m^2/ha^{-1}$	20.1	1.0	133.6	0.50
High forests ( $n = 257$ )	$x_g$ (cm)	24.4	11.3	58.4	0.43
	$N \cdot ha^{-1}$	2798.9	632.5	8590.1	69.1
	$G \cdot m^2 \cdot ha^{-1}$	13.2	1.8	58.1	0.45

Min. and Max. are minimal and maximal values of the dendrometric parameters; SE is the standard error of the mean.

**Table 2.**

Estimated parameters of Weibull for the two sylvicultural regimes;  $\alpha$ ,  $\beta$ , and  $\gamma$  are the Weibull position, scale, and shape parameters, respectively. *MLM* is the maximum likelihood method; *MOM* the method of moments and *MOP* the method of percentiles.

Method	$\hat{\alpha}$				$\hat{\beta}$				$\hat{\gamma}$			
	Mean	Min.	Max.	SE	Mean	Min.	Max.	SE	Mean	Min.	Max.	SE
a) Coppices regime												
<i>MLM</i>	8.14	0.00	17.05	0.08	12.78	0.01	55.58	0.23	2.34	0.06	8.43	0.03
<i>MOM</i>	8.97	0.00	19.00	0.10	11.79	1.18	46.70	0.22	2.20	0.56	3.50	0.02
<i>MOP</i>	9.20	0.00	20.00	0.12	12.17	0.66	129.38	0.35	2.41	1.00	32.00	0.08
b) High forests regime												
<i>MLM</i>	9.89	0.00	19.88	0.22	15.47	1.91	65.11	0.53	3.07	0.92	6.30	0.06
<i>MOM</i>	11.28	0.00	19.96	0.25	13.94	2.29	67.08	0.49	2.52	0.68	3.52	0.03
<i>MOP</i>	10.14	0.00	19.62	0.32	18.50	1.32	180.15	1.31	3.72	1.00	39.98	0.27

*MLM* is the maximum likelihood method, *MOM* the method of moments and *MOP* the method of percentiles; Min and max are the minimum and the maximum respectively; SE is the standard error.

**Table 3.**

Comparison of the efficiency of the three parameters estimation methods: values of the error index of Reynolds (e%) and the Kolmogorov-Smirnov statistic (KS) for the two silvicultural regimes.

Methods	Coppices				High forests			
	e%		KS		e%		KS	
	Mean	SE	Mean	SE	Mean	SE	Moy	SE
MLM	13.03	0.40	0.29	0.01	12.60	0.77	0.27	0.02
MOM	11.35	0.31	0.26	0.01	10.12	0.56	0.21	0.01
MOP	19.26	1.15	0.43	0.03	13.12	1.28	0.28	0.03

Consequently, *MOM* was chosen as the most appropriate method for modeling the distribution parameters of the Weibull function for private teak plantations. The models revealed significant differences between coppices and high forests for all three parameters estimated using *MOM* (Table 4).

**Parameters’ Model Development**

The results of the SURE analysis for coppices are presented in Table 5. Stepwise regression revealed that the best subset of two stand characteristics is  $P_{25}$  and  $P_{75}$  for  $\hat{\gamma}$ ,  $P_{75}$  and  $G^2$  for  $\hat{\beta}$ , and  $LP_{50}$  and  $P_{75}$  for  $\hat{\gamma}$  with  $(LP_{50} = \ln(P_{50}/x_g))$ . The final regression equations are:

$$\begin{aligned} \hat{\alpha} &= 6.736 + 0.405P_{25} - 0.167P_{75} \\ \hat{\beta} &= 5.5895 + 0.243P_{75} + 0.001G^2 \\ \hat{\gamma} &= 1.989 + 0.024P_{75} + 8.116\ln(P_{50}/x_g) \end{aligned}$$

The associated  $P_{value}$  are less than 0.001. The model to predict marketable products from coppices is:

$$F(x) = 1 - \exp\left[-\left(\frac{x - 6.736 - 0.405P_{25} + 0.167P_{75}}{5.5895 + 0.243P_{75} + 0.001G^2}\right)^\nu\right]$$

where  $\nu = 1.989 + 0.024P_{75} + 8.116LP_{50}$ ,  $F(x)$  is the Weibull function,  $x$  is the girth of the tree,  $G$  is the basal area,  $P_{25}$ ,  $P_{50}$ , and  $P_{75}$  are the 25<sup>th</sup>, 50<sup>th</sup>, and 75<sup>th</sup> are the  $i$ th percentiles, and  $LP_{50}$  the weighted percentile. Some dependent variables such as  $P_{25}$  and  $P_{75}$  were computed with the best adjustment as follows:

$$P_{25} = -9.79 + 5.69\sqrt{x_g} \text{ and } P_{75} = -26.6 + 11.1\sqrt{x_g}$$

with  $P_{value} = 0.000$ . The results of the SURE analysis for high forest are presented in Table 6 and the best models equations with  $P_{value}$  less than 0.023 are:

$$\begin{aligned} \hat{\alpha} &= 5.249 + 0.476x_{min} - 0.111G \\ \hat{\beta} &= 10.567 + 0.004G^2 - 23.519\ln(P_{50}/x_g) \\ \hat{\gamma} &= 1.867 + 8.283\ln(P_{50}/x_g) + 0.014N^{0.5} \end{aligned}$$

The equation to predict the marketable products for high forests is then:

$$F(x) = 1 - \exp\left[-\left(\frac{x - 5.249 - 0.476x_{min} + 0.111G}{10.567 + 0.004G^2 - 23.519LP_{50}}\right)^w\right]$$

where  $w = 1.867 + 8.283LP_{50} + 0.014\sqrt{N}$ ,  $F(x)$  is the Weibull

**Table 4.**

Comparison of the silvicultural regimes according to estimated parameters of Weibull function by the best estimator method *MOM*.

Parameters	Coppices		High forests		$t$	$P$
	Mean	SE	Mean	SE		
$\hat{\alpha}$	8.97	0.10	11.28	0.25	8.58	0.000
$\hat{\beta}$	11.79	0.22	13.94	0.49	4.00	0.000
$\hat{\gamma}$	2.20	0.02	2.52	0.03	8.88	0.000

$\hat{\alpha}$ ,  $\hat{\beta}$ , and  $\hat{\gamma}$  are the Weibull position, scale, and shape parameters, respectively. *MOM* is the method of moments;  $t$  is the statistic of student, and  $P$  the associated probability value.

distribution function,  $x$  is the girth of the tree,  $x_{min}$  is the minimum girth,  $G$  is the basal area,  $N$  is the density,  $P_{50}$  is the 50<sup>th</sup> percentile or median, and  $LP_{50}$  the weighted 50<sup>th</sup> percentile.

**Discussion**

**Efficiency of the Parameter Recovery Methods**

Regardless of the parameter recovery method used, the observed and theoretical distributions of the plantations were much closed according to the Kolmogorov-Smirnov test. Moreover, for both silvicultural practices, coppices and high forests, the average value of the shape parameter  $\gamma$  of the Weibull distribution was lower than 3.6, suggesting that the distribution of trees is left-skewed. Using the optimal method (*MOM*), the parameter  $\alpha$ , whose value is associated with the minimum girth, was 8.97 for coppices and 11.28 for high forest, both were close to the minimum girth of this study (10 cm).

The parameter  $\beta$ , which gives an idea of the central value of samples, had a maximum value of 46.70 for coppices and 67.08 for high forests, while the values measured, with the completed inventory of 18 plantations, were 51.6 cm for coppices and 58.4 cm for high forests. These results are similar to previous findings, which illustrated that  $\alpha$  is a good predictor of the minimum diameter (Frazier, 1981; Knoebel et al., 1986; Leduc et al., 2001). They also support those of Lei (2008), who demonstrated that the two-parameter Weibull distribution and *MOM* provided the best estimation of the diameter distribution of Chinese pine (*Pinus tabulaeformis*). These results are also consistent with those found by Liu et al. (2004) in their study of the diameter distribution of unthinned plantations of black spruce (*Picea mariana*) in central Canada, although in their study *MOP* was the preferred method. Zhang et al. (2003) previously dem-

**Table 5.**

Regression coefficients of the predictors and statistics resulting from the SURE (Seemingly Unrelated Regression Estimation) analysis for coppices for the response variables i.e.  $\hat{\alpha}$ ,  $\hat{\beta}$ , and  $\hat{\gamma}$  respectively the Weibull position, scale and shape parameters.

	$\hat{\alpha}$			$\hat{\beta}$			$\hat{\gamma}$		
	Constant	$P_{25}$	$P_{75}$	Constant	$P_{75}$	$G^2$	Constant	$P_{75}$	$LP_{50}$
Coef.	6.74	0.405	-0.167	5.895	0.243	0.001	1.989	0.024	8.116
$P$	<0.00	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001
$t$	16.46	8.69	-6.100	7.720	7.130	4.650	25.410	6.810	10.060
SE	0.41	0.05	0.027	0.763	0.034	0.000	0.078	0.004	0.807

Coef. is the regression coefficients,  $G$  is the basal area,  $P_{25}$  and  $P_{75}$  are the 25<sup>th</sup> and 75<sup>th</sup> percentiles, and  $LP_{50} = \ln(P_{50}/x_g)$  where  $x_g$  is the girth of the tree of mean basal area and  $P_{50}$  is the 50<sup>th</sup> percentile,  $t$  is the statistic of Student and  $P$  the associated probability value.

**Table 6.**

Regression coefficients values and their significant appreciation statistics resulting from the SURE (Seemingly Unrelated Regression Estimation) analysis for high forests with the response variables  $\hat{\alpha}$ ,  $\hat{\beta}$ , and  $\hat{\gamma}$  respectively the Weibull position, scale and shape parameters.

	$\hat{\alpha}$			$\hat{\beta}$			$\hat{\gamma}$		
	Constant	$x_{min}$	$G$	Constant	$G^2$	$LP_{50}$	Constant	$LP_{50}$	$N^{1/2}$
Coef	5.249	0.476	-0.111	10.567	0.004	-23.519	1.867	8.283	0.014
$P$	<0.001	<0.001	<0.001	<0.001	0.003	0.023	<0.001	0.003	<0.001
$t$	6.040	7.440	-4.060	15.000	3.640	-2.290	9.800	3.700	4.320
SE	0.870	0.064	0.027	6.705	0.001	23.519	0.191	2.241	0.003

Coef. is the regression coefficients,  $x_{min}$  is the minimum girth,  $G$  is the basal area,  $N$  is the density and  $LP_{50} = \ln(P_{50}/x_g)$  where  $x_g$  is the girth of tree of mean basal area and  $P_{50}$  is the 50<sup>th</sup> percentile,  $t$  is the statistic of Student and  $P$  the associated probability value, SE is the standard error.

onstrated the effectiveness of the Weibull distribution for describing the diameter distribution of natural stands of red spruce (*Picea rubens*) and balsam fir (*Abies balsamea*) in the north-east of North America. Meanwhile, Bailey & Dell (1973) have shown that *MOM* is more efficient than *MOP* for estimating parameters of the Weibull distribution, but it requires very complex calculations. Nanang (1998) in a study on the diameter distribution of *Azadirachta indica* plantations in Ghana asserted the same thing. It was also argued that *MOM* assures compatibility between the characteristics of the observed population used in parameter recovery and those obtained through simulation (Mateus & Tomé, 2011). The differences observed between coppices and high forests for all the three parameters estimated by *MOM* confirm the need to build separate models for different silvicultural regimes.

**Predicting the Weibull Parameters**

Parameters of the Weibull distribution were functions of most of the dendrometric characteristics. In all cases the regression coefficients were statistical significant with  $P$  value less than 0.001 for coppices and 0.023 for high forests. These findings are in agreement with Liu et al. (2004), who modeled the three parameters of a Weibull distribution using four characteristics of black spruce stands (age, basal area, average height, and site index). In most cases, the probability values associated to the regressions were less than 0.0001. Since the distribution model of marketable products from coppices differed from high

forests, a global model combining data from these two silvicultural regimes would not be suitable. This is supported by the observed differences between the estimated parameters. The parameter of the distribution shape,  $\gamma$ , is more influenced by  $LP_{50}$  for both silvicultural practices. The location parameter  $\alpha$  depends on the positional parameters  $P_{25}$  and  $P_{75}$  in coppices, and on the minimum girth and basal area in high forests.  $\beta$  is more influenced by the square of the basal area for both silvicultural regimes. These results differ from those of Torres-Rojo et al. (2000), who found that the shape of the distribution is strongly influenced by the diameter, mean basal area, density, and dominant height of the trees; and that  $\beta$  is influenced by diameter and mean basal area of trees. Moreover, several studies have previously found that the minimum girth most often influences the value of  $\alpha$  (Frazier, 1981; Knoebel et al., 1986; Lejeune, 1994; Leduc et al., 2001).

**Conclusion**

In this study, models have been developed to assess marketable teak resource produced by private teak plantations. The main advantage of modeling parameter with stand characteristics is the compatibility between the characteristics of the observed populations and those obtained through theoretical distributions. Results indicated that the three methods compared were generally suitable for modeling the distribution of marketable products. However, the relative performance of each method depends on its ability to predict the observed girth size

class frequencies. The method based on moment (*MOM*) appears to be the most appropriate.

Distribution models for marketable products were developed for coppices and for high forests using stand variables and *MOM*.

### Acknowledgements

This study was sponsored by the Belgian “*Commission Universitaire pour le Développement (CUD)*”—through le Projet Interuniversitaire Cibl : Contribution au d veloppement d’une fili re du teck au d part des for ts priv es du Sud-B nin (D partement de l’Atlantique)”.

### REFERENCES

- Anyonge, C. H., & Roshetko, J. M. (2003). Farm-level timber production: Orienting farmers towards the market. *Unasylva*, 54, 48-56.
- Aoudji, A. K. N., Ad gbidi, A., Agbo, V., Atindogb , G., Toyi, M. S. S., Y vid , A. S. I., Ganglo, J. C., & Lebailly, P. (2012). Functioning of farm-grown timber value chains: Lessons from the smallholder-produced teak (*Tectona grandis* L.f.) poles value chain in Southern Benin. *Forest Policy and Economics*, 15, 98-107. <http://dx.doi.org/10.1016/j.forpol.2011.10.004>
- Atindogb , G. (2012). *Evaluation et caract risation de la ressource en teck (Tectona grandis L.f.) dans les plantations priv es du Sud-B nin*. Th se de doctorat, B nin: Universit  d’Abomey-Calavi.
- Bailey, R. L., & Dell, T. R. (1973). Quantifying diameter distributions with Weibull function. *Forest Sciences*, 19, 97-104.
- Borders, B. E., Souter, R. A., Bailey, R. L., & Ware, K. D. (1987). Percentile-based distributions characterize forest stand tables. *Forest Sciences*, 33, 570-576.
- Frazier, J. R. (1981). *Compatible whole-stand and diameter distribution models for loblolly pine stands*. Ph.D. Thesis, Blacksburg: Virginia Polytechnic Institute.
- Hafley, W. L., & Schreuder, H. T. (1977). Statistical distributions for fitting diameter and height data in even-aged stands. *Canadian Journal of Forest Research*, 7, 481-487. <http://dx.doi.org/10.1139/x77-062>
- Harrison, S. R., Herbohn, J. L., & Niskanen, A. J. (2002). Non-industrial, smallholder, small-scale and family forestry: What’s in a name? *Small-Scale Forest Economics, Management and Policy*, 1, 1-11.
- Kilkki, P., & Paivinen, R. (1986). Weibull function in the estimation of the basal area dbh-distribution. *Silva Fennica*, 20, 149-156.
- Kilkki, P., Maltamo, M., Mykkanen, R., & Paivinen, R. (1989). Use of the Weibull function in estimating the basal area dbhdistribution. *Silva Fennica*, 23, 311-318.
- Knoebel, B. R., Burkhardt, H. E., & Beck, D. E. (1986). A growth and yield model for thinned stands of yellow-poplar. *Forest Sciences Monograph*, 27, 64.
- Lafond, V., Cordonnier, T., De Coligny, F., & Courbaud, B. (2012). Reconstructing harvesting diameter distribution from aggregate data. *Annals of Forest Science*, 69, 235-243. <http://dx.doi.org/10.1007/s13595-011-0155-2>
- Leduc, D. J., Matney, T. G., Belli, K. L., & Baldwin, V. C. (2001). *Predicting diameter distributions of longleaf pine plantations: A comparison between artificial neural networks and other accepted methodologies*. USDA For Serv Res Pap SRS-25.
- Lei, Y. (2008). Evaluation of three methods for estimating the Weibull distribution parameters of Chinese pine (*Pinus tabulaeformis*). *Forest Sciences*, 54, 566-571.
- Lejeune, P. (1994). Construction d’un mod le de r partition des arbres par classes de grosseur pour des plantations d’ pic a commun (*Picea abies* L Karst) en Ardenne belge. *Annals of Forest Science*, 51, 53-65. <http://dx.doi.org/10.1051/forest:19940104>
- Lindsay, S. R., Wood, G. R., & Woollons, R. C. (1996). Stand table modeling through the Weibull distribution and usage of skewness information. *Forest Ecology and Management*, 81, 19-23. [http://dx.doi.org/10.1016/0378-1127\(95\)03669-5](http://dx.doi.org/10.1016/0378-1127(95)03669-5)
- Little, S. N. (1983). Weibull diameter distributions for mixed stands of western conifers. *Canadian Journal of Forest Research*, 13, 85-88. <http://dx.doi.org/10.1139/x83-012>
- Liu, C., Zhang, S. Y., Lei, Y., Newton, P. F., & Zhang, L. (2004). Evaluation of three methods predicting diameter distributions of black spruce (*Picea mariana*) plantations in central. *Canadian Journal of Forest Research*, 34, 2424-2432. <http://dx.doi.org/10.1139/x04-117>
- Louppe, D. (2008). *Tectona grandis* (L. f). In D. Louppe, A. A. Oteng-Amoako, & M. Brink (Eds.), *Ressources v g tales de l’Afrique Tropicale. Bois d’oeuvre 1*. [Traduction de: *Plant Resources of Tropical Africa. Timbers 1*. 2008]. Wageningen, Pays-Bas: Fondation PROTA; Leiden, Pays-Bas: Backhuys Publishers; Wageningen, Pays-Bas: CTA.
- Maldonado, G., & Louppe, D. (1999). Les plantations villageoises de teck en C te d’Ivoire. *Bois et For ts des Tropiques*, 262, 19-30.
- Mateus, A., & Tom , M. (2011). Modelling the diameter distribution of eucalyptus plantations with Johnson’s SB probability density function: Parameters recovery from a compatible system of equations to predict stand variables. *Annals of Forest Science*, 68, 325-335. <http://dx.doi.org/10.1007/s13595-011-0037-7>
- Murthy, P. D. N., Xie, M., & Jiang, R. (2004). *Weibull models*. Wiley series in probability and statistics. Hoboken.
- Nanang, D. M. (1998). Suitability of the normal, log-normal and Weibull distributions for fitting diameter distributions of neem plantations in Northern Ghana. *Forest Ecology and Management*, 103, 1-7. [http://dx.doi.org/10.1016/S0378-1127\(97\)00155-2](http://dx.doi.org/10.1016/S0378-1127(97)00155-2)
- Nawir, A. A., Kassa, H., Sandewall, M., Dore, D., Campbell, B., Ohlsson, B., & Bekele, M. (2007). Stimulating smallholder tree planting—Lessons from Africa and Asia. *Unasylva*, 58, 53-59.
- Newton, P. F., & Amponsah, I. G. (2005). Evaluation of Weibull-based parameter prediction equation systems for black spruce and jack pine stand types within the context of developing structural stand density management diagrams. *Canadian Journal of Forest Research*, 35, 2996-3010. <http://dx.doi.org/10.1139/x05-216>
- Niskanen, A. (1998). Financial and economic profitability of reforestation in Thailand. *Forest Ecology and Management*, 104, 57-68. [http://dx.doi.org/10.1016/S0378-1127\(97\)00263-6](http://dx.doi.org/10.1016/S0378-1127(97)00263-6)
- Odiwe, A. F., Adewumi, R. A., Alami, A. A., & Ogunsanwo, O. (2012). Carbon stock in topsoil, standing floor litter and above ground biomass in *Tectona grandis* plantation 10-years after establishment in Ile-Ife, Southwestern Nigeria. *International Journal of Biological and Chemical Sciences*, 6, 3006-3016.
- Parresol, B. R., Fonseca, T. F., & Marques, C. P. (2010). *Numerical details and SAS programs for parameter recovery of the SB distribution*. Forest Service, Southern Research Station, General Technical Report SRS-122, United States Department of Agriculture, 31 p.
- Pauwels, D. (2003). *Conception d’un syst me d’aide   la decision pour le choix d’un Scenarrio sylvicole: Application aux peuplements de m l ze en R gion wallonne*. Th se de Doctorat, Gembloux: Facult  Universitaire des Sciences Agronomiques.
- Razali, A. M., Salih, A. A., & Mahdi, A. A. (2009). Best estimate for the parameters of the three parameter Weibull distribution. *Proceedings of the 5th Asian Mathematical Conference, Malaysia*, 2009.
- Rennolls, K., Geary, D. N., & Rollinson, T. J. D. (1985). Characterizing diameter distributions by the use of the Weibull distribution. *Forestry*, 58, 57-66.
- Reynolds, M. R., Burk, T. E., & Huang, W. C. (1988). Goodness of fit tests and model selection procedures for diameter distribution models. *Forest Sciences*, 34, 373-399.
- Rondeux, J., Laurent, C., & Thibaut, A. (1992). Construction d’une table de production pour le douglas (*Pseudotsuga menziesii* Mirb. Franco) en Belgique. *Bulletin des Recherches Agronomiques de Gembloux*, 27, 327-347.
- Scherr, S. J. (2004). Building opportunities for small-farm agroforestry to supply domestic wood markets in developing countries. *Agroforestry Systems*, 61-62, 357-370. <http://dx.doi.org/10.1023/B:AGFO.0000029010.97567.2b>
- Torres-Rojo, J. M., Maga a, O. S., & Acosta, M. (2000). Metodolog a para mejorar la predicci n de par metros de distribuciones diam tricas. *Agrociencia*, 34, 627-637.

WorldClim (2005). *WorldClim Global Climate Data* (GIS Data).  
<http://www.worldclim.org/>

Zhang, L., Packard, K. C., & Liu, C. (2003). A comparison of estimation methods for fitting Weibull and Johnson's SB distributions to

mixed spruce-fir stands in northeastern North America. *Canadian Journal of Forest Research*, 33, 1340-1347.  
<http://dx.doi.org/10.1139/x03-054>