

Abundance study of two magnetic B-type stars in the Orion Nebula Cluster

T. Morel

Institut d'Astrophysique et de Géophysique, Allée du 6 Août, 4000 Liège, Belgium
email: morel@astro.ulg.ac.be

Abstract. We present the results of an abundance analysis of two magnetic B-type stars in the Orion Nebula Cluster that support the lack of a direct relationship between the existence of a magnetic field and a nitrogen excess in the photosphere.

1. Motivation

The effects of magnetic fields on internal mixing in massive stars remain largely unknown. Collecting He and CNO abundances for magnetic OB stars is needed to assess the efficiency of rotational mixing and to constrain evolutionary and interior models (e.g., Meynet et al. 2011). We present the determination of the atmospheric parameters and chemical abundances of two magnetic B-type stars in the very young Orion Nebula Cluster (ONC) based on high-resolution optical spectra acquired with FIES at NOT.

2. The targets

The two ONC stars analysed are LP Ori (HD 36982; B1.5 Vp) and NU Ori (HD 37061; B0.5 V). They have been shown to host a strong magnetic field with a polar strength in the range 600-900 G assuming a bipolar geometry (Petit et al. 2008; Petit & Wade 2012). LP Ori and NU Ori are moderate to fast rotators with $v \sin i \sim 80$ and 200 km s⁻¹, respectively (Petit & Wade 2012; Simón-Díaz et al. 2011). NU Ori is of particular interest because it is one of the fastest-rotating magnetic B stars known.

3. Determination of parameters and chemical abundances

The analysis is based on the non-LTE line-formation code DETAIL/SURFACE and Kurucz model atmospheres. For LP Ori, T_{eff} is derived from Si II/III ionisation balance and $\log g$ from fitting the Balmer line wings. The abundances are inferred from a classical curve-of-growth analysis (Morel 2012, and references therein). The set of unblended lines is selected from a comparison with a spectrum of a slowly-rotating analogue (δ Ceti). A specific method is used for NU Ori because of the high $v \sin i$ (see details in Rauw et al. 2012, and Cazorla et al., these proceedings). The atmospheric parameters and abundances are estimated by finding the best match between a set of observed H, He, and CNO line profiles and a grid of rotationally-broadened, synthetic profiles (see Fig.1 for some illustrative fits). The star is part of an SB1 binary system with $\mathcal{P} \sim 19$ d and a mass ratio above 0.19 (Abt et al. 1991). To explore the impact on our results, we have repeated the analysis using composite, synthetic spectra. We assumed a luminosity ratio of 30%, i.e., that the secondary is a main-sequence B1 star of $\sim 10 M_{\odot}$ with $T_{\text{eff}} = 26\,000$ K, $\log g = 4.3$, and abundances typical of B stars. This may be considered as a rather extreme case (the lines of the secondary are not visible in our spectrum; see Fig.1).




Figure 1. Examples of fit to the He lines in NU Ori (black and red lines: observed and best-fitting synthetic profiles, respectively). The dashed, red lines show the line profiles computed for the final, mean parameters. The quality of the fit was not evaluated within the grey-shaded areas.

Table 1. Results of the analysis.

	LP Ori	NU Ori	
		Companion ignored	Companion taken into account
T_{eff} [K]	$21\,500 \pm 1000$	$30\,800 \pm 1000$	$31\,600 \pm 1000$
$\log g$	4.30 ± 0.15	4.30 ± 0.15	4.30 ± 0.15
ξ [km s^{-1}]	[5] (fixed)	[10] (fixed)	[10] (fixed)
y	0.071 ± 0.018	0.097 ± 0.030	0.101 ± 0.030
$\log \epsilon(\text{C})$	8.27 ± 0.13	7.96 ± 0.19	8.04 ± 0.19
$\log \epsilon(\text{N})$	7.91 ± 0.36	7.68 ± 0.26	7.55 ± 0.26
$\log \epsilon(\text{O})$	8.70 ± 0.42	8.48 ± 0.27	8.53 ± 0.27
[N/C]	-0.36 ± 0.29	-0.28 ± 0.32	-0.49 ± 0.32
[N/O]	-0.79 ± 0.33	-0.80 ± 0.17	-0.98 ± 0.17

4. Results and discussion

The analysis proved difficult because of the substantial line broadening due to rotation. The abundances are weakly constrained, but neither star shows evidence for a He or a N excess (Table 1). The data for NU Ori are compatible with a star that just arrived on the ZAMS and is too young to have dredged up CNO-cycled material to the surface or dramatically spun down because of magnetic braking. These results confirm the diversity of surface nitrogen abundances found in magnetic, early B-type stars (Morel 2012).

Acknowledgements

I acknowledge financial support from Belspo for contract PRODEX GAIA-DPAC.

References

- Abt, H. A., Wang, R., & Cardona, O. 1991, *ApJ* 367, 155
Meynet, G., Eggenberger, P., & Maeder, A. 2011, *A&A* 525, L11
Morel, T. 2012, Vol. 465 of *Astronomical Society of the Pacific Conference Series*, p. 54
Petit, V. & Wade, G. A. 2012, *MNRAS* 420, 773
Petit, V., Wade, G. A., Drissen, L., Montmerle, T., & Alecian, E. 2008, *MNRAS* 387, L23
Rauw, G., Morel, T., & Palate, M. 2012, *A&A* 546, A77
Simón-Díaz, S., García-Rojas, J., Esteban, C., et al. 2011, *A&A* 530, A57