



Université  
de Liège




Measuring the size of the BLR in  
quasars with  
*Microensing-aided reverberation  
mapping*

Dominique Sluse

(University of Liège, Belgium)


with Malte Tewes (AlfA – Bonn - DE)

# Introduction




(Credit: Courbin, Saha, Schechter 2003; Claesken+ 2006)


# Introduction




COSMOGRAIL lightcurve




# Introduction



# Introduction



# Introduction




(Credit: Courbin, Saha, Schechter 2003; Claesken+ 2006)



# Microlensing and broad band filters

Monitoring data are BROAD band: mix several emission components



$\eta_0$  = micro-lens  
Einstein radius

$$\eta_0 = \sqrt{\frac{4G\langle M \rangle}{c^2} \frac{D_{\text{os}} D_{\text{ls}}}{D_{\text{ol}}}}$$
$$\sim 2.03 \times 10^{16} \sqrt{\frac{\langle M \rangle}{0.3 M_{\odot}}} \text{ cm}$$

Continuum  $\sim 0.2 \eta_0$

BLR  $\geq 1 \eta_0$


NLR  $> 10 \eta_0$

# Simulating lightcurves

$$F_1(\lambda, t) = M F_M + M \mu(t) F_{M\mu}$$
$$F_M = f_{\text{BLR}} * l(t), \quad F_{M\mu} = c(t)$$

$$c(t) = \mathcal{GP} \{ \bar{c}, \kappa(t, t') \}$$

$$l(t) = \int \Psi(t - t') c(t') dt'$$



$$\tau = 100 \text{ days}$$

$$f_{\text{BLR}} = 0.2$$

$$\mu = 0.5$$



# Simulating lightcurves

$$F_1(\lambda, t) = M F_M + M \mu F_{M\mu}$$

$$F_2(\lambda, t) = F_M + F_{M\mu}$$

$$M=1, \mu = 0.5, F_M = f_{\text{BLR}} * I(t), F_{M\mu} = c(t)$$


$$\tau = 100 \text{ days}$$

$$f_{\text{BLR}} = 0.2$$


Constant  $\mu$ L:

$$\mu = 0.5$$

M = macro-mag.  
(strong-lensing)



# Measuring a time lag: CCF



# Estimating the microlensing flux

$$\begin{aligned} F_1(t) &= M \overset{\text{BLR}}{F_M} + M \mu_1 \overset{\text{Continuum}}{F_{M\mu}} \\ F_2(t) &= F_M + \mu_2 F_{M\mu} \end{aligned}$$

We can define  $A(t) = \mu_1(t)/\mu_2(t) \times M = M \times \mu(t)$

# Estimating the microlensing flux

BLR

Continuum

$$F_1(t) = M F_M + M \mu_1 F_{M\mu}$$
$$F_2(t) = F_M + \mu_2 F_{M\mu}$$

We can define  $A(t) = \mu_1(t)/\mu_2(t) \times M = M \times \mu(t)$

$$F_M(t) = \frac{-A(t)}{A(t) - M} \left( \frac{F_1(t)}{A(t)} - F_2(t) \right)$$

$$\mu_2(t) F_{M\mu}(t) = \frac{M}{A(t) - M} \left( \frac{F_1(t)}{M} - F_2(t) \right)$$



# Estimating the microlensing flux

$$\begin{aligned} F_1(t) &= M \overset{\text{BLR}}{F_M} + M \mu_1 \overset{\text{Continuum}}{F_{M\mu}} \\ F_2(t) &= F_M + \mu_2 F_{M\mu} \end{aligned}$$

We can define  $A(t) = \mu_1(t)/\mu_2(t) \times M = M \times \mu(t)$

Note that we have  $A(t) = F_1/F_2$  if  $F_M = 0$

In practice we may use an empirical estimate of  $A(t)$ :


$$\hat{A}(t) = \frac{\overset{\text{Spectrum at } t=t_1}{A(t_1)}}{\hat{F}_{12}(t_1)} \hat{F}_{12}(t)$$

Model of large scale variations of  $F_1/F_2$




# Estimating the microlensing flux

Deriving  $A(t_1)$  : Illustration



# Estimating the microlensing flux

$$\hat{A}(t) = \frac{A(t_1)}{\hat{F}_{12}(t_1)} \hat{F}_{12}(t)$$



# Estimating the microlensing flux

BLR                      Continuum


$$F_1(t) = M F_M + M \mu_1 F_{M\mu}$$
$$F_2(t) = F_M + \mu_2 F_{M\mu}$$




$$F_M(t) = \frac{-A(t)}{A(t) - M} \left( \frac{F_1(t)}{A(t)} - F_2(t) \right)$$

$$\mu_2(t) F_{M\mu}(t) = \frac{M}{A(t) - M} \left( \frac{F_1(t)}{M} - F_2(t) \right)$$

# Estimating the microlensing flux




# Estimating the microlensing flux








# Effect of gaps and sampling




# Effect of gaps and sampling




# Realistic microlensing



# Realistic microlensing



# Already in data ?








# Microensing of the BLR

Possible source of noise / additional complexity

HE 0047-1756 (A-B):



CIII]



MgII

# Conclusions

*Proof of concept* that long lightcurves of gravitationally lensed quasars (such as those that will be obtained by LSST) can be used to perform photometric reverberation mapping ... with single band data ! (See [Sluse and Tewes 2014, A&A, 576, A60](#) )

Signal possibly already present in existing data of gravitationally lensed quasars [**STAY TUNED**]

This is complementary to the use of quasar-microlensing to probe the properties of the BLR and of the accretion disc