

Long-Term Safety Follow-up of a Randomized Trial of Darbepoetin Alpha and Intravenous Iron Following Autologous Hematopoietic Cell Transplantation

Aurélie Jaspers, MD*,¹, Frederic Baron, MD PhD²,³, Johan Maertens, MD PhD*,⁴, Bernard J. M. De Prijck, MD*,¹, Henri (Rik) Schots, MDPhD⁵, Christophe Bonnet, MD PhD*,¹,², Kaoutar Hafraoui, MD*,¹, Evelyne Willems, MD PhD*,¹, Sophie Servais, MD PhD*,³, Georges Fillet, MD PhD¹,², and Yves Beguin, MD PhD³,²

+ Author Affiliations

Article	Info & Metrics	E-Letters	PDF
Abstract			
We rece	ently reported the effic	acy of darbepoetin alpha (DA)	(Aranesp®) and
intraver	nous (IV) iron (Venofer	®) on erythroid reconstitution	following
autolog	ous hematopoietic ce	ll transplantation (auto-HCT).	Although short-
term di	sease progression, inf	ections or other safety events	were not

intravenous (IV) iron (Venofer®) on erythroid reconstitution following autologous hematopoietic cell transplantation (auto-HCT). Although short-term disease progression, infections or other safety events were not influenced by DA or IV iron, a longer follow-up was required as some meta-analyses suggest increased mortality in patients with cancer receiving erythropoiesis-stimulating agents (ESA), particularly when no concomitant chemotherapy is administered. Besides, some in vitro and animal studies raised the possible role of iron in tumor growth. Therefore, we analyzed long-term outcome to verify the safety of DA and IV iron.

We included 127 patients with myeloma (n=76), Hodgkin's lymphoma (n=8), aggressive (n=26) or indolent (n=12) non-Hodgkin's lymphoma and other diagnoses (n=5). Groups were well balanced for age, disease, disease stage and disease status at time of transplantation. They were randomized between no erythropoietic therapy (n=25), DA from day 28 post-auto-HCT (n=52) or DA + IV iron on days 28, 42 and 56 (n=50). We collected data about infections and other complications, disease status and overall survival (OS) from auto-HCT (between March 2004 and January 2008) up to January 2014. Patients given a second transplant were censored at that time.

During long-term follow-up (mean: 3.4 years), the incidence of infection remained comparable in the 3 groups. Specifically, 3 patients in the control group (12%), 8 in the DA group (15%) and 6 in the DA+iron group (12%) experienced at least one infection. The number of infections per patient

← Previous		
Volume: 124 Issue: 21 Pages: 3948 - 3948 DOI: http://dx.doi.oi	g/	
Email		Save to My Folders
Citation Alert		© Request Permission
Correction Aler	t	Share
Citation Tools		
Article Info & Metrics		
E-Letters		
Related Articles		-

between day 126 (end of the study) and the last follow-up was 0.2 ± 0.7 , 0.2 ± 0.4 and 0.1 ± 0.3 in control, DA and DA+iron groups, respectively (NS).

During follow-up, 4/25 (16%) patients in control group, 13/52 (25%) in DA group and 6/50 (12%) in the DA+iron group experienced at least one complications (NS). While 3 patients (2 in the control group and 1 in the DA+iron group) presented a thrombosis on study, no other thrombo-embolic event occurred during follow-up. Two patients (1 in control group and 1 in DA group) developed a benign tumor, whereas 4 patients (3 in DA group and 1 in DA+iron group) presented a secondary malignancy (NS). Other complications were: hypertension (n=2, 1 in control and 1 in DA groups), ischemic cardiomyopathy (3 in DA group) with 1 myocardial infarction, 4 arrhythmias (1 in control and 3 in DA groups) with pacemaker requirement in 2 (1 in control and 1 in DA groups), peripheral arteriopathy (1 in DA group), diabetes (n=3, 2 in DA and 1 in DA+iron groups), radiopneumonitis (1 in control group), vasculitis (1 in DA group), nephritis (1 in DA and 1 in DA+iron groups), thyroiditis (1 in DA group), spondylarthritis (1 in DA+iron group), jaw osteonecrosis in myeloma patients treated with zoledronic acid (1 in DA and 1 in DA+iron groups), hip osteonecrosis (1 in DA+iron group), peripheral neuropathy (1 in DA and 1 in DA+iron groups), sudden transient deafness (1 in DA group) and sarcoidosis (1 in DA+iron group).

We did not observe any difference in survival. Indeed, 1-year OS were 100%, 88% and 100% and 5-year OS were 86%, 78% and 91% in the control, DA and DA+iron groups, respectively (p=0.43). Progression-free survival (PFS) were also similar in the 3 groups: 1-year and 5-year PFS were 85% and 71% in the control group, compared to 86% and 57% in the DA group and 94% and 78% in the DA+iron group (p=0.30).

In conclusion, DA and IV iron therapy following auto-HCT did not affect safety, disease outcome or survival in long-term analyses.

Disclosures Beguin: *Amgen:* Consultancy, Speakers Bureau; *Vifor:* Consultancy, Speakers Bureau.

 —* Asterisk with author names denotes non-ASH members.

© 2014 by The American Society of Hematology

▲ Back to top

Articles by Jaspers, A.
Articles by Beguin, Y.

Articles by Jaspers, A.
Articles by Beguin, Y.







Leading the way in experimental and clinical research in hematology

American Society of Hematology 2021 L Street NW, Suite 900, Washington, DC 20036 Phone 202-776-0544 | Fax 202-776-0545

Current Issue Subscriptions		Submit to <i>Blood</i>	Information for:		
First Edition	About Blood	Alerts	Authors		
Topics	Newsroom	RSS	Subscribers		
Collections	Public Access	Blood App	Institutions/Librarians		
All Issues	Permissions	Contact Us	Advertisers		
Abstracts	Order Reprints	Feedback	ASH Privacy Policy		

	ASH Home	Research	Education	Advocacy	Meetings	ASH Store	f	¥	in	
--	----------	----------	-----------	----------	----------	-----------	---	---	----	--

Copyright © 2015 by American Society of Hematology