

Treatment of multiple myeloma with high-risk cytogenetics: a consensus of the International Myeloma Working Group

Pieter Sonneveld,¹ Hervé Avet-Loiseau,² Sagar Lonial,³ Saad Usmani,⁴ David Siegel,⁵ Kenneth C. Anderson,⁶ Wee-Joo Chng,⁷ Philippe Moreau,⁸ Michel Attal,⁹ Robert A. Kyle,¹⁰ Jo Caers,¹¹ Jens Hillengass,¹² Jesús San Miguel,¹³ Niels W. C. J. van de Donk,¹⁴ Hermann Einsele,¹⁵ Joan Bladé,¹⁶ Brian G. M. Durie,¹⁷ Hartmut Goldschmidt,¹⁸ María-Victoria Mateos,¹⁹ Antonio Palumbo,²⁰ and Robert Orlowski²¹

¹Department of Hematology, Erasmus MC Cancer Institute, Rotterdam, The Netherlands; ²Laboratory for Genomics in Myeloma, University Cancer Center of Toulouse, Toulouse, France; ³Winship Cancer Institute, Emory University Medical School, Atlanta, GA; ⁴Levine Cancer Institute/Carolinas Healthcare System, Charlotte, NC; ⁵John Theurer Cancer Center, Hackensack University Medical Center, Hackensack, NJ; ⁶Jerome Lipper Multiple Myeloma Center, Dana-Farber Cancer Institute, Boston, MA; ⁷Division of Hematology, National University Cancer Institute, Singapore; ⁸Department of Hematology, University Hospital, Nantes, France; ⁹Department of Hematology, Purpan University Hospital, Toulouse, France; ¹⁰Department of Laboratory Medicine and Pathology, Mayo Clinic, Minneapolis, MN; ¹¹Department of Hematology, Centre Hospitalier Universitaire de Liège, Liège, Belgium; ¹²Medical Clinic V, Section for Multiple Myeloma, University of Heidelberg, Heidelberg, Germany; ¹³Clinica Universidad de Navarra, Centro de Investigación Médica Aplicada, Pamplona, Spain; ¹⁴Department of Hematology, Free University Medical Center Amsterdam, Amsterdam, The Netherlands; ¹⁵Medical Clinic 2, Universitätsklinik Würzburg, Würzburg, Germany; ¹⁶University Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer, Barcelona, Spain; ¹⁷International Myeloma Foundation, North Hollywood, CA; ¹⁸University Hospital Heidelberg and National Center for Tumor Diseases Heidelberg, Heidelberg, Germany; ¹⁹Department of Hematology, University Hospital of Salamanca, Salamanca, Spain; ²⁰Myeloma Unit, University of Torino, Torino, Italy; and ²¹Division of Cancer Medicine, Department of Lymphoma/Myeloma, MD Anderson Cancer Center, Houston, TX

The International Myeloma Working Group consensus updates the definition for high-risk (HR) multiple myeloma based on cytogenetics. Several cytogenetic abnormalities such as t(4;14), del(17/17p), t(14;16), t(14;20), nonhyperdiploidy, and gain(1q) were identified that confer poor prognosis. The prognosis of patients showing these abnormalities may vary with the choice of therapy. Treatment strategies have shown promise for HR cytogenetic diseases, such as proteasome inhibition in combination with lenalidomide/

pomalidomide, double autologous stem cell transplant plus bortezomib, or combination of immunotherapy with lenalidomide or pomalidomide. Careful analysis of cytogenetic subgroups in trials comparing different treatments remains an important goal. Cross-trial comparisons may provide insight into the effect of new drugs in patients with cytogenetic abnormalities. However, to achieve this, consensus on definitions of analytical techniques, proportion of abnormal cells, and treatment regimens is needed. Based on data available today,

bortezomib and carfilzomib treatment appear to improve complete response, progression-free survival, and overall survival in t(4;14) and del(17/17p), whereas lenalidomide may be associated with improved progression-free survival in t(4;14) and del(17/17p). Patients with multiple adverse cytogenetic abnormalities do not benefit from these agents. FISH data are implemented in the revised International Staging System for risk stratification. (*Blood*. 2016;127(24):2955-2962)

Introduction

Multiple myeloma (MM) is a proliferation of monoclonal plasma cells that produce a monoclonal protein.¹ Indications for treatment are based on end-organ damage (hypercalcemia, renal impairment, anemia, bone lesions) and markers of active disease (ie, an involved:uninvolved serum-free light-chain ratio ≥ 100 , bone marrow plasma cells $\geq 60\%$, or >1 lesion found on magnetic resonance imaging).²

Response to treatment and survival of newly diagnosed MM (NDMM) is heterogeneous, with median overall survival (OS) ranging from 2 to >10 years. MM is characterized by chromosomal instability, and cytogenetic abnormalities (CA) have an impact on prognosis.¹⁻⁴ This perspective will define high-risk (HR) CA and provide recommendations for treatment of HR NDMM patients.

Methods

We will describe techniques for identification of CA followed by a discussion of prognostic impact and treatments. This perspective was developed by an

international expert panel based on evidence of published studies through November 15, 2015. The statement was drafted and circulated among all panel members, followed by rounds of revision.

Diagnostic techniques for CA

Conventional karyotyping. Karyotyping reveals CA in 20% to 30% of patients, those being mainly numerical abnormalities. Several translocations including t(4;14) are not detected. The normal karyotype in patients with a low proliferation index corresponds to the kinetics of normal bone marrow cells. Abnormal karyotype had an unfavorable impact in the Total Therapy programs.⁵ Because more sensitive techniques reveal CA in nearly all MM, karyotyping is not a routine test.

Fluorescence in situ hybridization. Fluorescence in situ hybridization (FISH) is performed in interphase cells, thereby overcoming the problem of karyotyping. Purification of CD138-expressing plasma cells or dual staining for cytoplasmic immunoglobulin (Ig) and FISH are required for FISH. Currently, FISH is the standard technique for analysis of CA. Samples are usually screened for CA, which occur in $>1\%$ of patients. FISH is a practical cytogenetic tool to detect genomic aberration in situ and to

Table 1. Primary and secondary genetic events that can be identified by FISH

Primary genetic events			Secondary genetic events		
IgH translocation	Gene(s)	Frequency (%)	Deletion	Gene(s)	Frequency (%)
t(4;14)	<i>FGFR3/MMSET</i>	15	1p	<i>CDKN2C, FAF1, FAM46C</i>	30
t(6;14)	<i>CCND3</i>	4	6q		33
t(11;14)	<i>CCND1</i>	20	8p		25
t(14;16)	<i>MAF</i>	4	13	<i>RB1, DIS3</i>	44
t(14;20)	<i>MAFB</i>	1	11q	<i>BIRC2/BIRC3</i>	7
			14q	<i>TRAF3</i>	38
			16q	<i>WWOX, CYLD</i>	35
			17p	<i>TP53</i>	7
Hyperdiploidy			Gain		
Trisomies of chromosomes 3, 5, 7, 9, 11, 15, 19, 21	NA	50	1q	<i>CKS1B, ANP32E</i>	40

enumerate the percentage of cells harboring such abnormalities. It does not detect single-nucleotide variants.⁶ For example, TP53 on chromosome 17p is deleted in 7% of myeloma, yet mutated at a much higher frequency in myeloma based on exome sequencing. Knowing these restrictions, FISH testing may include gain(1q), del(1p), t(4;14)(p16;q32), t(14;16)(q32;q23), del(17p13), and a marker for aneuploidy (Table 1). For routine diagnosis, testing of t(4;14) and del(17p13) suffices.

Single-nucleotide polymorphism-based mapping arrays. High-resolution genome-wide analysis (GWAS) of single-nucleotide polymorphisms (SNP) detects regions with loss of heterozygosity and numerical abnormalities. SNP mapping arrays identify copy number variations (CNV). Translocations are not usually detected and will require additional FISH.

Comparative genomic hybridization. Array-based comparative genomic hybridization is a tool for genome-wide classification of CNVs, which primarily detects numerical abnormalities.

Gene expression profiling. Gene expression profiling (GEP) is a technique to identify expression of genes and pathways. Based on RNA expression using microarrays, subgroups of patients are identified with a unique GEP phenotype that partly corresponds to the TC classification.⁷ GEP from patients in clinical trials can be used to identify HR profiles with significant prognostic significance.⁸

Consensus statement. FISH is the standard approach for identification of primary genetic events and secondary numerical events. SNP-based mapping arrays and CGH are more sensitive techniques to detect small numerical aberrations, and therefore these can be used in clinical trials. GEP profiling is useful for prognostication and may require bioinformatics support.

High-risk CA

IgH translocations. In MM, primary events are chromosome translocations involving the immunoglobulin heavy chain (IgH) locus and hyperdiploidy, with multiple copies of odd-numbered chromosomes (Table 1).⁹ IgH translocations are observed in 40% of cases. Frequently involved partner chromosomes/loci are 4p16 (*FGFR3/MMSET*) (12%-15%), 11q13 (*CCND1*) (15%-20%), 16q23 (*MAF*) (3%), 6p21 (*CCND3*) (<5%), and 20q11 (*MAFB*) (1%).¹⁰

Translocation t(4;14) leads to deregulation of fibroblast growth factor receptor 3 (*FGFR3*) and multiple myeloma SET domain (*MMSET*).^{11,12} Because *FGFR3* is not expressed in one third of patients with t(4;14), the target gene is most likely *MMSET*.¹³ t(4;14) is associated with impaired progression-free survival/overall survival (PFS/OS).¹⁴ Importantly, bortezomib seems to improve the negative prognostic impact of t(4;14).¹⁵⁻¹⁸ Prolonged survival was reported in t(4;14) treated with high-dose therapy (HDT) and tandem autologous stem cell transplant (ASCT).^{19,20} SNP arrays showed the heterogeneous adverse impact of t(4;14) related to concomitant CA.²¹

Translocation t(14;16) results in deregulation of the *c-MAF* proto-oncogene and predicts poor outcome.^{11,12,22} An Intergroupe Francophone de Myelome (IFM) analysis showed no adverse impact of t(14;16), possibly because 60% of patients received a double ASCT.²³ Translocation t(14;20) results in deregulation of the *MAFB* oncogene and confers a poor prognosis.¹²

Translocation t(11;14) results in upregulation of cyclinD1 and was identified as favorable in some studies, whereas it had no impact in others.^{14,24,25} This translocation is associated with CD20 expression and a lymphoplasmacytic morphology. In general t(6;14), t(11;14), gain(5q), and hyperdiploidy do not confer poor prognosis.

Genomic imbalance. Hyperdiploidy, which occurs in ~50% of NDMM, is associated with improved PFS/OS.^{11,25} In the MRC IX trial, coexisting hyperdiploidy did not abrogate the poor prognosis of adverse CA.²⁶ In contrast, in a retrospective analysis, PFS of patients with t(4;14) was negatively affected by del(1p32), del22q, and >30 structural CA, whereas del(6q) worsened PFS and del(1p32) worsened OS, and >8 numerical changes improved OS in del(17p).²¹ Modern techniques (GWAS) identify additional CNV above karyotypic hyperdiploidy.²⁷

Del(13q) predicts impaired PFS/OS when detected by karyotyping.²⁸ The adverse impact of del(13q) by FISH is associated with del(17p) and t(4;14). del(13q) as single CA does not confer poor survival.^{11,12,25,29}

Del(17p) or del(17) has a negative impact on PFS/OS. Deletion of *TP53* induces clonal immortalization and survival of tumor cells.³⁰

Patients with ≥ 3 copies of 1q have a worse treatment outcome, reflecting a dosage effect of genes such as *CKS1B*.^{12,29,31} Gain(1q) frequently coincides with del(1p32), which confers poor prognosis.^{21,32-34} Hypodiploidy is regarded as a poor prognostic CA.

It is currently unclear which minimum percentage of cells carrying del(17p) is required for an adverse prognosis or whether this varies with the choice of therapy and stage of disease. Minimal percentages of 20% and 60% have been recommended for del(17p).^{12,21} An international effort will address this issue in a meta-analysis.

The prognostic impact of CA may vary from diagnosis to (refractory) disease because of the selection of subclonal disease.³⁵ In solitary plasmacytoma or extramedullary disease, del(17p) may occur more frequently.^{36,37}

Multiple adverse CA. Among patients with an adverse IgH translocation 62% have gain(1q) compared with 32.4% in controls.¹² The frequency of del(17p) is similar in patients without adverse IgH translocations. Among patients with an IgH translocation and/or gain 1q or del(17p), 20% shared ≥ 2 CA. When CA occurred in isolation, each lesion had a similar impact on OS. The triple combination of an adverse IGH translocation, gain(1q), and del(17p) was associated with a median OS of 9.1 months,¹² demonstrating the progressive impact of cosegregation of multiple adverse CA on OS. The IFM showed that in 110 patients displaying either t(4;14) or del(17p), 25 had both abnormalities. In patients with t(4;14), PFS was worse with concomitant del(1p32), del(22q), and/or >30 structural changes, whereas del(13q14), del(1p32), and higher number of CA shortened OS. In patients with del(17p), del(6q) reduced PFS, whereas gain15 and del14 had a protective effect. Del(1p32) shortened OS, whereas >8 numerical changes improved OS.²¹

Good combined with adverse CA. Gain of 5(q31) improved outcome with hyperdiploid MM.³⁸ Among patients with hyperdiploidy, trisomies 3 and 5 confer a favorable prognosis.²¹

Table 2. Summary of cytogenetic risk features

	High-risk	Standard-risk
Cytogenetic abnormality	FISH: t(4;14), t(14;16), t(14;20), del(17/17p), gain(1q) Nonhyperdiploid karyotype Karyotype del(13) GEP: high-risk signature	All others including: FISH: t(11;14), t(6;14)

In the Myeloma IX study, 58% of patients had hyperdiploidy.³⁹ Of these, 61% had ≥ 1 adverse lesion (t(4;14), t(14;16), t(14;20), gain1q, or del(17p)). OS and PFS were worse in patients with hyperdiploidy plus an adverse lesion, compared with hyperdiploidy alone (median PFS, 23 vs 15.4 months; median OS, 60.9 vs 35.7 months). Alternatively, presence of hyperdiploidy did not change the outcome in patients with an adverse lesion.

Presence of trisomies in patients with t(4;14), t(14;16), t(14;20), or *TP53* deletion in MM reduced their adverse impact.⁴⁰

Cytogenetic risk classifications

The definition of HR disease is subject to diagnostic and treatment options. With median PFS and OS of transplant-eligible (TE) patients approaching 4 and 10 years, most investigators consider HR disease as OS <3 years, with ultra-HR disease having a survival <2 years. For non-transplant-eligible patients (NTE) OS, <2 years is considered HR.^{41,42} It is important to define HR disease based on objective criteria.

Risk classifications based on FISH. IMWG proposed a model of HR MM defined as at least one of the following: del17p, t(4;14), or t(14;16) determined by FISH.⁴³ The Mayo Clinic classification added hypodiploidy and t(14;20) to the definition of HR MM (Table 2).⁴⁴ Later classifications attempted to separate MM into several risk groups. In MRC IX, 3 groups were identified (ie, favorable risk [FR: no adverse IgH translocation, del(17p), or gain(1q)], intermediate risk (IR: 1 adverse CA), and HR (>1 adverse CA). Median PFS/OS of patients with FR, IR, or HR was 23.5, 17.8, and 11.7 months and 60.6, 41.9, and 21.7 months, respectively.¹² Ultra-HR was defined as ≥ 3 CA (2%; median OS, 9 months). These classifications may change with treatment modalities. An example is t(4;14), which may be IR rather than HR when novel agents are given.^{15,45,46} IMWG stated that HR MM should include t(4;14), t(14;16), or del(17p).⁴⁷

Risk classifications based on FISH and ISS. The combination of International Staging System (ISS) with HR CA reflects tumor mass, patient condition, and genetics. IMWG showed that t(4;14) and/or del(17p) separates

Table 3. Survival of MM patients with high-risk FISH compared with those without high-risk FISH

FISH	Np/Na	End point	Therapy	Present	Absent	Comment	Ref
Conventional therapy							
t(4;14)	42/290	3-y OS	VBMCP	24%	64%	E9486	13
	100/616	3-y OS	VAD + ASCT \times 2	55%	80%	IFM-99	25
	98/414	3-y OS	VAD + ASCT \times 1/2	40%	72%	IFM-2005	68
del17p	37/308	3-y OS	VBMCP	32%	68%	E9486	13
	58/474	3-y OS	VAD + ASCT \times 2	50%	78%	IFM-99	25
	119/393	3-y OS	VAD + ASCT \times 1	49%	82%	IFM-2005	68
Unfavorable FISH	141/166	3-y OS	CVAD + ASCT \times 1	58%	81%	MRC IX intensive	62
	90/125	3-y OS	MP	26%	48%	MRC IX non-intensive	61
	98/129	3-y OS	Placebo maintenance	69%	72%	MRC IX maintenance	39
	18/111	3-y OS	VBMCP/VBAD +Bz \times 2 + ASCT \times 1	48%	84%	GEM2005 <65	63
Thalidomide							
t(4;14)	57/181	3-y PFS	TD + ASCT \times 2 + TD	20%	48%	GIMEMA	102
	26/156	3-y OS	VAD + ASCT \times 1 + Thal maintenance	44%	79%	HOVON65/GMMG-HD4	29
del17p	21/161	3-y OS	VAD + ASCT \times 1 + Thal maintenance	17%	79%	HOVON65/GMMG-HD4	29
Unfavorable FISH	43/302	5-y OS	Thal induction, consolidation, maintenance	56%	72%	Total Therapy 2	18
	152/167	3-y OS	CTD + ASCT \times 1	59%	82%	MRC IX intensive	62
	96/129	3-y OS	CTDa	58%	78%	MRX IX non-intensive	61
	99/126	3-y OS	Thalidomide maintenance	45%	76%	MRX IX maintenance	39
	17/110	3-y OS	TD + ASCT \times 1	56%	86%	GEM2005 < 65	63
Lenalidomide							
t(4;14)	28/102	Median OS	RD in RRMM	18 m	23 m	MM-016	103
	26/158	Median OS	RD in RRMM	9 m	15 m	IFM	83
	152/355	Median PFS	Lenalidomide maintenance	27 m	42 m	IFM-2005	68
del17p	12/118	Median OS	RD in RRMM	4 m	23 m	MM-016	103
	6.6%	Median PFS	Lenalidomide maintenance	29 m	42 m	IFM-2005	68
Unfavorable FISH	16/84	3-y OS	RD	77%	86%	Mayo Clinic	76
	21/105	2-y OS	RD	76%	91%	E4A03	104
Bortezomib							
t(4;14)	106/401	4-y OS	VD + ASCT \times 1	63%	85%	IFM-2005	68
	53/183	3-y PFS	VTD + ASCT \times 2 + BzTD	65%	61%	GIMEMA	102
	24/148	3-y OS	VAD + ASCT \times 1 + Bz	66%	82%	HOVON65/GMMG-HD4	29
del17p	54/453	4-y OS	VD + ASCT \times 1	50%	79%	IFM-2005	68
	16/158	3-y OS	VAD + ASCT \times 1 + Bz	69%	82%	HOVON65/GMMG-HD4	29
	18/112	3-y OS	VTD + ASCT \times 1	60%	88%	GEM2005 <65	63
Unfavorable FISH	44/188	3-y OS	VMP/BzTP, BzT/BzP	55%	73%	GEM2005 <65	73
	28/140	3-y OS	VMP	56%	71%	VISTA	72

Table 4. Survival of high-risk genetic subgroups in randomized, controlled clinical trials of newly diagnosed MM: effect of treatment modalities and novel drugs

FISH	N1/N2	End point	Arm 1	Arm 2	Arm 1 (%)	Arm 2 (%)	Comment	Ref
t(4;14)	26/24	3-y OS	PAD/ASCT/thalidomide*	VAD/ASCT/bortezomib*	44	66	HOVON65/GMMG- HD4	15
	98/106	4-y OS	VAD	VD	32	63*	IFM-2005	68
	21/23	2-y OS	Thalidomide*	Placebo*	67	87	TT2	18
	21/29	2-y OS	Thalidomide-TT2	Bortezomib TT3	67	97*	TT2 vs TT3	70
Del(17p)	21/16	3-y OS	VAD/ASCT/thalidomide	PAD/ASCT/bortezomib*	17	69*	HOVON65/GMMG-HD4	15
	119/54	4-y OS	VAD	V D	36	50	IFM-2005	68
Nonhyperdiploid	92	3-y OS	VTD	VMP	53	72*	PETHEMA	63
Unfavorable FISH	152/141	3-y OS	CTD	VAD-cyclophosphamide	58	56	MRC IX intensive	62
	96/90	3-y OS	CTD	Placebo MP	34	26	MRC IX nonintensive	61
	99/98	3-y OS	Thalidomide	Placebo	45	69*	MRC IX maintenance	39

Adapted from Bergsagel et al.⁵⁸

*Significant better survival outcome.

2 groups with different event-free survival (EFS) and OS within each ISS stage.⁴⁸ Combining t(4;14) and del(17p) with ISS stage improved prognostic staging.⁴⁸ Neben et al combined ISS with t(4;14) or del(17p).¹¹ Median PFS after ASCT were 2.7, 2.0, and 1.2 years for the FR group (ISS I, no HR CA), IR (ISS I and HR CA or ISS II/III without HR CA), and HR (ISS II/III and HR CA). Five-year OS were 72%, 62%, and 41%, respectively. Identical results were obtained in the Hovon-65/GMMG-HD4 trial.²⁹

The MRC IX study combined ISS and the presence of 0, 1, or >1 adverse CA. Median OS in the ultra-HR group, defined by ISS II or III plus >1 adverse CA, were 9.9 and 19.4 months, compared with OS 67.8 months in the favorable group.¹²

Risk classifications based on FISH, ISS, and lactodehydrogenase.

A meta-analysis of randomized trials in NDMM confirmed that combining ISS, serum lactodehydrogenase (LDH), and FISH identifies 4 risk groups including a very HR population (5%-8%). Patients with ISS stage III, elevated LDH, and t(4;14) or del(17p) have a 2-year OS of only 54.6%.⁴⁹ More recently, the revised ISS was defined, incorporating HR FISH (t(4;14), t(14;16), and del(17p) with ISS and LDH.⁵⁰

Gene expression profiling.

The prognostic impact of GEP by microarray was examined in several studies. The UAMS identified a 70-gene subset as an independent prognosticator.⁵¹ Presence of a HR signature (13.1%) resulted in inferior EFS (5-year EFS: 18% vs 60%) and OS (5-year OS: 28% vs 78%). In this signature, several genes mapped to chromosome 1(q) and (1p).²⁰ The same group performed GEP analysis 48 hours after bortezomib dosing in TT3.⁵² Based on GEP changes, the UAMS-80 signature was constructed.

The EMC-92 signature was derived from patients in the Hovon-65/GMMG-HD4 trial. When combined with ISS, it predicts impaired PFS and OS across treatments.^{53,54} OS of HR patients (21%) at 5 years was 10% compared with 72% for others (79%). Other GEP-based risk models include the IFM-15 and MRC IX-6 gene signatures.⁵⁵ In general, these GEP signatures are useful for prognostication, whereas prediction has to be validated.⁵⁶

mSMART.

The Mayo Stratification of Myeloma and Risk-Adapted Therapy (mSMART) criteria use a combination of FISH, plasma cell labeling index, and GEP as tools to identify 3 risk categories (standard risk [SR], IR, HR) for prognostication of patients with NDMM.⁵⁷ Patients can be stratified for different therapeutic approaches.⁴⁷ However, risk-adapted therapy has not been validated in prospective studies.

Consensus statement.

Translocations t(4;14), t(14;16), t(14;20), and del(17/17p) and any nonhyperdiploid karyotype are HR cytogenetics in NDMM regardless of treatment. Gain(1q) is associated with del(1p) carrying poor risk. Combinations of ≥3 CA confer ultra-HR with <2 years survival. Routine testing should include t(4;14) and del(17p). Clinical classifications may combine these lesions with ISS, serum LDH, or HR gene expression signatures. CA may differ in first and later relapse because of clonal evolution, which may influence the effect of salvage treatment.

Treatment options for high-risk disease

IMWG recommends using the combination of FISH, LDH, and ISS stage for risk stratification in NDMM.⁴⁷ Other features such as renal failure, plasma cell proliferative rate, and presence of extramedullary disease also contribute to risk.

GEP is emerging as a prognostic tool for risk stratification. New approaches to predict survival include analysis of microRNAs, custom capture mutation analysis, and evaluation of methylation and splicing patterns.

Here we address the treatment choices for patients with HR NDMM based on cytogenetic profile. Recently, 2 reviews addressed the issue of general treatment strategies for HR myeloma.^{58,59} This review covers treatment options for t(4;14) and/or del(17/17p).

Thalidomide.

Thalidomide does not overcome adverse impact of HR CA. In the UAMS trial for RRMM, del(13q) by karyotyping had a shorter survival with thalidomide.⁶⁰ Three trials studying thalidomide during induction in NDMM (MRC IX: CTD vs CTDa; HOVON50/GMMG-HD2: VAD vs TAD; GEM2005:TD) observed shorter OS in HR CA.⁶¹⁻⁶⁴ Thalidomide maintenance did not improve survival in HR CA in 3 trials, MRC IX (3-year OS 45% vs 69%), HOVON50 (3-year OS 17% vs 69%) trials and Total Therapy 2 (TT2) (5-year OS 56% vs 72%).^{15,18,25,61,62,65} In MRC IX, 3-year OS was worse in patients with HR-CA (45%).⁶⁶ In HOVON50/GMMG-HD2, first PFS was better with thalidomide treatment, but second PFS was significantly shorter, resulting in a reduced OS.⁶⁴ In TT2, presence of CA was associated with inferior survival, and a benefit with thalidomide was only observed in a subgroup of patients after 10 years.⁶⁷

Consensus statement.

Thalidomide does not abrogate the adverse effect of t(4;14), t(14;16), t(14;20), and del(17) or del(17p) and gain(1q) CA in TE patients. Conclusive data for elderly or frail patients are not available.

Bortezomib.

Several randomized trials have evaluated bortezomib for induction, consolidation, or maintenance treatment in cytogenetic subgroups. In IFM-2005-01, bortezomib/dexamethasone showed a superior response and OS compared with vincristin/doxorubicin/dexamethasone. This combination resulted in a better EFS and OS for patients with t(4;14), but did not improve outcome in del(17p) (4-year OS 50% vs 79%).⁶⁸ In HOVON65/GMMG-HD4, bortezomib-based induction and maintenance showed an improved outcome for patients with del(17p) (median PFS 26 vs 12 months; 3-year OS 69% vs 17%). At long-term follow-up, this advantage is still present. However, OS remains inferior to patients without del(17p) (3-year OS 85%). In patients with t(4;14), PFS was not better with bortezomib (25 vs 22 months), whereas OS was improved (3-year OS 69% vs 44%) compared with 85% in patients without t(4;14).²⁹ In the GEM 2005 trial, bortezomib/thalidomide/dexamethasone (VTD) followed by ASCT and maintenance did not improve OS in HR CA (3-year OS 60% vs 88%).⁶³ The GIMEMA group compared VTD with thalidomide/dexamethasone (TD) for induction and consolidation with double ASCT. In the subgroup of 25% with t(4;14), OS was 69% vs 37% in favor of VTD compared with 74% vs 63% without t(4;14) and/or del(17p).¹⁷ A meta-analysis of 4 randomized trials showed that the odds of posttransplantation complete response (CR) + near CR in bortezomib-treated patients were similar for HR (del(17p) + t(4;14)) and SR cytogenetics (2.44 vs 1.67, n.s.).¹⁶ These trials (1874 patients) showed that bortezomib plus ASCT was superior (PFS 41 vs 33 months) ($P < .0001$). In patients with HR FISH, this was 32 vs 22 months ($P < .0001$). PFS benefit was observed in patients with t(4;14) but lacking del(17p) (36 vs 24 months, $P = .001$) and in del(17p) lacking t(4;14) (27 vs 19 months, $P = .014$), but not in patients carrying both CA.⁶⁹ In TT3, OS was significantly shorter in patients with a HR profile (2-year OS 56% vs 88%) compared with SR GEP profile, with the

exception of low TP53 expression.⁷⁰ Addition of bortezomib improved OS compared with TT2 in LR MM.^{70,71}

Data in NTE patients are scarce. The VISTA trial combined melphalan/prednisone (MP) with MP + bortezomib (VMP). In patients treated with VMP, HR-CA did not influence outcome when compared with SR (OS 56% vs 71%).⁷² In a Pethema trial comparing VMP with bortezomib/thalidomide/prednisone (VPT) followed by maintenance with bortezomib/thalidomide vs bortezomib/prednisone, HR patients had shorter PFS than SR patients from the first (24 vs 33 months) and second randomization (17 vs 27 months) and shorter survival (3-year OS: 55% vs 77%).⁷³ The GIMEMA group compared VMP with VMP/thalidomide. In this bortezomib-dense treatment, HR vs SR patients had similar PFS.⁷⁴ The IFM group observed that across bortezomib regimens, no benefit was achieved in HR-CA NTE patients.⁷⁵

Consensus statement. Bortezomib partly overcomes the adverse effect of t(4;14) and possibly del(17p) on CR, PFS, and OS. There is no effect in t(4;14) combined with del(17p) in TE patients. In non-TE patients, VMP may partly restore PFS in HR cytogenetics.

Lenalidomide and pomalidomide. Experience with lenalidomide in first-line therapy for HR-CA patients is limited. In HR-CA, PFS with lenalidomide was inferior compared with SR patients (18 vs 26 months).⁷⁶ In the GIMEMA trial comparing high-dose melphalan with MPR, there was a trend for better PFS with lenalidomide maintenance in SR compared with HR-CA (HR 0.38 [0.24-0.62] vs 0.73 [0.37-1.42]). However, there was no effect on OS.⁷⁷ In the IFM 2005-02 trial, lenalidomide maintenance did not overcome the poor prognosis of t(4;14) (27 vs 24 months) and only partly of del(17p) (29 vs 14 months vs 42 months in all patients).⁷⁸ Convincing data for continuous lenalidomide in CA groups are lacking.^{79,80} Subgroup analysis of the FIRST trial in NDMM did not demonstrate a benefit of continuous lenalidomide in HR CA.⁸¹ In relapse MM, carfilzomib combined with lenalidomide and dexamethasone (K-RD) was effective across HR and SR patients (23 vs 29 months, $P = NS$), whereas RD showed less activity (13 vs 19 months, $P = .004$).⁸² Data of IFM did not show a benefit of RD in relapse/refractory multiple myeloma (RRMM) with del(13q) or t(4;14).⁸³ In the Eloquent 2 trial for RRMM, elotuzumab with RD (E-RD) improved outcome over RD in del(17p).⁸⁴ Recent data of the effect of pomalidomide with dexamethasone in patients with RRMM show that this combination does not abrogate overall adverse outcome in HR-CA, whereas OS may improve in del(17p).⁸⁵ In phase 2 trials, a response benefit of pomalidomide with dexamethasone was shown in patients with del(17p).⁸⁶

Consensus statement. Lenalidomide partly improves the adverse effect of t(4;14) and del(17p) on PFS, but not OS, in TE patients. In non-TE patients, there are no data suggesting that the drug may improve outcome with HR cytogenetics. Pomalidomide with dexamethasone showed promising results in RRMM with del(17p).

Combined proteasome inhibition and lenalidomide. Bortezomib combined with RD (VRD) in a phase 1/2 trial in 66 patients with NDMM showed 18-month PFS of 100% in 13 patients with del(17p) and/or t(4;14).⁸⁷ The EVOLUTION trial examined several schedules including VRD in NDMM. One-year PFS was similar in HR-CA (17% of all patients) and SR patients.⁸⁸ VRD in TE patients with NDMM had similar 3-year PFS (86%) in patients with >60% del(17p) or t(4;14) or del(13q) compared with all patients.⁸⁹

Carfilzomib monotherapy did not improve PFS/OS in t(4;14) or del(17p) in RRMM.⁹⁰ Carfilzomib combined with pomalidomide/dexamethasone had equivalent PFS and OS in HR vs SR RRMM.⁹¹ In the Aspire trial, in RRMM, KRd was superior to RD for PFS across cytogenetic risk groups, suggesting that this combination (partly) abrogates the negative impact of t(4;14) and del(17p).⁸² Similarly, in Tourmaline-MM1, ixazomib combined with RD showed identical PFS in patients with del(17p) or t(4;14) or no CA (20 vs 18 vs 20 m).⁹² More recently, carfilzomib combined with lenalidomide (KRd) or thalidomide (KTd) and dexamethasone in NDMM showed similar CR rate (>60%) and PFS between HR and SR patients.^{93,94}

Recently, favorable responses were observed with monoclonal antibodies against CD38 (daratumumab) or SLAMF7 (elotuzumab) combined with RD in RRMM across cytogenetic subgroups.⁹⁵

Consensus statement. Combining a proteasome inhibitor with lenalidomide and dexamethasone greatly reduces the adverse effect of t(4;14) and del(17p) on PFS in NDMM. Carfilzomib with lenalidomide and dexamethasone seems effective in patients with HR cytogenetics. However, with exception

of Aspire and Tourmaline, most data were obtained in nonrandomized studies and long-term follow-up has not been reported. The group advises treating NDMM patients with HR cytogenetics with the combination of a proteasome inhibitor with lenalidomide or pomalidomide and dexamethasone.

High-dose therapy and ASCT. In TE patients with NDMM, the hallmark of first-line treatment is high-dose therapy and ASCT combined with novel agents. This strategy has significantly improved PFS and OS. Therefore, it is difficult to address the role of HDT/ASCT for HR-CA. Few studies have investigated the effect of a second ASCT. In TT3, the addition of bortezomib to double ASCT improved outcome in patients with t(4;14), indicating that the effect of HDT/ASCT varies with induction and consolidation/maintenance.⁵ Similarly, addition of RVD for consolidation and maintenance after ASCT may improve PFS in HR MM.^{96,97} A meta-analysis of 4 European trials showed that double ASCT combined with bortezomib-based treatment partially abrogates poor PFS in patients carrying both t(4;14) and del(17p).⁶⁹

Consensus statement. HDT with ASCT is standard therapy for TE patients with NDMM. It contributes to improved outcome across prognostic groups. Double HDT/ASCT combined with bortezomib may improve PFS in patients with t(4;14) or del(17p), and in those with both abnormalities. Although results from stratified randomized trials are not yet available, HDT plus double ASCT is recommended for patients with HR cytogenetics. The results from clinical trials with bortezomib and thalidomide combinations with/without HDT + ASCT in HR cytogenetics are summarized in Tables 3 and 4.

Allogeneic stem cell transplantation. Allogeneic SCT has been proposed as a treatment of HR younger patients. Data on CA are scarce and partly based on classic karyotyping. In a trial of 73 NDMM patients, tandem auto-allo-transplantation yielded similar 5-year PFS (24% vs 30%) and OS (50% vs 54%) in patients without t(4;14) or del(17p).⁹⁸ The EBMT-NMAM2000 study showed better OS in patients treated with ASCT/RIC-allo or ASCT alone: 49% vs 36% at 96 months, respectively ($P = .030$). Unfortunately, convincing FISH data are lacking.⁹⁹ A retrospective analysis in 143 patients indicated that patients with del(13q) or t(4;14) or del(17p) or t(11;14) had similar 3-year PFS and OS as patients without abnormality.¹⁰⁰ A study of allo-SCT in 101 relapsed MM showed worse 4-year PFS (28 vs 43%) and OS (30 vs 49%) in 16 patients with del(17p), whereas in 16 patients with t(4;14) no impact was observed.¹⁰¹

Consensus statement. Allogeneic SCT or tandem auto-allo-SCT may improve PFS in patients with t(4;14) or del(17p). Results are better in an early stage of the disease. The novel treatments may challenge the role of allo-SCT and its use should be restricted to clinical trials.

Concluding remarks and future perspectives

Risk stratification in MM is important to predict survival and to define a treatment strategy. Cytogenetic abnormalities by FISH currently are clinically relevant prognostic factors in MM. The IMWG consensus panel on FISH advises to test for the presence of del(17p), t(4;14), and possibly t(14;16). An extended panel, which may be incorporated in clinical trials, includes t(11;14), t(14;20), gain(1q), del(1p), del(13q), and ploidy status. Bortezomib and lenalidomide may partially abrogate the adverse effect of del(17p). Bortezomib combined with iMDS may improve outcome in t(4;14). Double HDT/ASCT plus bortezomib may improve outcome in patients with both adverse CA. Application of these risk factors may be a first step toward precision medicine in patients with MM.

Acknowledgments

The following members of the International Myeloma Working Group participated in this study: Kenneth C. Anderson, Dana-Farber Cancer Institute, Boston, MA; Michel Attal, Purpan Hospital, Toulouse, France; Hervé Avet-Loiseau, University of Toulouse,

Toulouse, France; Joan Bladé, Hospital Clinic, IDIBAPS, University of Barcelona, Spain; Michele Cavo, University of Bologna, Bologna, Italy; Wee-Joo Chng, National University Health System, Singapore; Jo Caers, Centre Hospitalier Universitaire de Liège, Liège, Belgium; Niels van de Donk, VU University Medical Center Amsterdam, Amsterdam, The Netherlands; Dominik Dytfeld, Karol Marcinkowski University of Medical Sciences, Poznan, Poland; Michael O'Dwyer, National University of Ireland, Ireland; Hermann Einsele, Universitätsklinik Würzburg, Würzburg, Germany; Laurent Garderet, Hôpital Saint Antoine, Paris, France; Hartmut Goldschmidt, University Hospital Heidelberg and National Center for Tumor diseases Heidelberg, Heidelberg, Germany; Jens Hillengass, University of Heidelberg, Heidelberg, Germany; Shinsuke Ida, Nagoya City University Medical School, Nagoya, Japan; Robert A. Kyle, Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN; Sagar Lonial, Emory University Medical School, Atlanta, GA; Jin Lu, Peoples Hospital, Beijing University, Beijing, China; Marivi Mateos, University of Salamanca, Spain; Amitabha Mazumder, NYU Comprehensive Cancer Center, New York, NY; Philippe Moreau, University Hospital, Nantes, France; Amara Nouel, Hospital Rutz y Paez, Bolivar, Venezuela; Robert Orłowski, MD Anderson Cancer Center, Houston, TX; Antonio Palumbo, University of Torino, Torino, Italy; Anthony Reiman, Saint John Regional Hospital, Saint John, New Brunswick, Canada; Eloísa Riva Serra, Hospital de Clinicas, Montevideo, Uruguay; Javier de la Rubia, Hospital Universitario La Fe, Valencia, Spain; Kenneth R. Romeril,

Wellington Hospital, Wellington, New Zealand; Murielle Roussel, University of Toulouse, Toulouse, France; Jesús San Miguel, Clinica Universitaria de Navarra, CIMA, Pamplona, Spain; Orhan Sezer, Memorial Sisli Hastanesi, Istanbul, Turkey; Kazuyuki Shimizu, Tokai Central Hospital, Kakamigahara, Japan; David Siegel, Hackensack University Medical Center, Hackensack, NJ; Pieter Sonneveld, Erasmus MC, Rotterdam, The Netherlands; Evangelos Terpos, University of Athens School of Medicine, Athens, Greece; Saad Usmani, Levine Cancer Institute/Carolinas Healthcare System, Charlotte, NC; Sonja Zweegman, VU University Medical Center Amsterdam, Amsterdam, The Netherlands.

This study was supported by Amgen/Onyx, Celgene, Janssen, Karyopharm, and SkylineDx, and by honoraria from Amgen/Onyx, Celgene, Janssen, and Karyopharm.

Authorship

Contribution: P.S. wrote, revised, and approved the manuscript; and all authors revised and approved the manuscript.

Correspondence: Pieter Sonneveld, Department of Hematology, Rm Na822, Erasmus MC Cancer Institute, Erasmus MC, PO Box 2040, 3000 CA Rotterdam, The Netherlands; e-mail: p.sonneveld@erasmusmc.nl.

References

- Palumbo A, Anderson K. Multiple myeloma. *N Engl J Med*. 2011;364(11):1046-1060.
- Rajkumar SV, Dimopoulos MA, Palumbo A, et al. International Myeloma Working Group updated criteria for the diagnosis of multiple myeloma. *Lancet Oncol*. 2014;15(12):e538-e548.
- Morgan GJ, Walker BA, Davies FE. The genetic architecture of multiple myeloma. *Nat Rev Cancer*. 2012;12(5):335-348.
- Hervé AL, Florence M, Philippe M, et al. Molecular heterogeneity of multiple myeloma: pathogenesis, prognosis, and therapeutic implications. *J Clin Oncol*. 2011;29(14):1893-1897.
- Usmani SZ, Crowley J, Hoering A, et al. Improvement in long-term outcomes with successive Total Therapy trials for multiple myeloma: are patients now being cured? *Leukemia*. 2013;27(1):226-232.
- Ross FM, Avet-Loiseau H, Ameye G, et al. European Myeloma Network. Report from the European Myeloma Network on interphase FISH in multiple myeloma and related disorders. *Haematologica*. 2012;97(8):1272-1277.
- Bergsagel PL, Kuehl WM, Zhan F, Sawyer J, Barlogie B, Shaughnessy J Jr. Cyclin D dysregulation: an early and unifying pathogenic event in multiple myeloma. *Blood*. 2005;106(1):296-303.
- Chng WJ, Chung TH, Kumar S, et al. Gene signature combinations improve prognostic stratification of multiple myeloma patients. *Leukemia*. 2015.
- Munshi NC, Avet-Loiseau H. Genomics in multiple myeloma. *Clin Cancer Res*. 2011;17(6):1234-1242.
- Kuehl WM, Bergsagel PL. Multiple myeloma: evolving genetic events and host interactions. *Nat Rev Cancer*. 2002;2(3):175-187.
- Neben K, Jauch A, Bertsch U, et al. Combining information regarding chromosomal aberrations t(4;14) and del(17p13) with the International Staging System classification allows stratification of myeloma patients undergoing autologous stem cell transplantation. *Haematologica*. 2010;95(7):1150-1157.
- Boyd KD, Ross FM, Chiecchio L, et al; NCR I Haematology Oncology Studies Group. A novel prognostic model in myeloma based on co-segregating adverse FISH lesions and the ISS: analysis of patients treated in the MRC Myeloma IX trial. *Leukemia*. 2012;26(2):349-355.
- Fonseca R, Blood E, Rue M, et al. Clinical and biologic implications of recurrent genomic aberrations in myeloma. *Blood*. 2003;101(11):4569-4575.
- Gertz MA, Lacy MQ, Dispenzieri A, et al. Clinical implications of t(11;14)(q13;q32), t(4;14)(p16.3;q32), and -17p13 in myeloma patients treated with high-dose therapy. *Blood*. 2005;106(8):2837-2840.
- Sonneveld P, Schmidt-Wolf IG, van der Holt B, et al. Bortezomib induction and maintenance treatment in patients with newly diagnosed multiple myeloma: results of the randomized phase III HOVON-65/ GMMG-HD4 trial. *J Clin Oncol*. 2012;30(24):2946-2955.
- Sonneveld P, Goldschmidt H, Rosiñol L, et al. Bortezomib-based versus nonbortezomib-based induction treatment before autologous stem-cell transplantation in patients with previously untreated multiple myeloma: a meta-analysis of phase III randomized, controlled trials. *J Clin Oncol*. 2013;31(26):3279-3287.
- Cavo M, Tacchetti P, Patriarca F, et al; GIMEMA Italian Myeloma Network. Bortezomib with thalidomide plus dexamethasone compared with thalidomide plus dexamethasone as induction therapy before, and consolidation therapy after, double autologous stem-cell transplantation in newly diagnosed multiple myeloma: a randomised phase 3 study. *Lancet*. 2010;376(9758):2075-2085.
- Barlogie B, Pineda-Roman M, van Rhee F, et al. Thalidomide arm of Total Therapy 2 improves complete remission duration and survival in myeloma patients with metaphase cytogenetic abnormalities. *Blood*. 2008;112(8):3115-3121.
- Moreau P, Attal M, Garban F, et al; SAKK; IFM Group. Heterogeneity of t(4;14) in multiple myeloma. Long-term follow-up of 100 cases treated with tandem transplantation in IFM99 trials. *Leukemia*. 2007;21(9):2020-2024.
- Shaughnessy JD Jr, Zhan F, Burington BE, et al. A validated gene expression model of high-risk multiple myeloma is defined by deregulated expression of genes mapping to chromosome 1. *Blood*. 2007;109(6):2276-2284.
- Hebraud B, Magrangeas F, Cleynen A, et al. Role of additional chromosomal changes in the prognostic value of t(4;14) and del(17p) in multiple myeloma: the IFM experience. *Blood*. 2015;125(13):2095-2100.
- Narita T, Inagaki A, Kobayashi T, et al. t(14;16)-positive multiple myeloma shows negativity for CD56 expression and unfavorable outcome even in the era of novel drugs. *Blood Cancer J*. 2015;5:e285.
- Avet-Loiseau H, Malard F, Campion L, et al; Intergroupe Francophone du Myélome. Translocation t(14;16) and multiple myeloma: is it really an independent prognostic factor? *Blood*. 2011;117(6):2009-2011.
- Fonseca R, Blood EA, Oken MM, et al. Myeloma and the t(11;14)(q13;q32); evidence for a biologically defined unique subset of patients. *Blood*. 2002;99(10):3735-3741.
- Avet-Loiseau H, Attal M, Moreau P, et al. Genetic abnormalities and survival in multiple myeloma: the experience of the Intergroupe Francophone du Myélome. *Blood*. 2007;109(8):3489-3495.
- Pawlyn C, Melchor L, Murison A, et al. Coexistent hyperdiploidy does not abrogate poor prognosis in myeloma with adverse cytogenetics and may precede IGH translocations. *Blood*. 2015;125(5):831-840.

27. Walker BA, Leone PE, Chiecchio L, et al. A compendium of myeloma-associated chromosomal copy number abnormalities and their prognostic value. *Blood*. 2010;116(15):e56-e65.
28. Chiecchio L, Protheroe RK, Ibrahim AH, et al. Deletion of chromosome 13 detected by conventional cytogenetics is a critical prognostic factor in myeloma. *Leukemia*. 2006;20(9):1610-1617.
29. Neben K, Lokhorst HM, Jauch A, et al. Administration of bortezomib before and after autologous stem cell transplantation improves outcome in multiple myeloma patients with deletion 17p. *Blood*. 2012;119(4):940-948.
30. Teoh PJ, Chung TH, Sebastian S, et al. p53 haploinsufficiency and functional abnormalities in multiple myeloma. *Leukemia*. 2014;28(10):2066-2074.
31. Zhan F, Colla S, Wu X, et al. CKS1B, overexpressed in aggressive disease, regulates multiple myeloma growth and survival through SKP2- and p27Kip1-dependent and -independent mechanisms. *Blood*. 2007;109(11):4995-5001.
32. Boyd KD, Ross FM, Walker BA, et al; NCRI Haematology Oncology Studies Group. Mapping of chromosome 1p deletions in myeloma identifies FAM46C at 1p12 and CDKN2C at 1p32.3 as being genes in regions associated with adverse survival. *Clin Cancer Res*. 2011;17(24):7776-7784.
33. Chang H, Qi X, Jiang A, Xu W, Young T, Reece D. 1p21 deletions are strongly associated with 1q21 gains and are an independent adverse prognostic factor for the outcome of high-dose chemotherapy in patients with multiple myeloma. *Bone Marrow Transplant*. 2010;45(1):117-121.
34. Hebraud B, Leleu X, Lauwers-Cances V, et al. Deletion of the 1p32 region is a major independent prognostic factor in young patients with myeloma: the IFM experience on 1195 patients. *Leukemia*. 2014;28(3):675-679.
35. Walker BA, Boyle EM, Wardell CP, et al. Mutational Spectrum, Copy Number Changes, and Outcome: Results of a Sequencing Study of Patients With Newly Diagnosed Myeloma. *J Clin Oncol*. 2015;33(33):3911-3920.
36. Billecke L, Murga Penas EM, May AM, et al. Cytogenetics of extramedullary manifestations in multiple myeloma. *Br J Haematol*. 2013;161(1):87-94.
37. López-Anglada L, Gutiérrez NC, García JL, Mateos MV, Flores T, San Miguel JF. P53 deletion may drive the clinical evolution and treatment response in multiple myeloma. *Eur J Haematol*. 2010;84(4):359-361.
38. Avet-Loiseau H, Li C, Magrangeas F, et al. Prognostic significance of copy-number alterations in multiple myeloma. *J Clin Oncol*. 2009;27(27):4585-4590.
39. Morgan GJ, Gregory WM, Davies FE, et al; National Cancer Research Institute Haematological Oncology Clinical Studies Group. The role of maintenance thalidomide therapy in multiple myeloma: MRC Myeloma IX results and meta-analysis. *Blood*. 2012;119(1):7-15.
40. Kumar S, Fonseca R, Ketterling RP, et al. Trisomies in multiple myeloma: impact on survival in patients with high-risk cytogenetics. *Blood*. 2012;119(9):2100-2105.
41. Ludwig H, Durie BG, Bolejack V, et al. Myeloma in patients younger than age 50 years presents with more favorable features and shows better survival: an analysis of 10 549 patients from the International Myeloma Working Group. *Blood*. 2008;111(8):4039-4047.
42. Gay F, Larocca A, Wijermans P, et al. Complete response correlates with long-term progression-free and overall survival in elderly myeloma treated with novel agents: analysis of 1175 patients. *Blood*. 2011;117(11):3025-3031.
43. Fonseca R, Bergsagel PL, Drach J, et al; International Myeloma Working Group. International Myeloma Working Group molecular classification of multiple myeloma: spotlight review. *Leukemia*. 2009;23(12):2210-2221.
44. Stewart AK, Bergsagel PL, Greipp PR, et al. A practical guide to defining high-risk myeloma for clinical trials, patient counseling and choice of therapy. *Leukemia*. 2007;21(3):529-534.
45. Rajkumar SV, Gahrton G, Bergsagel PL. Approach to the treatment of multiple myeloma: a clash of philosophies. *Blood*. 2011;118(12):3205-3211.
46. Gutiérrez NC, Castellanos MV, Martín ML, et al; GEM/PETHEMA Spanish Group. Prognostic and biological implications of genetic abnormalities in multiple myeloma undergoing autologous stem cell transplantation: t(4;14) is the most relevant adverse prognostic factor, whereas RB deletion as a unique abnormality is not associated with adverse prognosis. *Leukemia*. 2007;21(1):143-150.
47. Chng WJ, Dispenzieri A, Chim CS, et al; International Myeloma Working Group. IMWG consensus on risk stratification in multiple myeloma. *Leukemia*. 2014;28(2):269-277.
48. Avet-Loiseau H, Durie BG, Cavo M, et al; International Myeloma Working Group. Combining fluorescent in situ hybridization data with ISS staging improves risk assessment in myeloma: an International Myeloma Working Group collaborative project. *Leukemia*. 2013;27(3):711-717.
49. Moreau P, Cavo M, Sonneveld P, et al. Combination of international scoring system 3, high lactate dehydrogenase, and t(4;14) and/or del(17p) identifies patients with multiple myeloma (MM) treated with front-line autologous stem-cell transplantation at high risk of early MM progression-related death. *J Clin Oncol*. 2014;32(20):2173-2180.
50. Palumbo A, Avet-Loiseau H, Oliva S, et al. Revised International Staging System for Multiple Myeloma: A Report From International Myeloma Working Group. *J Clin Oncol*. 2015;33(26):2863-2869.
51. Zhan F, Hardin J, Kordsmeier B, et al. Global gene expression profiling of multiple myeloma, monoclonal gammopathy of undetermined significance, and normal bone marrow plasma cells. *Blood*. 2002;99(5):1745-1757.
52. Shaughnessy JD Jr, Qu P, Usmani S, et al. Pharmacogenomics of bortezomib test-dosing identifies hyperexpression of proteasome genes, especially PSMD4, as novel high-risk feature in myeloma treated with Total Therapy 3. *Blood*. 2011;118(13):3512-3524.
53. Kuiper R, Broyl A, de Knecht Y, et al. A gene expression signature for high-risk multiple myeloma. *Leukemia*. 2012;26(11):2406-2413.
54. Kuiper R, van Duin M, van Vliet MH, et al. Prediction of high- and low-risk multiple myeloma based on gene expression and the International Staging System. *Blood*. 2015;126(17):1996-2004.
55. Decaux O, Lodé L, Magrangeas F, et al; Intergroupe Francophone du Myélome. Prediction of survival in multiple myeloma based on gene expression profiles reveals cell cycle and chromosomal instability signatures in high-risk patients and hyperdiploid signatures in low-risk patients: a study of the Intergroupe Francophone du Myélome. *J Clin Oncol*. 2008;26(29):4798-4805.
56. Amin SB, Yip WK, Minvielle S, et al. Gene expression profile alone is inadequate in predicting complete response in multiple myeloma. *Leukemia*. 2014;28(11):2229-2234.
57. Mikhael JR, Dingli D, Roy V, et al; Mayo Clinic. Management of newly diagnosed symptomatic multiple myeloma: updated Mayo Stratification of Myeloma and Risk-Adapted Therapy (mSMART) consensus guidelines 2013. *Mayo Clin Proc*. 2013;88(4):360-376.
58. Bergsagel PL, Mateos MV, Gutierrez NC, Rajkumar SV, San Miguel JF. Improving overall survival and overcoming adverse prognosis in the treatment of cytogenetically high-risk multiple myeloma. *Blood*. 2013;121(6):884-892.
59. Bianchi G, Richardson PG, Anderson KC. Best treatment strategies in high-risk multiple myeloma: navigating a gray area. *J Clin Oncol*. 2014;32(20):2125-2132.
60. Singhal S, Mehta J, Desikan R, et al. Antitumor activity of thalidomide in refractory multiple myeloma. *N Engl J Med*. 1999;341(21):1565-1571.
61. Morgan GJ, Davies FE, Gregory WM, et al; NCRI Haematological Oncology Study Group. Cyclophosphamide, thalidomide, and dexamethasone (CTD) as initial therapy for patients with multiple myeloma unsuitable for autologous transplantation. *Blood*. 2011;118(5):1231-1238.
62. Morgan GJ, Davies FE, Gregory WM, et al; National Cancer Research Institute Haematological Oncology Clinical Studies Group. Cyclophosphamide, thalidomide, and dexamethasone as induction therapy for newly diagnosed multiple myeloma patients destined for autologous stem-cell transplantation: MRC Myeloma IX randomized trial results. *Haematologica*. 2012;97(3):442-450.
63. Rosiñol L, Oriol A, Teruel AI, et al; Programa para el Estudio y la Terapéutica de las Hemopatías Malignas/Grupo Español de Mieloma (PETHEMA/GEM) group. Superiority of bortezomib, thalidomide, and dexamethasone (VTD) as induction pretransplantation therapy in multiple myeloma: a randomized phase 3 PETHEMA/GEM study. *Blood*. 2012;120(8):1589-1596.
64. Lokhorst HM, van der Holt B, Zweegman S, et al; Dutch-Belgian Hemato-Oncology Group (HOVON). A randomized phase 3 study on the effect of thalidomide combined with adriamycin, dexamethasone, and high-dose melphalan, followed by thalidomide maintenance in patients with multiple myeloma. *Blood*. 2010;115(6):1113-1120.
65. Attal M, Harousseau JL, Leyvraz S, et al; Intergroupe Francophone du Myélome (IFM). Maintenance therapy with thalidomide improves survival in patients with multiple myeloma. *Blood*. 2006;108(10):3289-3294.
66. Boyd KD, Ross FM, Tapper WJ, et al; NCRI Haematology Oncology Studies Group. The clinical impact and molecular biology of del(17p) in multiple myeloma treated with conventional or thalidomide-based therapy. *Genes Chromosomes Cancer*. 2011;50(10):765-774.
67. Barlogie B, Anaissie E, van Rhee F, et al. The Arkansas approach to therapy of patients with multiple myeloma. *Best Pract Res Clin Haematol*. 2007;20(4):761-781.
68. Avet-Loiseau H, Leleu X, Roussel M, et al. Bortezomib plus dexamethasone induction improves outcome of patients with t(4;14) myeloma but not outcome of patients with del(17p). *J Clin Oncol*. 2010;28(30):4630-4634.
69. Cavo M, Salwender H, Rosinol L, et al. Double vs single autologous stem cell transplantation after bortezomib-based induction regimens for multiple myeloma: an integrated analysis of

- patient-level data from phase European III studies [abstract]. *Blood*. 2013;122(21). Abstract 767.
70. Nair B, van Rhee F, Shaughnessy JD Jr, et al. Superior results of Total Therapy 3 (2003-33) in gene expression profiling-defined low-risk multiple myeloma confirmed in subsequent trial 2006-66 with VRD maintenance. *Blood*. 2010; 115(21):4168-4173.
 71. Pineda-Roman M, Zangari M, Haessler J, et al. Sustained complete remissions in multiple myeloma linked to bortezomib in total therapy 3: comparison with total therapy 2. *Br J Haematol*. 2008;140(6):625-634.
 72. San Miguel JF, Schlag R, Khuageva NK, et al; VISTA Trial Investigators. Bortezomib plus melphalan and prednisone for initial treatment of multiple myeloma. *N Engl J Med*. 2008;359(9): 906-917.
 73. Mateos MV, Gutiérrez NC, Martín-Ramos ML, et al. Outcome according to cytogenetic abnormalities and DNA ploidy in myeloma patients receiving short induction with weekly bortezomib followed by maintenance. *Blood*. 2011;118(17):4547-4553.
 74. Palumbo APBS, Rossi D, Berretta S, et al. A phase III study of VMPT versus VMP in newly diagnosed elderly myeloma patients. *J Clin Oncol*. 2009;27:8515a.
 75. Avet-Loiseau H, Hulin C, Campion L, et al. Chromosomal abnormalities are major prognostic factors in elderly patients with multiple myeloma: the intergroupe francophone du myélome experience. *J Clin Oncol*. 2013;31(22): 2806-2809.
 76. Kapoor P, Kumar S, Fonseca R, et al. Impact of risk stratification on outcome among patients with multiple myeloma receiving initial therapy with lenalidomide and dexamethasone. *Blood*. 2009;114(3):518-521.
 77. Palumbo A, Cavallo F, Gay F, et al. Autologous transplantation and maintenance therapy in multiple myeloma. *N Engl J Med*. 2014;371(10): 895-905.
 78. Attal M, Lauwers-Cances V, Marit G, et al; IFM Investigators. Lenalidomide maintenance after stem-cell transplantation for multiple myeloma. *N Engl J Med*. 2012;366(19):1782-1791.
 79. Palumbo A, Hajek R, Delforge M, et al; MM-015 Investigators. Continuous lenalidomide treatment for newly diagnosed multiple myeloma. *N Engl J Med*. 2012;366(19): 1759-1769.
 80. McCarthy PL, Owzar K, Hofmeister CC, et al. Lenalidomide after stem-cell transplantation for multiple myeloma. *N Engl J Med*. 2012;366(19): 1770-1781.
 81. Benboubker L, Dimopoulos MA, Dispenzieri A, et al; FIRST Trial Team. Lenalidomide and dexamethasone in transplant-ineligible patients with myeloma. *N Engl J Med*. 2014;371(10): 906-917.
 82. Stewart AK, Rajkumar SV, Dimopoulos MA, et al; ASPIRE Investigators. Carfilzomib, lenalidomide, and dexamethasone for relapsed multiple myeloma. *N Engl J Med*. 2015;372(2): 142-152.
 83. Avet-Loiseau H, Soulier J, Ferman J, et al; IFM and MAG groups. Impact of high-risk cytogenetics and prior therapy on outcomes in patients with advanced relapsed or refractory multiple myeloma treated with lenalidomide plus dexamethasone. *Leukemia*. 2010;24(3): 623-628.
 84. Lonial S, Dimopoulos M, Palumbo A, et al; ELOQUENT-2 Investigators. Elotuzumab Therapy for Relapsed or Refractory Multiple Myeloma. *N Engl J Med*. 2015;373(7):621-631.
 85. Dimopoulos MA, Weisel KC, Song KW, et al. Cytogenetics and long-term survival of patients with refractory or relapsed and refractory multiple myeloma treated with pomalidomide and low-dose dexamethasone. *Haematologica*. 2015; 100(10):1327-1333.
 86. Leleu X, Karlin L, Macro M, et al; Intergroupe Francophone du Myélome (IFM). Pomalidomide plus low-dose dexamethasone in multiple myeloma with deletion 17p and/or translocation (4;14): IFM 2010-02 trial results. *Blood*. 2015; 125(9):1411-1417.
 87. Richardson PG, Weller E, Lonial S, et al. Lenalidomide, bortezomib, and dexamethasone combination therapy in patients with newly diagnosed multiple myeloma. *Blood*. 2010; 116(5):679-686.
 88. Kumar S, Flinn I, Richardson PG, et al. Randomized, multicenter, phase 2 study (EVOLUTION) of combinations of bortezomib, dexamethasone, cyclophosphamide, and lenalidomide in previously untreated multiple myeloma. *Blood*. 2012;119(19):4375-4382.
 89. Roussel M, Lauwers-Cances V, Robillard N, et al. Front-line transplantation program with lenalidomide, bortezomib, and dexamethasone combination as induction and consolidation followed by lenalidomide maintenance in patients with multiple myeloma: a phase II study by the Intergroupe Francophone du Myélome. *J Clin Oncol*. 2014;32(25):2712-2717.
 90. Jakubowiak AJ, Siegel DS, Martin T, et al. Treatment outcomes in patients with relapsed and refractory multiple myeloma and high-risk cytogenetics receiving single-agent carfilzomib in the PX-171-003-A1 study. *Leukemia*. 2013; 27(12):2351-2356.
 91. Shah JJ, Stadtmauer EA, Abonour R, et al. Carfilzomib, pomalidomide, and dexamethasone for relapsed or refractory myeloma. *Blood*. 2015; 126(20):2284-2290.
 92. Moreau P, Masszi T, Grzasko N, et al. Ixazomib, an Investigational Oral Proteasome Inhibitor (PI), in Combination with Lenalidomide and Dexamethasone (IRd), Significantly Extends Progression-Free Survival (PFS) for Patients (Pts) with Relapsed and/or Refractory Multiple Myeloma (RRMM): The Phase 3 Tourmaline-MM1 Study (NCT01564537) [abstract]. *Blood*. 2015;126(23). Abstract 727.
 93. Sonneveld P, Asselbergs E, Zweegman S, et al. Phase 2 study of carfilzomib, thalidomide, and dexamethasone as induction/consolidation therapy for newly diagnosed multiple myeloma. *Blood*. 2015;125(3):449-456.
 94. Jakubowiak AJ, Dytfield D, Griffith KA, et al. A phase 1/2 study of carfilzomib in combination with lenalidomide and low-dose dexamethasone as a frontline treatment for multiple myeloma. *Blood*. 2012;120(9):1801-1809.
 95. van de Donk NW, Moreau P, Plesner T, et al. Clinical efficacy and management of monoclonal antibodies targeting CD38 and SLAMF7 in multiple myeloma. *Blood*. 2016; 127(6):681-695.
 96. Nooka AK, Kastiris E, Dimopoulos MA, Lonial S. Treatment options for relapsed and refractory multiple myeloma. *Blood*. 2015;125(20): 3085-3099.
 97. Nooka AK, Kaufman JL, Muppidi S, et al. Consolidation and maintenance therapy with lenalidomide, bortezomib and dexamethasone (RVD) in high-risk myeloma patients. *Leukemia*. 2014;28(3):690-693.
 98. Kröger N, Badbaran A, Zabelina T, et al. Impact of high-risk cytogenetics and achievement of molecular remission on long-term freedom from disease after autologous-allogeneic tandem transplantation in patients with multiple myeloma. *Biol Blood Marrow Transplant*. 2013; 19(3):398-404.
 99. Gahrton G, Iacobelli S, Björkstrand B, et al; EBMT Chronic Malignancies Working Party Plasma Cell Disorders Subcommittee. Autologous/reduced-intensity allogeneic stem cell transplantation vs autologous transplantation in multiple myeloma: long-term results of the EBMT-NMAM2000 study. *Blood*. 2013;121(25): 5055-5063.
 100. Roos-Weil D, Moreau P, Avet-Loiseau H, et al; Société Française de Greffe de Moelle et de Thérapie Cellulaire (SFGM-TC). Impact of genetic abnormalities after allogeneic stem cell transplantation in multiple myeloma: a report of the Société Française de Greffe de Moelle et de Thérapie Cellulaire. *Haematologica*. 2011;96(10): 1504-1511.
 101. Schilling G, Hansen T, Shimoni A, et al. Impact of genetic abnormalities on survival after allogeneic hematopoietic stem cell transplantation in multiple myeloma. *Leukemia*. 2008;22(6):1250-1255.
 102. Cavo M, Pantani L, Petrucci MT, et al; GIMEMA (Gruppo Italiano Malattie Ematologiche dell'Adulto) Italian Myeloma Network. Bortezomib-thalidomide-dexamethasone is superior to thalidomide-dexamethasone as consolidation therapy after autologous hematopoietic stem cell transplantation in patients with newly diagnosed multiple myeloma. *Blood*. 2012;120(1):9-19.
 103. Reece D, Song KW, Fu T, et al. Influence of cytogenetics in patients with relapsed or refractory multiple myeloma treated with lenalidomide plus dexamethasone: adverse effect of deletion 17p13. *Blood*. 2009;114(3): 522-525.
 104. Jacobus SJ, Kumar S, Uno H, et al. Impact of high-risk classification by FISH: an eastern cooperative oncology group (ECOG) study E4A03. *Br J Haematol*. 2011;155(3):340-348.



blood[®]

2016 127: 2955-2962

doi:10.1182/blood-2016-01-631200 originally published
online March 21, 2016

Treatment of multiple myeloma with high-risk cytogenetics: a consensus of the International Myeloma Working Group

Pieter Sonneveld, Hervé Avet-Loiseau, Sagar Lonial, Saad Usmani, David Siegel, Kenneth C. Anderson, Wee-Joo Chng, Philippe Moreau, Michel Attal, Robert A. Kyle, Jo Caers, Jens Hillengass, Jesús San Miguel, Niels W. C. J. van de Donk, Hermann Einsele, Joan Bladé, Brian G. M. Durie, Hartmut Goldschmidt, María-Victoria Mateos, Antonio Palumbo and Robert Orlowski

Updated information and services can be found at:

<http://www.bloodjournal.org/content/127/24/2955.full.html>

Articles on similar topics can be found in the following Blood collections

[Free Research Articles](#) (4070 articles)

[Lymphoid Neoplasia](#) (2378 articles)

[Multiple Myeloma](#) (343 articles)

[Perspectives](#) (188 articles)

Information about reproducing this article in parts or in its entirety may be found online at:

http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>