

## **Magma storage conditions and processes at Calbuco volcano (Central Southern Volcanic Zone, Chile)**

S.MONTALBANO<sup>1</sup>, NAMUR O.<sup>1,2</sup>, SCHIANO P.<sup>3</sup>, BOLLE O.<sup>1</sup>,  
VANDER AUWERA J.<sup>1</sup>

<sup>1</sup> Université de Liège, Département de géologie (Belgium),  
s.montalbano@ulg.ac.be

<sup>2</sup> Leibniz Universität Hannover (Germany)

<sup>3</sup> Laboratoire Magmas et Volcans, Clermont-Ferrand (France)

Despite the hazardous character of Calbuco stratovolcano, evidenced by the April 2015 eruption, its differentiation processes have not yet been investigated in detail. A series of 155 samples were collected in the different units of the volcano and analyzed for mineral and bulk rock composition. The macrocrysts are usually abundant (26-47 %) and comprise olivine (Fo<sub>56-88</sub>) usually with a peritectical reaction rim of orthopyroxene, plagioclase (An<sub>43-94</sub>), clinopyroxene (Mg#59-81), orthopyroxene (Mg#52-75), amphibole (Mg#59-75) locally with a dehydration rim due to decompression and titanomagnetite with rare exsolution lamellae of ilmenite.

In a TAS diagram, the samples range from basalts to andesites with a marked predominance of basaltic andesites. The differentiation trend is mainly calc-alkaline except for the basalts that plot in the tholeiitic field. In variation diagrams, CaO, MgO, FeO, Sc, Ni decrease with increasing SiO<sub>2</sub> whereas K<sub>2</sub>O, Rb, Zr increase. A few samples are significantly lower in MgO and FeO and higher in Al<sub>2</sub>O<sub>3</sub> and CaO (Low-MgO samples) than the samples of the main differentiation trend (High-MgO samples). This is interpreted as resulting from a higher degree of plagioclase accumulation in the former. Chondrite-normalized spiderdiagrams show enrichment in LILE and negative anomalies in Nb-Ta.

The combination of compatible and incompatible elements in appropriate equations shows that fractional crystallization is the dominant differentiation process. A three steps least square regression model indicates that the most evolved andesitic composition is reached after 43 % crystallization. Several geothermobarometers and the plagioclase hygrometer respectively constrain the depth of the main magma storage at 3-4 kb and the H<sub>2</sub>O content of 3-4 wt.%. This high water content is supported by the appearance of amphibole as a liquidus phase after about 39 % fractionation.