



## Research Article

# Testicular and Body Morphometric Traits of Mature Rams of Djallonke and Ouda Breeds Reared in North Benin

Koutinhoun G. Benoît<sup>1</sup>, Tougan Polycarpe Ulbad<sup>1,2\*</sup>, Boko K. Cyrille<sup>3</sup>, Douada I-H<sup>2</sup>, Zannou M. Serge<sup>4</sup>, Aholou A. B. Raoul<sup>1</sup>, Beckers Yves<sup>6</sup>, Hanzen Ch<sup>5</sup>, and Thewis André<sup>6</sup>

<sup>1</sup>Research Unit of Animal Husbandry and Meat Products Quality Control of Animal Production and Health Department; <sup>3</sup>Department of Animal Production and Health, Polytechnic School of Abomey-Calavi, 01 BP 2009, Cotonou, Republic of Benin, Benin; <sup>2</sup>Faculty of Agronomy (FA), University of Parakou, Republic of Benin; <sup>4</sup>Regional Excellence Centre of Avian Sciences of the University of Lome, BP 1515, Lome, Republic of Togo.

<sup>5</sup>Service de Thétiogénologie, Département clinique des animaux de production, Faculté de Médecine Vétérinaire Université de Liège; <sup>6</sup>Animal Sciences Unit, Gembloux Agro Bio Tech, University of Liege, ULg-Gx ABT-Passage des Deportés, 2-5030, Gembloux, Belgium

\*Corresponding author: ulcaless71@yahoo.fr (Dr. Tougan Polycarpe Ulbad)

**Article History:** Received: October 22, 2016      Revised: May 25, 2017      Accepted: May 26, 2016

### ABSTRACT

The study aims to compare the testicular and body morphometric traits of Djallonke and Ouda sheep breeds reared on natural pasture in Benin. Data on testicles and body morphometric parameters were collected on 70 mature rams at 12 months of age, including 45 Djallonke rams and 25 Ouda rams, from January 2015 to September 2016. These data were analyzed with SAS software (2006). It comes out from this study that Ouda rams were significantly heavier ( $P < 0.001$ ) than Djallonke rams at 12 months old. In the same way, the head length (19.29cm) and whither's height (57.39 cm) of Ouda Sheep were higher ( $P < 0.01$ ) than those of Djallonke sheep (17.92cm and 53.9 cm). Nevertheless, the scapulo-ischium length and chest circumference were similar ( $P > 0.05$ ) for both sheep breeds. However, the Djallonke sheep had presented the lower values of shoulder width and pelvis length (19.22 cm vs 22.61cm).

The scrotal length was significantly affected ( $P < 0.01$ ) by the sheep breed with the weakest value (13.73 cm) recorded within Djallonke sheep breed comparatively to Ouda rams (15.01 cm). In the same way, the scrotal circumference, paired testes weight, mean testes length, testes diameter and mean testes volume varied significantly ( $P < 0.001$ ) according to the sheep breed with the highest values observed in Ouda sheep.

**Key words:** Breed, Sheep, Testis, Morphometric traits, Variability

### INTRODUCTION

Sheep production is a widespread activity in Benin, as in most countries in sub-Saharan Africa (Gbangboché *et al.*, 2005a, SWAC-OECD / ECOWAS 2008; Babatunde *et al.*, 2010). As ruminants, they play a very vital role in the livelihood of rural populations as sales of the animals and their products help to stabilize household income. Djallonke and Ouda breeds are the main sheep breeds met in Benin with a predominance of Djallonke breed (Gbangboché *et al.*, 2004) because of its perfect adaptation to the local climatic conditions and its resistance (Mawuena 1987; Gbangboché *et al.*, 2005a, Gbangboché *et al.*, 2005b).

The West African Dwarf sheep of Djallonke breed is a hairy sheep breed found all over West and Central

Africa of 14° south latitude. The characteristics of WAD sheep have been described by several authors (Rombaut, 1980; Pagot, 1985; Larrat, 1989; Fournier, 2006; Gbangboché *et al.*, 2005). It is a compact breed with a small mature size and short horizontal lop ears. Coat colour varies from spotted black and white to solid black or white. Some have tan or brown coat colour and black bellies. Rams are horned and females usually polled. WAD sheep are capable of limiting parasite multiplication and remain productive in tsetse-infested areas where other breeds can't survive without treatment (Ayuk *et al.*, 2014). However, West African dwarf sheep have low productivity (Yapi, 1994) and weak reproduction and growth performance (Adjibodè, 2012) varying with latitude from north to the south of Benin (Gbangboché *et al.*, 2005b; Youssao *et al.*, 2008).

**Cite This Article as:** Benoît KG, TP Ulbad, BK Cyrille, IH Douada, ZM Serge, C Hanzen and T André, 2017. Testicular and body morphometric traits of mature rams of Djallonke and Ouda breeds reared in North Benin. *Inter J Vet Sci*, 6(2): 108-113. www.ijvets.com (©2017 IJVS. All rights reserved)

Found in northern Nigeria, southern Niger, Northern Benin, central TC had, northern Cameroon and western Sudan, the Ouda sheep breed is one of the hair sheep breeds of the Sahel type (Mason, 1996). It is a meat breed with distinctive markings. The front half of the body is black or brown and the back half is white. The rams of the Ouda are horned and the ewes are usually polled. When compared to the West African Dwarf, the main difference is that the Sahel-type sheep are taller, heavier, of poor mutton conformation and rams do not have a throat ruff or mane. The Sahel-type are usually white, white and brown, or white and black with lop ears. The males display a long twisting pattern to the horns and the females are usually polled. It is adapted to extensive grazing and survival under hot and dry environment. In general, the Ouda sheep inhabit the semi-arid mono-modal rainfall lowlands and adjoining arid areas of southern Niger, northern Benin, northern Nigeria, central Chad, western Sudan and parts of Cameroon. It does not survive well outside its ecological zone.

Due to their relative simplicity compared to the rearing of large animals (Vallerand and Blanckaertr, 1975; London and Weninger, 1996), sheep rearing in Benin increase over the years with a national herd increasing from 690,000 heads in 2003 to 860,000 heads in 2013 (County Stat, 2016).

However, the productivity of those sheep breed in Benin is low (Gbangboché *et al.*, 2005b; Youssao *et al.*, 2008), amongst others, due to a low fertility rate in the breeding herds. Thus, to keep up with the increasing demand of meat production and the productivity of sheep, there is a need for sustainable improvement strategies. Improvement of sheep reproductive performance requires effective actions on its various components, with prolificacy being one of the most important (Yakubu and Musa-Azara, 2013) such as body and gonads morphometric traits.

Morphometric traits are continuous parameters describing aspects of animal body conformation (Riva *et al.*, 2004; Cervantes *et al.*, 2009) and are an essential component of breed characterization (Gizaw *et al.*, 2007). Morphological parameters such as chest circumference, wither's height and scapulo-ischium length can be used for rapid selection of large size animals for constitution of elite herds (Dossa *et al.*, 2007). Variations in morphometric traits between different sheep populations can provide a soundness basis for livestock management.

Few studies (Gbangboché *et al.*, 2004; Akouedegni *et al.*, 2013; Awohuedji *et al.*, 2013) are carried on the morphometric characterization of sheep breeds reared in Benin. Testicular measurements and live weight were reported to generally indicate the production of viable spermatozoa by the male (Agga *et al.*, 2011). The morpho-biometrical analysis of testicular development is of great importance since it is significantly correlated with reproductive activity (Emsen, 2005).

The aim of this study was to compare the body and testicular morphometric traits of Djallonke and Ouda sheep breeds reared under traditional sheep breeding in Benin.

## MATERIALS AND METHODS

### Study area

The study was thus conducted in the department of Borgou in the North of Benin. Situated between the

latitudes of 6°20' and 12°30' north and between the longitudes of 1°30' and 3°45' East, the republic of Benin covers an area of 113440 km<sup>2</sup> with a population of 10448647 inhabitants (INSAE, 2015) and a density of 60 hab/km<sup>2</sup>.

The department Borgou exhibits climatic conditions of Sudan type, characterized by only one rainy season (from April to October) and one dry season (November to March). Average rainfall varies between 900 and 1300 mm per year while the average annual temperature is 26°C with a maximum of 32°C in March and a minimum of 23°C from December to January. The relative humidity varies between 30 and 70%. Vegetation of Borgou department is a diversified savannah where tree density decreases towards the North.

### Data collection

Data on testicles and body morphometric parameters were then collected on 45 Djallonke rams and 25 Ouda rams of 12 months old, from from January 2015 to September 2016. These animals were all raised in a traditional system. Feeding was mainly based on natural pasture. The animals were put on pasture at about 7.30 a.m. and returned to the barns in the afternoon. They were then fed *ad libitum* a supplementary diet consisting of crop residues according to the traditional system. The sheep were treated for ectoparasites, drenched once every three months and given other veterinary attention when the need appeared.

Reproductive tracts of those 70 matured rams of 12 months old were obtained after slaughter within the department of Borgou in the North of Benin. The reproductive tracts were then immediately brought to the laboratory covered in ice and were processed on the same day.

The material used for data collection was composed of a data file for recording the testicular and body morphometric traits and usual morphometric traits recording materials.

The body weight of the rams was recorded using a scale of 40 kg of capability and 20g of accuracy. Scrotal circumference was measured using a tape at the broadest part of the scrotum. Shoulder width was determined with the aid of a tape measure, as the horizontal distance between the processes on the left shoulder and those of the right shoulder blade. Chest circumference was measured by using a measuring tape around the chest, just behind the front legs; body length was measured from the sternum to the aitch bone and hip or pelvis width was measured using a plastic measuring tape, while height at wither was measured vertically from thoracic vertebrae to the ground using a metal ruler.

The epididymis was carefully excised from the testis along the physiological joints. The testes and epididymis was separated free of adhering connective tissues and fats before the records of morphometric parameters. The following parameters were taking: Testicular length, Testicular diameter, Testicular volume, Testicular weight, Epididymal weight.

Testicles length was measured with the use of flexible tape in cm; testicles diameter was taken with the use of Vernier caliper; testicle volume was measured by the use of water displacement technique according to Alexandrou

(2001). Testicular and Epididymal weights were recorded in grams with the use of digital weighing balance. Also, the testicular density was obtained by dividing the testicular weight by the testicular volume as the following formula:

$$\text{Testicular density (g.cm}^{-1}\text{)} = \frac{\text{Testes weight (g)}}{\text{Testes volume (cm}^3\text{)}}$$

### Statistical analysis

Data collected were analyzed for the effect of breed (Ouda and Djallonke) using the software Statistical Analysis System (SAS, 2006). Student t-test and the one way analysis of variance (ANOVA) analysis were done. The mathematical expression of this model is as follows:

$$Y_{ij} = \mu + E_i + e_{ij}$$

Where

$Y_{ij}$ : the morphometric parameter of the animal j, of the breed i (Ouda or Djallonke);

$\mu$ : overall mean;

$E_i$ : fixed effect of breed i (Ouda or Djallonke);

$e_{ij}$ : Effect of random residual average performance of the animal j, of the breed i (Ouda or Djallonke).

## RESULTS

### Variability of body morphometric parameters in Djallonke and Ouda Sheep reared in Benin

The table 1 show the variability of body morphometric parameters in Djallonke and Ouda Sheep reared in Benin. It appears that Ouda rams were significantly heavier ( $P < 0.001$ ) than Djallonke rams at 12 months old. In the same way, the head length (19.29cm) and whither's height (57.39cm) of Ouda Sheep were higher ( $P < 0.01$ ) than those of Djallonke sheep (17.92cm

and 53.9cm). Nevertheless, the scapulo-ischium length and chest circumference were similar ( $P > 0.05$ ) for both sheep breeds. However, the Djallonke sheep had presented the lower values of shoulder width (14.66 cm vs 15.632cm) and pelvis length (19.22 cm vs 22.61cm).

### Variability testicular measurements in Djallonke and Ouda Sheep reared in Benin

The variability of testicular measurements parameters in Djallonke and Ouda Sheep reared in Benin is given in table 2. The scrotal length was significantly affected ( $P < 0.01$ ) by the sheep breed with the weakest value (13.73cm) recorded within Djallonke sheep breed comparatively to Ouda rams (15.01cm). In the same way, the scrotal circumference, paired testes weight, mean testes length, testes diameter and mean testes volume varied significantly ( $P < 0.001$ ) according to the sheep breed with the highest values observed in Ouda sheep. The paired epididymal weight and mean epididymal length of Ouda rams (29.21g and 21.12cm respectively) were significantly higher ( $P < 0.001$ ) than those of Djallonke breed (25.79g and 18.64cm respectively). Similarly, the mean epididymal volume of Djallonke sheep was 14.76ml to 16.62ml for Ouda breed ( $P < 0.01$ ). However, the mean testes density and the mean epididymal density of the both sheep breeds didn't vary significantly according to the breed ( $P > 0.05$ ).

## DISCUSSION

### Variability of body morphometric and testicular traits in Djallonke and Ouda Sheep breeding in Benin

Our results on measured body and testicular traits showed that the West African Dwarf (WAD) Sheep of Djallonke breed is a small format sheep breed comparatively to Ouda breed which is the heavier. Similar result was presented by Otoikhian (2008) and Okpeku *et*

**Table 1:** Variability of body morphometric parameters in Djallonke and Ouda Sheep breeding in Benin

Variables	Djallonke		Ouda		Breed effect
	Mean	SE	Mean	SE	
Live Weight (kg)	20.18	0.70	25.18	0.51	***
Head length (cm)	17.92	0.33	19.29	0.34	**
Whither's height (cm)	53.90	0.83	57.39	0.94	**
Scapulo-Ischium length (cm)	60.12	0.73	61.53	0.58	NS
Chest Circumference (cm)	70.82	1.02	71.70	0.86	NS
Shoulder or Back width (cm)	14.66	0.28	15.63	0.24	*
Pelvis length (cm)	19.22	0.58	22.61	0.47	***

SE: Standard Error, NS:  $P > 0.05$ ; \*:  $P < 0.5$ ; \*\*:  $P < 0.01$ ; \*\*\*:  $P < 0.001$ .

**Table 2:** Variability testicular measurements in Djallonke and Ouda Sheep breeding in Benin

Variables	Djallonke		Ouda		Breed effect
	Mean	SE	Mean	SE	
Scrotal length (cm)	13.728	0.266	15.012	0.346	**
Scrotal circumference (cm)	21.41	0.607	24.514	0.447	***
Paired testes weight (g)	279.13	6.53	318.36	5.4	***
Mean testes length (cm)	6.303	0.148	7.143	0.134	***
Testes diameter (cm)	4.395	0.102	4.9857	0.0933	***
Mean testes volume (ml)	133.28	3.14	151.27	2.82	***
Mean testes density (g/cm <sup>3</sup> )	1.0492	0.00598	1.0471	0.00485	NS
Paired epididymal weight (g)	25.785	0.602	29.214	0.508	***
Mean epididymal length (cm)	18.644	0.435	21.121	0.367	***
Mean epididymal volume (ml)	14.767	0.366	16.621	0.445	**
Mean epididymal density (g/cm <sup>3</sup> )	0.8782	0.0153	0.8857	0.0143	NS

SE: Standard Error, NS:  $P > 0.05$ ; \*\*:  $P < 0.01$ ; \*\*\*:  $P < 0.001$ .

*al.* (2011). According to these authors, the Ouda, Balami and Yankasa sheep breeds are heavier than the WAD sheep. Several authors concluded, that the genotype or the breeds of sheep reared under comparable conditions have frequently different age-type weight and average daily gains (Gbangboche 2005a; Doko Alou *et al.*, 2013). These authors linked it to several factors including the genetic type and natural selection (Lebbie and Ramsay, 1999; Gbangboche, 2005b; Youssao, 2008). This reality is highlighted by the results of our study which show that Ouda rams were significantly heavier than Djallonke Dwarf rams. This is in agreement with the report of Osinowo *et al.* (1989) and Otoikhian (2008). It was also observed that the variation in phenotypic traits of the studied sheep breeds found in Benin is relatively high. This is a good potential to explore for genetic improvement programmes for Benin native sheep breeds.

Poivey *et al.* (1982) reported high heritability for weight at three months of age and suggest taking this into account in the selection of dwarf sheep. It is therefore proven that the crossbreeds Sahelian x West African dwarf sheep express best growth performances than pure Djallonke breed (Amégé, 1984) and could be perceived as an alternative of improvement of the profitability of sheep rearing (Gbangboché *et al.*, 2002). Thus, the crossing between Djallonke sheep and Ouda sheep breeds can improve the growth performances of Djallonke sheep and preserve their resistance to disease.

The weights obtained at the 12<sup>th</sup> month old in Djallonke and Ouda rams are lower than the value of 30kg obtained by Amégé (2004) for Vogan sheep at 7 months old and the live weight of 37.3kg reported for Lohi sheep of 9 months old by Lashari and Tasawar (2010). These confirm that the growth performances of the sheep breeds reared in Benin have relatively very low growth performance and need improvement program, since body size is generally important parameters used in breeding soundness evaluation. Knowing the body weight of an animal is important for a number of reasons such as breeding, correct feeding, health matters, growth as well as classification.

In the same way, the head length and wither's height of Ouda Sheep were significantly higher than those of Djallonke sheep. However, the scapulo-ischium length and chest circumference were similar in the both sheep breeds in our study. The Djallonke sheep had also presented the lower values of shoulder width and pelvis length.

The variation of morphometric measurements observed herein according to the ecotype with the highest values of all measured parameters found in sheep of Ouda breed is in accordance with the report of Salako and Ngere (2002). Similar result was also reported by Traore *et al.* (2008) between the Sahelian and Sudan- Sahelian (Mossi) sheep of Burkina-Faso where Sahelian sheep had longer tail than Mossi sheep. According to FAO (2005), the characterization of body measurements of the Djallonke sheep presents great variability according to the latitudes and mediums of breeding.

#### **Variability testicular measurements in Djallonke and Ouda Sheep reared in Benin**

The testicular morphometric parameters in Djallonke and Ouda Sheep are given in table 2. It come out from this

table that, the scrotal length was significantly affected by the sheep breed with the weakest value recorded within Djallonke sheep breed comparatively to Ouda rams. In the same way, the scrotal circumference, paired testes weight, mean testes length, testes diameter and mean testes volume varied significantly according to the sheep breed with the highest values observed in Ouda sheep. The paired epididymal weight and mean epididymal length of Ouda rams were significantly higher than those of Djallonke breed. Similarly, the mean epididymal volume of Djallonke sheep was 14.76 ml to 16.62 ml for Ouda breed. This finding confirms the result of Ibrahim *et al.* (2012) who observed that the scrotal circumference of Uda was significantly higher than of the Balami and Yankasa, but there had no significant difference between Balami and Yankasa breeds. Similar reports of differences between breeds were earlier reported in goats (Raji *et al.*, 2008) and cattle (Abassi, 2011; Addass, 2011). According to these authors, these differences could be due to the effect of genotype or breed. Nevertheless, the means of scrotal circumference of Ouda breed obtained herein was higher than the value of  $21.50 \pm 0.61$ , reported by Ahemen and Bitto (2007) for West African dwarf rams but similar with the value recorded in Djallonke rams. Scrotal circumference is measured as it gives a good indication of rams breeding ability. Schoenian (2011) reported that ram lamb with scrotal circumference of less than 30 cm and adult rams with scrotal circumference of less than 32 cm should probably not be used for breeding. Similarly, Söderquist and Hultén (2006) reported that for mature rams (17 - 54 months old), the mean scrotal circumference was  $34.4 \pm 2$  cm in Gotlandie breed, and  $34.5 \pm 0.9$  cm for Dorper ram by Besta.

The differences found in testes size could be due to the effect of breed. Similar reports of differences between breeds were earlier reported in goats (Raji *et al.*, 2008) and cattle (Abassi, 2011; Addass, 2011).

Testes weight, a soundness index of semen producing ability had been found to depend on the breed in the current study. The results observed herein are similar to those found in other breeds of sheep within the tropics and sub-tropics, but significantly higher than the values varying between 134.48 g and 154 g as reported for WAD rams and Kajli rams in Nigeria and Pakistan by Ahemen and Bitto (2007) and Siddiqui *et al.* (2005) respively.

Moreover, it was reported that males with larger testes tend to sire daughters that reach puberty at an earlier age (Söderquist and Hultén, 2006). Brito *et al.* (2004) have reported that heavier testes produce more spermatozoa than the smaller testes in breeding animals. The significant higher testes weight of sheep Ouda found in our study would mean that those testes could contain more seminiferous tubule, interstitial endocrine cells and possibly more spermatozoa than the Djallonke breed. The variation in the testes volume for the different breed observed in this study agrees with the report of Ibrahim (2012) in different strains of chicken.

#### **Conclusion**

The study revealed that chest circumference, back width, pelvis length, paired testes weight, mean testes length, testes diameter, mean testes volume, mean testes density, paired epididymal weight, mean epididymal

length, mean epididymal volume and the mean epididymal density depend on the sheep breed with the best testicles size recorded in Ouda sheep. Further study must be carried out on the variability of semen quality of those sheep breeds reared in Benin.

#### Acknowledgement

The authors are very much grateful to the traditional sheep breeders and the lab of the National Agro-pastoral High School of the University of Parakou for giving all types of support in conducting this study.

#### REFERENCES

- Abbasi MA and F Ghafouri-Kesbi, 2011. Genetic (co) variance components for body weight and body measurements in Makooei sheep. *Asian-Aust J Anim Sci*, 24: 739-743.
- Addass PA, 2011. Genotype and seasonal variations in testes and paired epididymal sperm production among indigenous bull cattle in Mubi Adamawa State, Nigeria. *Agric Biol J N Am*, 2: 19-22.
- Adjibode AG, 2012. Facteurs non génétiques influençant les paramètres zootechniques de la chèvre naine en zone soudanienne du Bénin. Mémoire présenté en vue de l'obtention du diplôme de master en production et santé animales, EPAC/UAC, 91.
- Agga GE, U Udala, F Regassa and A Wudie, 2011. Body measurements of bucks of three goat breeds in Ethiopia and their correlation to breed, age and testicular measurements. *Small Rumin Res*, 95: 133-138.
- Ahemen T, and II Bitto, 2007. Sperm production rate, Gonadal and extragonadal sperm reserves of the West African Dwarf rams in Makurdi. *Proc. of the 32nd Annu Conf of Nig Soc for Anim Prod*, pp: 99-101.
- Akouèdègni CG, IG Tossa, E Ahoussi, and MS Hounzangbé-Adoté, 2013. Effects of the fresh leaves of *Spondias mombin* L. on milk production of West African Dwarf (WAD) ewes and their lamb's growth performance. *Global J Res Med Plants & Indigen Med*, 2: 126-134.
- Alexandrou A, 2001. Principles of fluid mechanics (1 edition). New Jersey. Prentice Hall Inc.
- Amégée Y, 1984. Le mouton de Vogon (croisé Djallonké x Sahélien) au Togo III. Performances d'engraissement et rendement des carcasses. *Revue Elev Méd vét Pays Trop*, 37: 97-106.
- Awohouedji DY, S Babatounde, JG Adoukpe, M Houinato and S Hounzangbe-Adote, 2013. Supplementing *Panicum maximum* with two medicinal forages in the diet of Djallonké sheep at the Benin national sheep center. *Sci J Anim Sci*, 2: 284-295.
- Ayuk A, E Iyayi, B Okon and J Ayuk, 2014. Growth Performance of West African Dwarf (WAD) Sheep Fed Biodegraded *Enterolobium cyclocarpum* Based Diets. *Agricultural Sciences*, 5, 710-715. doi: 10.4236/as.2014.58074.
- Babatounde S, Oumorou M, Tchabi IV, Lecomte T, Houinato M, Adandedjan C, 2010. Ingestion volontaire et préférences alimentaires chez des moutons Djallonké nourris avec des graminées et des légumineuses fourragères tropicales cultivées au Bénin. *Int J Biol Chem Sci*, 4: 1030-1043.
- Bitto II, and GN Egbunike, 2006. Seasonal variations in the morphometric characteristics of the pubertal West African Dwarf Bucks in its Native Tropical Environment. *Int J Morphol*, 24: 637-42.
- Cervantes I, R Baumung, A Molina, T Druml, JP Gutierrez, J Solkner, and M Valera, 2009. Size and shape analysis of morphofunctional traits in the Spanish Arab horse. *Livestock Sci*, 125: 43-49. doi: 10.1016/j.livsci.2009.03.006. [Cross Ref]
- Country Stat, 2016. Répartition des effectifs d'animaux vivants. <http://www.countrystat.org/home.aspx?c=BENandta=053SPD135andtr=21> consulté le 06/05/2016.
- CountryStat/Benin, 2013. Base de données statistiques, consulté à l'adresse, <http://countrystat.org/ben> ou <http://www.fao.org/economic/ess/countrystat/en/>. Consulté le 06/05/2016.
- Doko AS, S Farougou and FCC Hountondji, 2013. Impact of prophylactic measures and the use of local food resources on the viability and growth of pre-weaning lambs in Djougou, in the northern region of Benin. *J Anim Plant Sci*, 19: 2933-2940.
- Dossa LH, C Wollny, and M Gaulty, 2007. Spatial variation in goat populations from Benin as revealed by multivariate analysis of morphological traits. *Small Rumin Res* 73: 150-159. doi: 10.1016/j.smallrumres.2007.01.003. [Cross Ref]
- Emsen E, 2005. Testicular development and body weight gain from birth to 1 year of age of Awassi and Redkaraman sheep and their reciprocal crosses. *Small Rumin Res*, 29: 79-82.
- FAO, 2005. Global Livestock Production and Health Atlas. Food and Agriculture Organisation, Rome. <http://www.fao.org/ag/aga/glipha/index.jsp> (accessed March, 2005).
- Fournier A, 2006. Elevage des moutons. Edition Artemis, Slovaquie, 94p.
- Gbangboché AB, FA Abiola, JP Laporte, S Salifou, PL Leroy, 2002. Amélioration des ovins dans l'Ouémé et le Plateau en République du Bénin. Enjeux de croisement des ovins Djallonké avec les moutons du Sahel. *Tropicicultura*, 20: 70-75.
- Gbangboché AB, FA Abiola, C Michaux, and PL Leroy, 2005a. Paramètres génétiques et non-génétiques des caractères de croissance du mouton Djallonké au Bénin. *Renc. Rech. Ruminants*, 12.
- Gbangboché AB, JL Hornich, M Adamou-N'diaye, AP Etorh, F Fanir, FA Abiola, and PL Leroy, 2005b. Caractérisation et maîtrise des paramètres de la reproduction et de la croissance des ovins Djallonké (*Ovis aries*). *Ann Méd Vét*, 149: 170-182.
- Gbangboché AB, SM Hounzangbé-Adoté, SY Doko, F Farnir, J Detilleux, and PL Leroy, 2004. Production des ovins Djallonké en station dans la zone guinéenne au Bénin: 1. Performances de reproduction et influence des facteurs non génétiques. *RASPA* 2: 49-55.
- Gbangboché AB, FA Abiola, JP Laporte, S Salifou et PL Leroy, 2002. Amélioration des ovins dans l'Ouémé et le Plateau en République du Bénin. Enjeux de

- croisement des ovins Djallonké avec les moutons du Sahel. *Tropicultura*, 20: 70-75.
- Gizaw S, JAM van Arendonk, H Komen, JJ Windig, O Hanotte, 2007. Population structure, genetic variation and morphological diversity in indigenous sheep of Ethiopia. *Anim Genet*; 38:621-628. doi: 10.1111/j.1365-2052.2007.01659.x.
- Ibrahim AA, 2012. Assessments of the semen quality, testicular development and carcass characteristics of four strains of cockerels in a semi-arid environment of Kano State, Nigeria. MSc. Thesis. University of Maiduguri.
- Larrat R, 1989. Manuel vétérinaire des agents techniques de l'élevage tropical. Collection manuels et précis d'élevage, Ministère de la coopération & Développement, IEMVT, n°5, 2è édition, 533p.
- Lashari MH, and Z Tasawar, 2010. Genetic Potentials of local breed of sheep habitating around dera ghazi khan, Pakistan. *Sarhad J Agric*, 26: 251-258.
- Lebbie SHB and K Ramsay, 1999. A perspective on conservation and management of small ruminant genetic resources in the sub-Saharan Africa. *Small Rumin Res*, 34: 231-247.
- London JC, and JH Weniger, 1996. Investigation into traditionally managed Djallonké-sheep production in humid and subhumid zones of Asante, Ghana. V. Productivity indices. *J Anim Breed Genet*, 113: 483-492.
- Mason IL, 1996. A World Dictionary of Livestock Breeds, Types and Varieties. Fourth Edition. CAB International, 273 pp.
- Mawuena K, 1987. Haut degré de tolérance à la trypanosomose des moutons et des chèvres de race Naine Djallonké des régions sud-guinéennes du Togo: comparaison avec les bovins trypanotolérants. *Rev Elev Méd Vét Pays Trop*, 40: 55-58.
- Okpeku M, A Yakubu, SO Peters, MO Ozoje, CON Ikeobi, OA Adebambo and IG Imumorin, 2011. Application of multivariate principal component analysis to morphological characterization of indigenous goats in Southern Nigeria. *Acta Agric Slov*, 98: 101-109.
- Osinowo EO, SM Dennis, DIK Osori, ECT Molokwu, 1977. Scrotal and Testicular Dimensions in some West African Bulls. *Niger J Anim Prod* 4: 31-36.
- Otoikhian CSO, AM Otoikhian, OP Akporhwarho and C Isidahoman, 2008. Correlation of body weight and some body measurement parameters in Quda sheep under extensive management system. *Afr J Gen Agric*, 4: 129-133.
- Pagot J, 1985. L'élevage en pays tropicaux. Editions Maisonneuve et Larose (Paris), 526p.
- Poivey JP, E Landais, and Y Berger, 1982. Etude et amélioration génétique de la croissance des Djallonké. Résultats obtenus au Centre de Recherches Zootechniques de Bouaké (Côte-d'Ivoire). *Revue Elev Méd Vét Pays Trop*, 35: 421-433.
- Raji AO, JU Igwebuike, and J Aliyu, 2008. Testicular biometry and its relationship with body weight of indigenous goats in a semi-arid region of Nigeria. *ARNP J Agric Biol Sci*, 3: 6-9.
- Riva J, R Rizzi, S Marelli, and LG Cavalchini, 2004. Body measurements in Bergamasca sheep. *Small Rum Res*, 55: 221-227. doi: 10.1016/j.smallrumres.2003.12.010.
- Rombaut D, 1980. Comportement du mouton Djallonké en élevage rationnel. *Revue Elev Méd Vét Pays Trop*, 33: 427-439.
- Salako AE and LO Ngere, 2002. Application of multifactorial structural discriminant analysis in the morphometric structural differentiation of West African Dwarf and Yankassa sheep in South West Nigeria. *Niger J Anim Prod*, 29: 163-167.
- Schoenian S, 2011. Sheep 201. *Bigginers Guide to Raising Sheep*. Available in: <http://www.sheep101.info/201>
- Siddiqui HUR, AA Ahamad, and ZK Han, 2005. Biometrical studies of testes of the ram. *J Agric Soc Sci*, 1: 78-79.
- Söderquist L, and F Hultén, 2006. Normal values for the scrotal circumference in rams of gotlandic breed. *Reprod Domest Anim*, 41: 61-2.
- SWAC-OECD / ECOWA, 2008. Livestock and regional market in the Sahel and West Africa Potentials and challenges. 170p.
- Traoré A, HH Tamboura, A Kaboré, LJ Royo, I Fernández, I Álvarez, M Sangaré, D Bouchel, JP Poivey, L Sawadogo, and F Goyache, 2008. Multivariate analyses on morphological traits in Burkina Faso goat. *Arch Anim Breed*, 51: 588-600.
- Vallerand F, and R Branckaert, 1975. La race ovine Djallonké au Cameroun. Potentialités zootechniques, conditions d'élevage, avenir. *Revue Elev Méd Vét Pays Trop*, 28: 451-457.
- Yakubu A, and IS Musa-Azara, 2013. Evaluation of three mathematical functions to describe the relationship between body weight, body condition and testicular dimensions in Yankassa sheep. *Int J Morphol*, 31: 1376-1382.
- Yapi-Gnaoré CV, 1994. Caractères phénotypiques de différenciation et de croissance des agneaux de race pure Djallonké et croisés Sahélien x Djallonké. In: Proc. 2nd Biennial conference of the African Small Ruminant Res Network, Arusha, Tanzania, December 7-11, 215-219.
- Yapi-Gnaoré CV, A Oya, B Dagnogo, and M Ouattara, 1995. Influence de la sélection du poids du bélier sur la croissance de la descendance. <http://www.fao.org/wairdocs/ilri/x5473b/x5473b07.htm>.
- Youssao AKI, S Farougou, BG Koutinhouin, G Bio Bagou, and BD Kora, 2008. Aptitudes maternelles de la brebis Djallonké en élevage traditionnel dans la Commune de Banikoara au Bénin. *Revue Méd Vét*, 159: 538-544.