

First limnological characterization of Lakes Leqinat and Drelaj in Bjeshkët e Nemuna National Park, Kosovo

Robert Schabetsberger, Linda Grapci-Kotori, Halil Ibrahim, Astrit Bilalli, Zlatko Levkov, Christian D. Jersabek, Marcel Vorage, Mathieu Denoël, Roland Kaiser, Karin Pall, Ursula Eisendle, Erich Eder & Almedina Sadiku

Keywords: Balkan Peninsula, conservation, diversity, *Ichthyosaura alpestris*, plankton

Abstract

The Balkan Peninsula is a biodiversity hotspot and hosts numerous mountain lakes, which offer a refuge for a multitude of species. However, previous pristine habitats have been deeply affected by anthropogenic change, such as non-native fish introductions, which calls for multi-species considerations in the last remaining unaltered habitats. We carried out abiotic measurements and biodiversity assessments in two neighbouring alpine lakes, Lakes Leqinat and Drelaj in the Bjeshkët e Nemuna National Park in Kosovo, in August 2018. Lake Leqinat is a permanent, stratified water body and exhibits weak oxygen depletion below 3 m. Phytoplankton was dominated by chrysophycean, cryptophycean and chlorophycean algae. Zooplankton consisted of five rotifer species and *Daphnia longispina*. A mark-recapture experiment yielded a population of alpine newts (*Ichthyosaura alpestris*) of nearly 4000 adult individuals. In contrast, cold water from the surrounding karst seeps into Lake Drelaj, which is a well-oxygenated temporary lake. Hence phytoplankton and zooplankton biomasses were considerably lower than in Lake Leqinat. Phytoplankton was dominated by cryptophycean, chlorophycean, and bacillariophycean algae. Zooplankton consisted of the diaptomid copepod *Mixodiaptomus tatricus*, the cladoceran *Daphnia rosea*, and the anostracan *Chirocephalus diaphanus*. Conservation efforts should ensure that Lake Leqinat remains unstocked as introduced fish would probably destroy the natural community.

Profile

Protected area

Bjeshkët e Nemuna

National Park

Mountain range

Bjeshkët e Nemuna/

Prokletije

Country

Kosovo

Introduction

The Balkan Peninsula is as an important biodiversity hotspot (Griffiths et al. 2004). Mountain lakes harbour unique invertebrate communities often containing large zooplankton species like diaptomid copepods or anostracans (Belmonte et al. 2018; Mancinelli et al. 2019). Under natural, fish-free conditions, amphibians are the native top predators of these aquatic ecosystems (Lejeune et al. 2018; Schabetsberger & Jersabek 1995; Schabetsberger et al. 2006). Among them, newts show an impressive pattern of intraspecific diversity, with several locally described subspecies and phenotypes, especially in Montenegro (see e.g. Radovanović 1951, 1961; Džukić et al. 1990). In neighbouring Kosovo, Bjeshkët e Nemuna National Park was established in 2012 in the Albanian Alps (the Prokletije, or *Accursed Mountains*). It encompasses 630 km² of mountainous terrain with dense deciduous and coniferous forests and alpine landscapes. So far, only two limnological studies have been conducted in the park, providing information on algae, zooplankton and macrozoobenthos within the Gjeravica Lakes (Živić et al. 1997; Urošević 1997).

The biggest threat to organisms in closed, permanent alpine lakes is stocking with alien fish species. Introduced fish can drive sensitive prey species to ex-

tinction (Knapp et al. 2001; Schindler & Parker 2002; Denoël et al. 2005, 2009; Tiberti et al. 2013; Ventura et al. 2017 and references therein). Large (>1 mm) and often pigmented alpine zooplankton organisms can coexist with natural amphibian predators (Schabetsberger & Jersabek 1995), but they disappear after stocking as a result of size-selective predation by alien fish (Brancelj 1999; Schabetsberger et al. 2009). Most native amphibian species also disappear after fish introduction, as their larvae and sometimes even the adults are particularly vulnerable (Pilliod & Peterson 2001; Knapp et al. 2001, 2005; Denoël et al. 2005, 2009; Miró et al. 2018). Although terrestrial landscapes often remain wild or traditionally managed, large declines of newts have been reported due to fish stocking, the primary local driver of their extirpation in the Balkans. This region is a hotspot for paedomorphic populations. Larval structures, such as gills and gill slits, are retained in adult animals. Many paedomorphic populations have become extinct in the Balkans, including all larger populations from Montenegro (Denoël et al. 2005, 2019). The loss of this intraspecific heterochrony represents a massive loss to diversity and there is an urgent need for better conservation of the remaining populations. Lakes in national parks containing newts are becoming rare and are therefore important targets for monitoring.




Figure 1 – Bathymetric map of Lake Leqinat © Google Earth Pro, CNES, Airbus. Insert on top left shows the geographical position of Kosovo in Europe (red) © David Linzgo (Wikimedia Commons, licensed under Creative Commons-Lizenz by-sa-2.0-de, <http://creativecommons.org/licenses/by-sa/2.0/de/legalcode>) and an aerial photograph of Lakes Leqinat (right) and Drelaj (left) looking South into the Albanian Alps. © Dini Begolli.

The aim of this preliminary, descriptive study was to support conservation efforts by characterizing the abiotic conditions and the biodiversity of phyto- and zooplankton as well as selected benthic organisms (see Darwall et al. 2018) of Lakes Leqinat and Drelaj in the Bjeshkët e Nemuna National Park. Given the important regulatory role of alpine newts (*Ichthyosaura alpestris*, Laurenti 1761), which are the top predators, and their decline in the region, their population size within Lake Leqinat was assessed. Based on the results, the probability of the survival of alien fish is discussed.

Materials and Methods

Study sites

The two study lakes in western Kosovo are pristine, fishless, high-elevation water bodies just below the timberline that had never been investigated before and could serve as refuges for endemic species. Lake Leqinat is permanent ($42^{\circ} 40.110' N$, $20^{\circ} 5.455' E$; 1860 m a.s.l.; Figure 1), whereas Lake Drelaj ($42^{\circ} 40.041' N$, $20^{\circ} 5.956' E$; 1801 m a.s.l., Figures 2A and 2B), 0.7 km east of Leqinat, is temporary and dries out in summer



Figure 2 – Lake Drelaj: A on 5 August 2018; B on 27 September 2018. © Linda Grapci-Kotori

or autumn. Both lakes are free of fish and inhabited by alpine newts.

Sampling

Lake Leqinat

Lake Leqinat was sampled from 1 to 4 August 2018. It was sounded along 6 transect lines (108 soundings in total) across the lake to generate a bathymetric map. Surface area and volume were estimated using Fields R package version 10.0 (Nychka et al. 2017). Profiles of temperature, pH, conductivity and oxygen were measured at 1-m intervals at the deepest point of the lake (4.5 m). Water samples were taken with a 1.5-l Schindler trap. Temperature ($\pm 0.1^\circ\text{C}$) was measured using a thermometer mounted inside the trap. Dissolved oxygen, oxygen saturation, pH and conductivity were measured with a HACH HQ30d portable multi-meter. Integrated plankton samples were collected by towing a 30- μm plankton net vertically from the bottom to the surface. Quantitative phytoplankton and zooplankton samples were collected with the Schindler trap at depths of 1, 2 and 3 m. For phytoplankton, samples of 100 ml of unfiltered water were preserved with Lugol's solution (Schwoerbel 1993) and kept in brown glass bottles. Zooplankton samples were filtered through a 30- μm mesh sieve and preserved in 4% formaldehyde. Macrophytes were collected in the littoral area. Samples for diatoms were taken from stones, sand and mud with a spoon or a pipette. Stones were scraped using a toothbrush and the surface biofilm was removed by gentle washing. Parts of submerged macrophytes and mosses were collected by hand and put into plastic vials. All samples were preserved using 4% formaldehyde. Rotifers and nematodes were sampled by snorkeling and towing a small plankton net (20 cm diameter, 30- μm mesh size) through the dense benthic algal mats. Adult trichopterans were caught with entomological nets in the vicinity of the lake. A capture-mark-recapture experiment (Chapman estimator, Krebs 1989) was conducted to estimate the total population size of alpine newts (*Ichthyosaura alpestris*) from Lake Leqinat. The newts were caught by sight with dip nets around the shoreline and by snorkeling in deeper parts of the lake. They were anaesthetized in MS222 ($\sim 50 \text{ mg l}^{-1}$) and marked by clipping one toe for release and recapture. This method does not affect newt survival (Arntzen et al. 1999) and toes regenerate fast (Gutleb 1991). Snout-vent length (i.e. from the tip of the snout to the end of the cloaca) and total length (i.e. from the tip of the snout to the end of the tail) of 30 males and females each were measured using ruler calipers. Adulthood and sex were determined on the basis of the shape of the cloaca (Denoël 2017).

Lake Drelaj

Lake Drelaj was sampled on 5 August 2018. The surface area was calculated from Google Earth Pro. Temperature, pH, conductivity and oxygen levels were

Table 1 – Phytoplankton species in Lakes Leqinat and Drelaj.

Lake Leqinat
Cyanobacteria
<i>Aphanocapsa incerta</i> (Lemmermann) G. Cronberg & Komárek
<i>Aphanocapsa</i> sp.
<i>Gloeobacter violaceus</i> Rippka, J.B. Waterbury & Cohen-Bazire
<i>Microcystis</i> sp.
<i>Planktothrix</i> sp.
<i>Radiocystis geminata</i> Skuja
Chlorophyceae
<i>Botryococcus braunii</i> Kützing
<i>Desmodesmus brasiliensis</i> (Bohlin) E. Hegewald
<i>Didymocystis</i> sp.
<i>Elakatothrix genevensis</i> (Reverdin) Hindák
<i>Neglectella solitaria</i> (Witrock) Stenclová & Kastovsky
<i>Oocystis</i> sp.
<i>Planktosphaeria gelatinosa</i> G.M. Smith
<i>Scenedesmus</i> sp.
<i>Tetraëdron minimum</i> (A. Braun) Hansgirg
Conjugatophyceae
<i>Cosmarium</i> cf. <i>baileyi</i> Wolle
<i>Cosmarium</i> cf. <i>laeve</i> Rabenhorst
<i>Cosmarium</i> spp.
<i>Gonatozygon brebissonii</i> De Bary
<i>Staurastrum</i> sp.
<i>Staurodesmus</i> sp.
<i>Zygnema</i> sp.
Chrysophyceae
<i>Mallomonas</i> sp.
<i>Pseudopedinella</i> sp.
<i>Uroglena</i> sp.
Dinophyta
<i>Ceratium hirudinella</i> (O.F. Müller) Dujardin
<i>Glenodinium</i> sp.
<i>Gymnodinium</i> sp.
Cryptophyceae
<i>Cryptomonas</i> cf. <i>erosa</i> Ehrenberg
<i>Cryptomonas marssonii</i> Skuja
Bacillariophyceae
<i>Achnanthes</i> sp.
<i>Cymbella</i> sp.
<i>Melosira varians</i> C. Agardh
<i>Nitzschia acicularis</i> (Kützing) W. Smith
<i>Stauroneis</i> sp.
Lake Drelaj
Chlorophyceae
<i>Pseudopediastrum boryanum</i> (Turpin) E. Hegewald
<i>Scenedesmus</i> sp.
Conjugatophyceae
<i>Gonatozygon brebissonii</i> De Bary
Cryptophyceae
<i>Cryptomonas erosa</i> Ehrenberg + sp.
<i>Cryptomonas</i> cf. <i>marssonii</i> Skuja
Bacillariophyceae
<i>Amphora ovalis</i> (Kützing) Kützing
<i>Cyclotella</i> sp.
<i>Cymbella</i> spp.
<i>Cymbopleura inaequalis</i> (Ehrenberg) Krammer
<i>Navicula</i> sp.
<i>Nitzschia</i> sp.
<i>Surirella librile</i> (Ehrenberg) Ehrenberg

measured at the surface and above the sediment (3 m depth). One phyto- and one zooplankton sample were taken at 3 m depth. The same methods as described above for Lake Leqinat were applied for phyto-, zooplankton, diatoms and trichopterans. Both lakes were visited again for visual inspection on 26–27 September 2019 with students from Salzburg and Prishtina Universities.

Analyses

Algae were counted under an inverted microscope (Telaval 3, Jena; magnification 40–1000x), applying the method of Utermöhl (1958). Biovolumes were calculated by fitting geometric forms to cell dimensions (Deisinger 1984). The organic content of the samples for diatoms was removed by acid digestion, with the addition of 2 ml of a supersaturated solution of K_2MnO_4 and 4 ml of HCl to a small (ca. 2 ml) subsample. The acid was then removed through a series of water washes. Permanent slides of the cleaned material were mounted with Naphrax®. Slide observations were performed using a Nikon E–80i light microscope, and photomicrographs were taken with a Nikon Coolpix 600 digital camera (40–1000x). Trichopterans were identified under a dissecting microscope (Ibrahimi et al. 2019).

Zooplankton samples were stained with Rose Bengal. Subsamples were counted under the inverted microscope (magnification 40–100x). Biovolumes and biomasses were calculated by approximating natural shapes with geometric formulae for rotifers (Ruttner-Kolisko 1977), and from length–weight regressions for crustaceans (Downing & Rigler 1984). Periphytic nematode samples were mounted on slides in glycerin after a slow, two-step dehydration process (Decraemer et al. 2019).

Results

Lake Leqinat

Lake Leqinat reaches a maximum depth of just 4.5 m, in the southern part of the basin (Figure 1). It has an area of 16900 m² and a volume of 23000 m³. Depth profiles of abiotic parameters showed a weakly stratified, oxygen-rich upper layer down to 3 m depth. High pH-values (>9) indicate biogenic decalcification processes. Water of lower oxygen content and higher conductivity was found in the deepest layer (Figure 3).

The phytoplankton community (Table 1) was dominated by the chrysophycean genus *Uroglena* sp., reaching highest densities at around 2 m depth and accounting for 80.5% of the total biovolume (Figure 4A). The cryptophycean species *Cryptomonas marssonii* (9.5%) also reached its highest densities within this intermediate depth layer, while the dominant chlorophycean *Oocystis solitaria* was almost homogeneously distributed throughout the water column, contributing 5.0 to 43.3% of the total biovolume. A total of 391 benthic and periphytic diatom species were recorded in both

Table 2 – Rotifers, nematodes and crustaceans in Lakes Leqinat and Drelaj. Euplanktonic species are marked with an asterisk.

Lake Leqinat
Rotifera
<i>Anuraeopsis fissa</i> (Gosse)*
<i>Ascomorpha saltans indica</i> Koste*
<i>Cephalodella apocolea</i> Myers
<i>Cephalodella edax</i> Hollowday*
<i>Cephalodella forticula</i> (Ehrenberg)
<i>Cephalodella</i> sp.
<i>Cephalodella ventripes</i> (Dixon-Nuttall)
<i>Collotheca</i> sp.
<i>Colurella obtusa</i> (Gosse)
<i>Eothinia elongata</i> (Ehrenberg)
<i>Euchlanis dilatata</i> Ehrenberg
<i>Lecane closterocerca</i> (Schmarda)
<i>Lecane flexilis</i> (Gosse)
<i>Lecane luna</i> (Müller)
<i>Lecane lunaris</i> (Ehrenberg)
<i>Lepadella patella</i> (Müller)
<i>Monommata</i> sp.
<i>Mytilina mucronata</i> (Müller)
<i>Pleurotrocha petromyzon</i> Ehrenberg
<i>Polyarthra luminosa</i> Kutikova*
<i>Polyarthra</i> sp.*
<i>Proales fallaciosa</i> Wulfert
<i>Trichocerca bidens</i> (Lucks)
<i>Trichocerca rattus</i> (Müller)
<i>Trichocerca</i> cf. <i>relicta</i> Donner
Nematoda
<i>Eumonhystera</i> cf. <i>barbata</i> Andrassy
<i>Eumonhystera</i> cf. <i>pseudopulbosa</i> (Daday) Andrassy
<i>Mesodorylaimus</i> spec. 1
<i>Tridontulus</i> spec. 1
<i>Plectus</i> spec. 1
<i>Plectus</i> spec. 2
<i>Rhabdolaimus</i> cf. <i>terrestris</i> DeMan
Cladocera
<i>Daphnia longispina</i> (Müller)
Copepoda
Cylopidae Gen. sp.
Lake Drelaj
Rotifera
<i>Bdelloidea</i> Gen spp.
Cladocera
<i>Daphnia rosea</i> Sars
Copepoda
<i>Mixodiaptomus tatricus</i> (Wierzejski)
Anostraca
<i>Chirocephalus diaphanus</i> Prévost

lakes (no separate counts available; see Appendix A, Supplementary Data). A dense mat of *Chara contraria* A. Braun ex Kützing covered large areas of Lake Leqinat down to the deepest part. In shallow regions, *Potamogeton alpinus* Balb. was present. At the shoreline, *Carex vesicaria* L. dominated.

Zooplankton diversity was low, with only five euplanktonic rotifers and a single cladoceran species




Figure 3 – Profiles of temperature, pH, conductivity and oxygen saturation in Lake Leqinat on 2 August 2018.

recorded in the samples; copepods were encountered only as immature developmental stages of Cyclopoida. Densities of *Daphnia longispina* increased with depth. The zooplankton community was dominated by the rotifers *Anuraeopsis fissa* (20.3% of the biomass) and *Polyarthra luminosa* (42.4%), with peak densities at 1 m depth (Figure 5A). A more diverse rotifer fauna was recorded in the benthic *Chara* mats (Table 2). In the sediment and periphytic samples, 7 nematode taxa were observed (Table 2). Six adult trichopteran species were found at Lake Leqinat (*Limnephilus centralis*, *L. flavospinosus*, *L. flavicornis*, *L. stigma*, *Agrypnia varia*, *Oligotricha striata*).

The alpine newt was the only amphibian recorded. Only the metamorphic phenotype was found among the adults. A total of 108 females and 131 males were marked and released. After one day, 136 females and 145 males were recaptured. The capture-mark-recapture experiment yielded a total population size of 1492 female (683–2302; 95% confidence interval) and 2408 male (925–3891) alpine newts in Lake Leqinat. The total population was estimated to be 3980 adults (2262–5699). Snout-vent length and total length were 49.15 mm (41.9–58.8; median, minimum-maximum)

and 85.1 mm (77.1–94.1) for females, and 42.2 mm (38.1–45.4) and 73.2 mm (67.1–78.2) for males.

Lake Drelaj

Lake Drelaj (Figure 2) is 3 m deep but had dried out between September and October. Its maximum surface area is around 3100 m². Cold water from the surrounding karst seeps into the well-oxygenated lake, which in August fed a brook (~ 5 l s⁻¹) that emerged from the rocks about 30 m below the lake. On 5 August 2018, at the water surface and at 3 m depth, the temperatures were 9.3 and 5.5°C, Oxygen 9.48 mg l⁻¹ (105% saturation) and 9.27 mg l⁻¹ (98%), pH 8.29 and 8.17, and conductivity 192 µS cm⁻¹. The diatom *Cyclotella delicatula* dominated the open water community (84.5% of the biomass; Figure 4B). Overall phytoplankton biomass in the cold, clear lake was approximately one tenth of that in Lake Leqinat (Figures 4A and 4B). The zooplankton community consisted of three crustacean species: *Daphnia rosea*, *Mixodiptomus taticus* and *Chirocephalus diaphanus*. No rotifer species were recorded, other than a few tycho planktonic bdelloid species. Zooplankton biomass was 6 times less than in Lake Leqinat (Figures 5A and 5B).




Figure 4 – Phytoplankton. A: Depth profiles of phytoplankton biovolume in Lake Leqinat on 2 August 2018. B: Biovolume of planktonic algae at 3 m depth in Lake Drelaj on 5 August 2018.

The anostracan species *C. diaphanus* reached very high densities in the puddles that remained in September, where adult metamorphosed and larval alpine newts were also found. Two adult trichopteran species (*Allogamus uncatus*, *Limnephilus bipunctatus*) were recorded in the close vicinity of the lake.

Discussion

Abiotic conditions, phyto- and zooplankton communities

The two alpine lakes differ markedly in their hydrology. Whereas Lake Leqinat is permanent, Lake Drelaj can dry out in summer / autumn. The latter has a visible outflow and comparatively faster water renewal, indicated by colder water temperatures, higher conductivity and lower pH. Lake Leqinat exhibited signs of biogenic decalcification in the upper water column and weak oxygen depletion / ion enrichment in the deepest part of the lake, probably due to summer stagnation, while Lake Drelaj was well oxygenated throughout the water column. Hence phytoplankton and zooplankton biomasses were considerably lower in Lake Drelaj than in Lake Leqinat. Accordingly, species assemblages in the open water also differed markedly. In Lake Leqinat, chrysophycean and chlorophycean species dominated. Zooplankton diversity was low, with only five euplanktonic rotifers, and *Daphnia longispina* as the sole microcrustacean encountered. Although biomasses were low, species compositions of both phyto- (*Oocystis solitaria*, *Uroglena* sp.) and zooplankton (*Anuraeopsis fissa*) reflect a meso- to eutrophic character of the lake and are reminiscent of lowland ponds rather than of alpine lakes. In Lake Drelaj, an impoverished phytoplankton community with few small-sized (nanoplanktonic) species was observed together with three crustacean species in low abun-

dances, presumably as a consequence of the lake's hydrological regime. Euplanktonic rotifers were entirely missing.

No diaptomid copepod was found in Leqinat, which may be due to the high feeding pressure of the large alpine newt population (see Dodson 1970; Wissinger et al. 1999). On the other hand, Drelaj contains a considerably smaller newt population (see below) associated with *Mixodiaptomus tatricus*, a species typical of temporary high-altitude water bodies in the Alps and the Balkans (Jersabek et al. 2001; Mancinelli et al. 2019). We are aware that *Daphnia rosea* may be a morphotype and belongs to the *D. longispina* complex (Petrušek et al. 2008; Błędzki & Rybak 2016), but we acknowledge morphological differences and distinct ecological divergences between the two taxa, as known from numerous studies in the Austrian Alps (Gaviria-Melo et al. 2005), by assigning the taxon from Drelaj to *D. rosea*. *Chirocephalus diaphanus* is one of the most widespread and tolerant anostracan species in Europe (Brtek & Thiéry 1995; Demeter & Mori 2004). Its presence confirms regular drying of the lake, as the diapausing eggs usually need such drying up for successful development (Brendonck 1996; Zarattini et al. 2002).

Benthic communities

Diatom assemblage is very rich, indicating the high ecological status of the lakes. The large number of species results from the presence of different microhabitats such as macrophytes, stones and pebbles, and fine and coarse sediment, but it probably also reflects low human impact on the lakes. Several rare and endangered diatom species of the flora of Central and Eastern Europe (Lange-Bertalot & Steindorf 1996) have been recorded. The recently discovered *Neidiopsis borealis* (Vidakovic et al. 2019) was also found. Appre-




Figure 5 – Zooplankton. A: Depth profiles of zooplankton biomass in Lake Leqinat on 2 August 2018. B: Zooplankton biomass at 3 m depth in Lake Drelaj on 5 August 2018.

ciable macrophyte vegetation was detected only in the permanent Lake Leqinat. However, the three species registered (*Chara contraria*, *Potamogeton alpinus* and *Carex vesicaria*) typically co-occur in oligo- to mesotrophic alpine lakes. *Carex vesicaria* also grows around eutrophic lakes. Nematode taxa represent common sediment and periphytic taxa. The most abundant, *Monobrystera* and *Plectida*, are deposit feeders essential for ecological functioning at the base of any freshwater food web. *Tridontus* species are still rarely recorded, because they are often misidentified as *Eumonobrystera dispar*. Three of the trichopteran species are new records for Kosovo and are currently known only from Lake Leqinat: *Limnephilus flavospinosus*, *Limnephilus flavicornis* and *Oligotricha striata*, the last one being known in the Balkan Peninsula from only a few localities (Ibrahimi et al. 2019).

A large alpine newt population inhabits Lake Leqinat. Approximately 4000 metamorphosed adults were found, corresponding to a density of 0.24 individuals per square metre within the lake. Higher densities have been reported for similar-sized lakes in northern Greece (Denoël & Schabetsberger 2003; > 30000 individuals), but these included a large proportion of paedomorphic individuals. Hridsko Jezero from the Montenegrin part of the mountain range also used to host a population of paedomorphs (Denoël et al. 2001), but now only metamorphosed individuals occur (Denoël et al. 2019). In Leqinat, the sex ratio was slightly biased towards males (0.62). In other populations of alpine newts from the Balkans, sex ratios were found to be either male- or female-biased (Kalezić et al. 1989; Denoël 2003). This may be explained by different sex-specific payoffs across varied environmental conditions. We were unable to conduct a mark-recapture experiment in Drelaj, but the densities were lower, probably due to the particularly unfavourable cold breeding conditions (5°C at the bottom

in summertime). Both locations appear to be new records for the occurrence of the alpine newt (Džukić et al. 2015).

Conservation aspects

Although the lakes so far are pristine, some threats do exist. While in temporary Lake Drelaj, fish introduction poses no threat to the alpine newts, alien fish could potentially survive in Lake Leqinat. Since we have no information about the winter oxygen conditions under ice, we cannot speculate about the survival of salmonids, but it is very likely that minnows (*Phoxinus* sp.) or other species that tolerate low oxygen levels would live through the winter, destroy the alpine newt population, and strongly alter the community of prey organisms (Schabetsberger et al. 2006; Miró et al. 2018). Indeed, many major populations of alpine newts that inhabited mountain lakes in the Balkans, even within National Parks, have now vanished because of fish introductions. Such introductions were particularly detrimental in Montenegro, including in Hridsko Jezero, where fortunately fish have recently disappeared (Denoël et al. 2019). Although we are not aware of any immediate stocking campaigns, hydro-power development in the Rugova Valley is an issue and may be associated with stocking campaigns in the region (personal observation). Additionally, a large area of woodland near Lake Leqinat has been burnt deliberately. This calls for an enforcement of the protection of the lakes by rangers, and a visitor centre or at least thematic display panels near the lakes to raise awareness of conservation issues.

Acknowledgements

Funding for this study was provided by the Austrian Development Cooperation (HERAS project

K-04-2017). Field work was carried out under permit 2566/18/336-DMMU, issued by the Ministry of Environment and Spatial Planning (21/05/2018), Kosovo. We thank Nazmije Gervalla, Edonisa Hasanaj, Diellza Kadolli, Albulena Qamili and Qendrim Ramshaj for help in the field, and Tobias Schernhammer for help with species identification.

References

- Arntzen, J.W., A. Smithson & R.S. Oldham 1999. Marking and tissue sampling effects on body condition and survival in the newt *Triturus cristatus*. *Journal of Herpetology* 33: 567–576. Doi: 10.2307/1565573
- Belmonte, G., G. Alfonso, N. Beadini, P. Kotori, S. Mali, S. Moscatello & B. Shkurtaj 2018. An inventory of invertebrate fauna of Albania and Macedonia lakes. *Thalassia Salentina* 40: 27–38. Doi: 10.1285/i15910725v40sup2p27
- Błędzki, L.A. & J.I. Rybak 2016. *Freshwater crustacean zooplankton of Europe. Cladocera & Copepoda (Calanoida, Cyclopoida). Key to species identification, with notes on ecology, distribution, methods and introduction to data analysis*. Doi: 10.1007/978-3-319-29871-9
- Brancelj, A. 1999. The extinction of *Arctodiaptomus alpinus* (Copepoda) following the introduction of charr into a small Alpine lake Dvojno Jezero (NW Slovenia). *Aquatic Ecology* 33: 355–361. Doi: 10.1023/A:1009972108485
- Brendonck, L. 1996. Diapause, quiescence, hatching requirements: what we can learn from large freshwater branchiopods (Crustacea: Branchiopoda: Anostraca, Notostraca, Conchostraca), In: Alekseev, V.R. & G. Fryer (eds.), Diapause in the Crustacea. *Hydrobiologia* 320: 85–97. Doi: 10.1007/BF00016809
- Brtek, J. & A. Thiéry 1995. The geographic distribution of the European branchiopods (Anostraca, Notostraca, Spinicaudata, Laevicaudata). *Hydrobiologia* 298: 263–280. Doi: 10.1007/BF00033821
- Darwall, D., V. Bremerich, A. De Wever, A.I. Dell, J. Freyhof, M.O. Gessner, H.-P. Grossart, I. Harrison, K. Irvine, S.C. Jähnig, J.M. Jeschke, J.J. Lee, C. Lu, A.M. Lewandowska, M.T. Monaghan, J.C. Nejtgaard, H. Patricio, A. Schmidt-Kloiber, S.N. Stuart, M. Thieme, K. Tockner, E. Turak & O. Weyl 2018. The Alliance for Freshwater Life: A global call to unite efforts for freshwater biodiversity science and conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems* 28: 1015–1022. Doi: 10.1002/aqc.2958
- Decraemer, W., U. Eisendle-Flöckner & E. Abebe 2019. Nematoda. In: Thorp J.H. & A.P. Covich (eds.), *Freshwater Invertebrates, Fourth Edition, Volume IV, Keys to the Palaearctic Fauna*: 269–299. London. Doi: 10.1016/B978-0-12-804223-6.00001-9
- Deisinger, G. 1984. *Leitfaden zur Bestimmung der planktischen Algen der Kärntner Seen und ihrer Biomasse*. Kärntner Institut für Seenforschung. [In German]
- Demeter, L. & C. Mori 2004. Spatial distribution and habitat characteristics of *Chirocephalus diaphanus* (Branchiopoda: Anostraca) in the Retezat National Park (Southern Carpathians, Romania). *Biota* 5: 11–23.
- Denoël, M. 2003. Avantages sélectifs d'un phénotype hétérochronique. Eco-éthologie des populations pédomorphiques du Triton alpestre, *Triturus alpestris* (Amphibia, Caudata). *Cahiers d'éthologie fondamentale et appliquée, animale et humaine* 21: 1–327. [In French]
- Denoël, M. 2017. On the identification of paedomorphic and overwintering larval newts based on cloacal shape: review and guidelines. *Current Zoology* 63: 165–173. Doi: 10.1093/cz/zow054
- Denoël, M., R. Duguet, G. Džukić, M.L. Kalezić & S. Mazzotti 2001. Biogeography and ecology of paedomorphosis in *Triturus alpestris* (Amphibia, Caudata). *Journal of Biogeography* 28: 1271–1280. Doi: 10.1046/j.1365-2699.2001.00634.x
- Denoël, M., G. Džukić & M.L. Kalezić 2005. Effect of widespread fish introductions on paedomorphic newts in Europe. *Conservation Biology* 19: 162–170. Doi: 10.1111/j.1523-1739.2005.00001.x
- Denoël, M., G.F. Ficetola, R. Čirović, D. Radović, D. Džukić, M.L. Kalezić & T.D. Vukov 2009. A multi-scale approach to facultative paedomorphosis of European newts in the Montenegrin karst: distribution pattern, environmental variables and conservation. *Biological Conservation* 142: 509–517. Doi: 10.1016/j.biocon.2008.11.008
- Denoël, M., G.F. Ficetola, N. Sillero, G. Džukić, M.L. Kalezić, T. Vukov, I. Muhovic, V. Ikoćić & B. Lejeune 2019. Traditionally managed landscapes do not prevent amphibian decline and the extinction of paedomorphosis. *Ecological Monographs* 89: e01347. Doi: 10.1002/ecm.1347
- Denoël, M. & R. Schabetsberger 2003. Resource partitioning in two heterochronic populations of Greek Alpine newts, *Triturus alpestris veluchiensis*. *Acta Oecologica* 24: 55–64. Doi: 10.1016/S1146-609X(03)00043-2
- Dodson, S.I. 1970. Complementary feeding niches sustained by size-selective predation. *Limnology and Oceanography* 15: 131–137.
- Downing, J.A. & F.H. Rigler 1984. *A manual on methods for the assessment of secondary production in fresh waters*. IBP-Handbook 17, 2nd edition. Oxford.
- Džukić, G., M. Cvijanović, A. Urošević, T.D. Vukov, N. Tomasevic Kolarov, M. Slipjepcević, A. Ivanović & M.L. Kalezić 2015. The batrachological collections of the Institute for Biological Research “Siniša Stanković”. *Bulletin of the Natural History Museum* 8: 118–167. Doi: 10.5937/bnhmb1508118D
- Džukić, G., M.L. Kalezić, M. Tvrtković & A. Džorović 1990. An overview of the occurrence of paedomorphosis in Yugoslav newt (*Triturus*, Salamandridae) populations. *British Herpetological Society Bulletin* 34: 16–22.
- Gaviria-Melo, S., L. Forró, C.D. Jersabek & R. Schabetsberger 2005. Checklist and distribution of cladocerans and leptodorans (Crustacea: Branchiopoda) from Austria. *Annalen des Naturhistorischen Museums in Wien* 106: 145–216.

- Griffiths, H.I., B. Kryštek & J.M. Reed 2004. *Balkan Biodiversity. Pattern and Process in the European Hotspot*. Dordrecht. Doi: 10.1007/978-1-4020-2854-0
- Gutleb, B. 1991. Phalangengeneration und eine neue Methode zur Individualerkennung bei Bergmolchen, *Triturus alpestris* (Laurenti, 1768) (Caudata: Salamandridae). *Herpetozoa* 4: 117–125. [In German]
- Ibrahimi, H., L. Grapci-Kotori, A. Bilalli, A. Qamili & R. Schabetsberger 2019. Contribution to the knowledge of the caddisfly fauna (Insecta: Trichoptera) of Leqinat lakes and adjacent streams in Bjeshkët e Nemuna (Kosovo). *Natura Croatica* 28: 35–44. Doi: 10.20302/NC.2019.28.3
- Jersabek, C.D., F. Brancelj, F. Stoch & R. Schabetsberger 2001. Distribution and ecology of copepods in mountainous regions of the Eastern Alps. *Hydrobiologia* 453/454: 309–324. Doi: 10.1023/A:1013113327674
- Kalezić, M.L., G. Džukić & A. Popadić 1989. Paedomorphosis in Yugoslav Alpine newt (*Triturus alpestris*) populations: morphometric variability and sex ratio. *Arhiv Biološkib Nauka* 41: 67–79.
- Knapp, R.A., C.P. Hawkins, J. Ladau & J. McClory 2005. Fauna of Yosemite National Park has resistance but high resilience to fish introductions. *Ecological Applications* 15: 835–847. Doi: 10.1890/04-0619
- Knapp, R.A., K.E. Matthews & O. Sarnelle 2001. Resistance and resilience of Alpine lake fauna to fish introductions. *Ecological Monographs* 71: 401–421. Doi: 10.1890/0012-9615(2001)071[0401:RAROAL]2.0.CO;2
- Krebs, C.J. 1989. *Ecological methodology*. New York.
- Lange-Bertalot, H. & A. Steindorf 1996. Rote Liste der limnischen Kieselalgen (Bacillariophyceae) Deutschlands. *Schriftenreihe für Vegetationskunde* 28: 633–677. [In German]
- Lejeune, B., N. Sturaro, G. Lepoint & M. Denoël 2018. Facultative paedomorphosis as a mechanism promoting intraspecific niche differentiation. *Oikos* 127: 427–439. Doi: 10.1111/oik.04714
- Mancinelli, G., S. Mali & S. Belmonte 2019. Species richness and taxonomic distinctness of zooplankton in ponds and small lakes from Albania and North Macedonia: the role of bioclimatic factors. *Water* 11: 2384. Doi: 10.3390/w11112384
- Miró, A., I. Sabás & M. Ventura 2018. Large negative effect of non-native trout and minnows on Pyrenean lake amphibians. *Biological Conservation* 218: 144–153. Doi: 10.1016/j.biocon.2017.12.030
- Nychka, D., R. Furrer, J. Paige & S. Sain 2017. *Fields: Tools for spatial data, R package version 10.0*. Available at: <https://cran.r-project.org/web/packages/fields> Doi: 10.5065/D6W957CT
- Petrusek, A., A. Hobæk, J.P. Nilsen, M. Skage, M. Černý, N. Brede & K. Schwenk 2008. A taxonomic reappraisal of the European *Daphnia longispina* complex (Crustacea, Cladocera, Anomopoda). *Zoologica Scripta* 37: 507–519. DOI: 10.1111/j.1463-6409.2008.00336.x
- Pilliod, D.S. & C.R. Peterson 2001. Local and landscape effects of introduced trout on amphibians in historically fishless watersheds. *Ecosystems* 4: 322–333. Doi: 10.1007/s10021-001-0014-3
- Radovanović, M. 1951. A new race of the Alpine newt from Yugoslavia. *British Journal of Herpetology* 1: 93–97.
- Radovanović, M. 1961. Neue Fundorte neotenischer Bergmolche in Jugoslawien. *Zoologischer Anzeiger* 166: 206–218. [In German]
- Ruttner-Kolisko, A. 1977. Suggestions for biomass calculation of plankton rotifers. *Archiv für Hydrobiologie. Beihefte. Ergebnisse der Limnologie* 8: 71–76.
- Schabetsberger, R., S. Grill, G. Hauser & P. Wukits 2006. Zooplankton successions in neighboring lakes with contrasting impacts of amphibian and fish predators. *International Review of Hydrobiology* 91: 197–221. Doi: 10.1002/iroh.200610867
- Schabetsberger, R. & C.D. Jersabek 1995. Alpine newts (*Triturus alpestris*) as top predators in a high-altitude karst lake: daily food consumption and the impact on the copepod *Arctodiaptomus alpinus*. *Freshwater Biology* 33: 47–61. Doi: 10.1111/j.1365-2427.1995.tb00385.x
- Schabetsberger, R., M. Luger, G. Drozdowski & A. Jagsch 2009. Only the small survive: Monitoring long-term changes in the zooplankton community of an Alpine lake after fish introduction. *Biological Invasions* 11: 1335–1345. Doi: 10.1007/s10530-008-9341-z
- Schindler, D.W. & B.R. Parker 2002. Biological pollutants: Alien fishes in mountain lakes. *Water, Air, & Soil Pollution: Focus* 2: 379–397. Doi: 10.1023/A:1020187532485
- Schwoerbel, J. 1993. *Methoden der Hydrobiologie*. Stuttgart, Jena. [In German]
- Tiberti, R., A. von Hardenberg & A. Bogliani 2013. Ecological impact of introduced fish in high-altitude lakes: a case study from the European Alps. *Hydrobiologia* 724: 1–19. Doi: 10.1007/s10750-013-1696-1
- Urošević, V. 1997. Periphyton algae in the system of Djeravica Lakes on the spring branch of Erenik. *University Thought, Natural Sciences* 3: 11–21.
- Utermöhl, H. 1958. Zur Vervollkommnung der quantitativen Phytoplankton-methodik. *Internationale Vereinigung für Theoretische und Angewandte Limnologie: Mitteilungen* 9: 1–38. [In German]
- Ventura, M., R. Tiberti, T. Buchaca, D. Buñay, I. Sabás & A. Miró 2017. Why should we preserve fishless high mountain lakes? In: Catalan, J., J.M. Ninot & M. Aniz (eds.), High Mountain Conservation in a Changing World. *Advances in Global Change Research* 62: 181–205. Berlin, Germany. Doi: 10.1007/978-3-319-55982-7_8
- Vidakovic, D., P.H. Hamilton & Z. Levkov 2019. *Neidiopsis borealis* sp. nov., a new diatom species from the mountain Shar Planina, Macedonia. *Phytotaxa* 402: 21–28. Doi: 10.11646/phytotaxa.402.1.3
- Wissinger, S.A., A.J. Bohonak, H.H. Whiteman & W.S. Brown 1999. Subalpine wetlands in Colorado. Habitat permanence, salamander predation and invertebrate communities. In: Patzer, D.P., R.B. Rader & S.A. Wissinger (eds.), *Invertebrates in Freshwater Wetlands*

of *North America: Ecology and Management*: 757–789. New York.

Zarattini, P., V. Rossi, B. Mantovani & G. Mura 2002. A preliminary study in the use of RAPD markers in detecting genetic differences in hatching patterns of *Chirocephalus diaphanus* Prevost, 1803 (Crustacea: Anostraca). *Hydrobiologia* 486: 315–323. Doi: 10.1023/A:1021335826251

Živić, N., B. Miljanović, N. Labus & T. Jakšić 1997. Composition of zooplankton and macrozoobenthos in big and small Djeravica Lakes. *University Thought, Natural Sciences* 3: 51–56.

Authors

Robert Schabetsberger¹

is a zoologist at the University of Salzburg, Department of Biosciences. He works in freshwater and marine biology. E-mail: robert.schabetsberger@sbg.ac.at

Linda Grapci-Kotori² – corresponding author

is a zoologist and works at the University of Prishtina, *Hasan Prishtina*, Faculty of Mathematics and Natural Sciences, Department of Biology. Her work focuses on the ecology, diversity, conservation and taxonomy of fish. E-mail: linda.grapci@uni-pr.edu

Halil Ibrahimović² and Astrit Bilalli²

are entomologists and work at the University of Prishtina, *Hasan Prishtina*, Faculty of Mathematics and Natural Sciences, Department of Biology. Their work focuses on the taxonomy, diversity, ecology and biogeography of insects. E-mails: halil.ibrahimovic@uni-pr.edu; astritbilalli@yahoo.com

Zlatko Levkov³

has research interests in many aspects of diatom biology: taxonomy and biogeography, systematics and phylogeny, species flocks in ancient lakes, monitoring of pollution based on diatoms, and applications in forensic sciences. He works at the Ss. Cyril & Methodius University in Skopje, Institute of Biology. E-mail: zlevkov@yahoo.com

Christian D. Jersabek¹

is a freelance aquatic ecologist, with a specialization in zooplankton ecology and taxonomy. He also works as a Consulting Engineer for Limnology and Ecology to a number of official institutions and provincial governments in Austria. E-mail: christian.jersabek@sbg.ac.at

Marcel Vorage⁴

is a biologist, economist and geographer at the Salzburg University of Education Stefan Zweig. His work concerns environmental, social and economic sustainable development. E-mail: marcel.vorage@phsalzburg.at

Mathieu Denoël⁵

is Research Director of the Fonds de la Recherche Scientifique at the University of Liège (Belgium) where he heads the Laboratory of Ecology and Conservation of Amphibians (LECA). E-mail: mathieu.denoel@uliege.be

Roland Kaiser¹

is a plant ecologist and works as an environmental consultant. He works in botany, computational biology and digital cartography. E-mail: kardinal.eros@gmail.com

Karin Pall⁶

is a botanist at Systema GmbH, a private limnological company in Vienna, Austria. Her work focuses on the ecology and taxonomy of macrophytes, especially charophytes. E-mail: karin.pall@systema.at

Ursula Eisendle¹

works as a freelance associated postdoc at the University of Salzburg. Her main focus is on nematodes (e.g. taxonomic editor WoRMS) and aquatic environmental and educational research. E-mail: ursula.eisendle@sbg.ac.at

Erich Eder⁷

is a zoologist and Assistant Professor of evolutionary biology at the Sigmund Freud University, Medical School, Vienna. His work focuses on the ecology of large branchiopods, science didactics, and the background of creationism and paranormal beliefs. E-mail: erich.eder@med.sfu.ac.at

Almedina Sadiku²

is a biologist (M.Sc.) and currently works as a primary school teacher of biology. During her Master's, she studied *Ichthyosaura alpestris* in Lake Leqinat. E-mail: almedinasadiku06@gmail.com

¹ University of Salzburg, Department of Biosciences, Hellbrunnerstrasse 34, 5020 Salzburg, Austria

² University of Prishtina, *Hasan Prishtina*, Faculty of Mathematics and Natural Sciences, Department of Biology, Mother Teresa Street, 10000 Prishtina, Kosovo

³ Institute of Biology, Ss Cyril and Methodius University, Arhimedova 3, 1000 Skopje, Macedonia

⁴ Salzburg University of Education Stefan Zweig, Akademiestrasse 23, 5020 Salzburg, Austria

⁵ University of Liege (ULiege), Laboratory of Ecology and Conservation of Amphibians (LEPA), Freshwater and Oceanic science Unit of reSearch, 22 Quai van Beneden, 4020 Liege, Belgium

⁶ Systema GmbH, Bensasteig 8, 1140 Vienna, Austria

⁷ Sigmund Freud University in Vienna, Medical School, Freudplatz 3, 1020 Vienna, Austria