

## A SPECTROSCOPIC STUDY OF UM 673 A AND B: ON THE SIZE OF LYMAN-ALPHA CLOUDS<sup>1</sup>

A. SMETTE,<sup>2,3,4</sup> J. SURDEJ,<sup>3,5</sup> P. A. SHAVER,<sup>2</sup> C. B. FOLTZ,<sup>6</sup> F. H. CHAFFEE,<sup>6,7</sup> R. J. WEYMANN,<sup>8</sup>  
 R. E. WILLIAMS,<sup>9</sup> AND P. MAGAIN<sup>3,5</sup>

Received 1991 February 27; accepted 1991 September 10

### ABSTRACT

We present a study of the Ly $\alpha$  forest spectra (2 Å resolution) obtained for the A and B images of the gravitationally lensed high-redshift quasar UM 673. We also present higher resolution data of the brightest (A) image. In the 2 Å resolution spectra, all the absorption lines detected at 5  $\sigma$  in the spectrum of the fainter B image are present in the A image; however, we find two anticoincidences, i.e., two lines in A which do not have a counterpart in B at more than a 3  $\sigma$  confidence level. Given the fact that corresponding Ly $\alpha$  lines in the spectra of A and B have their equivalent widths well correlated, this proves that both light beams actually cross the same clouds. Most of the velocity differences between corresponding lines are compatible with 0 km s<sup>-1</sup> within the error bars, with a standard deviation of 17 km s<sup>-1</sup>.

As the comoving linear separation increases from virtually 0h<sub>50</sub><sup>-1</sup> kpc ( $H_0 = 50h_{50}$  km s<sup>-1</sup> Mpc<sup>-1</sup>,  $q_0 = 0$ ) at the redshift of the QSO to 2h<sub>50</sub><sup>-1</sup> kpc, we derive a best value of 12h<sub>50</sub><sup>-1</sup> kpc for the 2  $\sigma$  lower limit and of 160h<sub>50</sub><sup>-1</sup> kpc for the 2  $\sigma$  upper limit of the diameter of spherical Ly $\alpha$  clouds in the redshift range 2.1–2.7, by means of Monte Carlo simulations. However, if we interpret the two anticoincidences as due to a Mg II doublet at  $z = 0.4261$ , we find in this case a best value of 23h<sub>50</sub><sup>-1</sup> kpc to the 2  $\sigma$  lower limit of the Ly $\alpha$  cloud diameter, but we cannot derive any upper limit.

For the two major heavy-element systems detected in the spectrum of UM 673, we do not find any significant difference between the corresponding lines in the two spectra, indicating that these systems do not show dramatic variations over scales of 0.8h<sub>50</sub><sup>-1</sup> and 2.1h<sub>50</sub><sup>-1</sup> kpc, respectively.

*Subject headings:* gravitational lensing — quasars: absorption lines — quasars: individual (UM 673)

### 1. INTRODUCTION

The study of gravitational lens systems not only provides us with a unique tool to derive an independent estimate of the Hubble parameter  $H_0$  or the mass of the lensing galaxy and its associated cluster (Refsdal 1964) but also allows us to understand better the variations of the galactic or intergalactic medium over scales of the order of virtually zero to several kiloparsecs.

The most striking feature that appears in the blue wing of the Ly $\alpha$  emission line in the spectra of distant quasars is a forest of narrow absorption lines due to hydrogen clouds located along our line of sight (Sargent et al. 1980, hereafter SYBT). Various models have been invoked to describe their physical state: pressure-confined clouds in an expanding intergalactic medium (SYBT; Ikeuchi & Ostriker 1986), relics of primordial density fluctuations in the cold dark matter sce-

nario for the biased formation of galaxies (Bond, Szalay, & Silk 1988), gravitationally confined clouds in cold dark matter minihalos (Rees 1986), or thermally unstable shocks associated with the formation of protogalaxies (Lake 1988; Hogan 1987). Despite these various possible scenarios for the origin of the Ly $\alpha$  clouds, there is still no general agreement on values of basic physical parameters such as their typical size.

SYBT observed that the Ly $\alpha$  absorption lines reach zero intensity even on the Ly $\alpha$  emission-line wings, in 0.8 Å resolution spectra. Following the argument given by Rees (1970), they concluded that the clouds cover the emission-line region of the QSO and set a lower limit of the order of 10 pc for their characteristic size.

In a spectral range of 900 Å between 3450 and 4350 Å, covering 400 Å of the Ly $\alpha$  forest, Foltz et al. (1984) found two or three Ly $\alpha$  lines which are not common or are of substantially different strengths in comparing 1 Å resolution spectra for the two images of Q2345+007 (separation 7'3). Identifying the radius of a cloud with the separation between the light beams at the redshift ( $z_a = 1.951$ ) of the shortest wavelength common line, they determine a lower limit of roughly 5h<sub>50</sub><sup>-1</sup> kpc to 25h<sub>50</sub><sup>-1</sup> kpc for the radius of the Ly $\alpha$  clouds.

However, noticing that the column density distribution functions for Ly $\alpha$  clouds and heavy-element systems could be fitted by the same power law, Tytler (1987) proposed that they form a single population. The fact that no metal lines are found in the Ly $\alpha$  systems requires that they have total column densities of 10<sup>17</sup> cm<sup>-2</sup> or less, or, for those with  $N(\text{H I}) \sim 10^{17}$  cm<sup>-2</sup>, that they be nearly neutral. In such a picture the Ly $\alpha$  clouds are small ( $\leq 10^{18}$  cm) in general, with a wide variety of ionizations (and metal abundances up to 0.1 solar), while a few could be larger ( $\sim 10^{21}$  cm) and of high ionization. To explain

<sup>1</sup> Based on observations made at the MMT0, at ESO, and at the CTIO.

<sup>2</sup> European Southern Observatory, Karl-Schwarzschild-Strasse, 2, D-8046 Garching bei München, Germany.

<sup>3</sup> Institut d'Astrophysique, Université de Liège, Avenue de Coïnte, 5, B-4000 Coïnte, Belgium.

<sup>4</sup> Now at European Southern Observatory, Casilla 19001, Santiago 19, Chile.

<sup>5</sup> Also Chercheur Qualifié au Fonds National de la Recherche Scientifique (Belgium).

<sup>6</sup> Multiple Mirror Telescope Observatory, Smithsonian Institution and the University of Arizona, Tucson, AZ 85721.

<sup>7</sup> At Institute of Astronomy, Cambridge, England, from 1990 July to 1991 July.

<sup>8</sup> Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101.

<sup>9</sup> Cerro Tololo Inter-American Observatories, Casilla 603, La Serena, Chile.

the results of Foltz et al. (1984) most of the clouds should have high aspect ratios, of the order of  $10^5$  at least.

Recently, Pettini et al. (1990) obtained qualitatively the same conclusion about the size of the Ly $\alpha$  clouds: using a  $6 \text{ km s}^{-1}$  resolution spectrum of Q2206–199N and limiting themselves to low H I column density, they found that (1) most of the lines have a  $b$  parameter less than  $22 \text{ km s}^{-1}$ , (2) this  $b$  parameter is correlated with the cloud column density  $N_{\text{HI}}$  for  $\log N(\text{H I}) \approx 12.7\text{--}14.0$ .

From the low observed values of  $b$ , they concluded that Ly $\alpha$  clouds must be cold, hence neutral and small ( $10^{-5}$  pc) or sheetlike objects. They interpreted the observed correlation by attributing the line broadening more to large scale rather than to thermal motion, implying that *all* Ly $\alpha$  clouds are cooler than previously thought. From this point of view, Pettini et al. (1990) consider the result of Foltz et al. (1984) as a typical clustering length instead of an estimation of the cloud size; they invoke a possible second population of Ly $\alpha$  clouds, characterized by a higher H I column density than the ones they selected, in order to explain that some lines reach zero intensity in the wings of the Ly $\alpha$  emission line.

It should be pointed out that, if spherical, isolated Ly $\alpha$  clouds actually had the size quoted by Pettini et al. (1990), the Ly $\alpha$  forest of a single quasar would look completely different from one year to the other simply because of the relative motion of Earth, as already argued by Rees (1970).

On the other hand, a similar work conducted by Carswell et al. (1991) on the quasar Q1100–264 leads to different results, i.e., (1) the  $b$  parameter ranges from 12 to  $80 \text{ km s}^{-1}$  (with a different line selection), but few lines have low Doppler parameters; (2) the correlation reported between the Doppler parameter and the H I column density is contested.

Here we present results of the analysis of spectra obtained for the UM 673 A and B images (cf. Surdej et al. 1987, hereafter Paper I; Surdej et al. 1988, hereafter Paper II). In comparison with the data of Foltz et al. (1984), our data are characterized by a better signal-to-noise ratio, and they cover a much larger spectral range ( $1700 \text{ \AA}$ ), with  $1400 \text{ \AA}$  in the Ly $\alpha$  forest, but at  $2 \text{ \AA}$  resolution. An advantage of using the images of UM 673 as probes of the intergalactic medium is that the lens has been identified ( $z_d = 0.493$ ), but on the other hand, the separation between the A and B images is more than 3 times smaller ( $2''.22$ ) than for Q2345+007. In the remainder we shall use the value of 2.727 for the redshift of UM 673 (cf. Sargent, Boksenberg, & Steidel 1988, hereafter SBS). Preliminary results of this study have been reported by Smette et al. (1990).

## 2. SPECTROSCOPIC OBSERVATIONS

### 2.1. The 1 and 2 $\text{\AA}$ Resolution Spectra

Observations of the UM 673 A and B images were obtained (by C. F. and F. C.) using the Blue Channel of the MMT Spectrograph equipped with an intensified photon-counting Reticon detector. Details of the dates of observations, exposure times, spectral coverage, and resolutions are given in Table 1. All observations were taken through a pair of  $1'' \times 3''$  slits, with the long axis of the slit aligned along constant azimuth (i.e., the direction of atmospheric dispersion). Observations of image B were obtained only under the best seeing conditions; the seeing as estimated from examination of the images on the television guider monitor was at all times better than  $0''.7$  (FWHM). Furthermore, in order to monitor the contamination from image A, observations were periodically taken at

TABLE 1  
LOG OF OBSERVATIONS

UM 673 Image	UT Date	Exposure Time (s)	Wavelength Coverage ( $\text{\AA}$ )	Resolution (FWHM) ( $\text{\AA}$ )
MMT				
A.....	1987 Dec 11	7200	3300–4200	1
A.....	1988 Nov 10	600	3200–4800	2
A.....	1988 Nov 11	1200	3200–4800	2
B.....	1988 Nov 7	3600	3200–4800	2
B.....	1988 Nov 10	2400	3200–4800	2
B.....	1988 Nov 11	1200	3200–4800	2
CTIO 4 m Telescope				
A.....	1988 Nov 8	3600	3800–4700	0.33
A.....	1988 Nov 8	7200	3800–4700	0.33
A.....	1988 Nov 8	7200	3800–4700	0.33
A.....	1988 Nov 9	3600	3800–4700	0.33
A.....	1988 Nov 9	2268	3800–4700	0.33
A.....	1988 Nov 10	6000	3800–4700	0.33
A.....	1988 Nov 10	5400	3800–4700	0.33
B.....	1988 Nov 9	9000	3800–4700 <sup>a</sup>	0.33
ESO 3.6 m Telescope				
A.....	1987 Sep 3	4800	4000–5000	0.6
A.....	1987 Sep 3	4800	4000–5000	0.6
A.....	1987 Dec 3	2700	5000–6000	1

<sup>a</sup> Because the poor signal-to-noise ratio, only a  $200 \text{ \AA}$  interval has been illustrated in Fig. 2.

a position at the same elevation as that of image B, on the opposite side of the line of constant azimuth that intersected image A. *In no case was the contamination from image A found to be more than about 5% of the signal from image B.*

Data reductions followed standard procedures. The spectrograph is a two-channel device, allowing nearly simultaneous sky subtraction. Neither extinction correction nor flux calibration was carried out. Therefore, the spectra were reduced to units of photons  $\text{pixel}^{-1} \text{ s}^{-1}$  versus heliocentric wavelengths. Since the detector is a photon counter, the variance of each spectrum was calculated on a pixel-to-pixel basis assuming that all noise resulted from photon noise in the object, sky, and dark signal. Individual spectra covering the same wavelength regions were then combined, weighted by the inverses of their corresponding variances.

The  $1 \text{ \AA}$  spectrum (A image only) was calibrated with He-Ar spectra. The residuals to the fits were  $0.18 \text{ \AA}$  in both spectrograph apertures. Shortward of  $3500 \text{ \AA}$ , though, where there is a dearth of He-Ar lines, they were somewhat worse, perhaps twice as large. The  $2 \text{ \AA}$  spectra were calibrated with He-Ar and Hg-Cd spectra, which give a larger number of lines at shorter wavelengths. The residuals to the fits were found to be  $0.4 \text{ \AA}$  or less.

### 2.2. The 0.6 $\text{\AA}$ Resolution Spectrum

We have also used two CASPEC spectra of the A image obtained by P. M. at the coude focus of the ESO 3.6 m telescope at La Silla. The detector was an RCA CCD, with a pixel size of  $30 \mu\text{m}$ ; the readout noise had been reduced by using a binning of  $2 \times 2$  pixels during the reading. The grism had  $36.1 \text{ lines mm}^{-1}$  and the central wavelength was  $4500 \text{ \AA}$ . A white internal lamp has been used for the flat field, while exposures of

a thorium lamp immediately before and after the object observations were made for the calibration in wavelength. To calibrate the spectra in flux, exposures of the standard star HD 19445 were taken immediately afterward. The resolution is about  $0.6 \text{ \AA}$ . The reduction followed the standard procedure described in MIDAS (Banse et al. 1988). The two spectra were then added. The spectrum has been rebinned to vacuum heliocentric wavelengths. A rough estimate of the noise has been calculated using the difference between these two spectra. Unfortunately, it was not possible to correct properly for the blaze function at the shortest wavelengths, affecting particularly a region between 4055 and 4070  $\text{\AA}$ .

### 2.3. The $0.33 \text{ \AA}$ Resolution Spectra

The last spectra covering the Ly $\alpha$  forest were obtained by R. J. W. and R. E. W. at the Ritchey-Chrétien focus of the 4 m telescope at Cerro Tololo with the echelle spectrograph. Refer to Table 1 for the log book of these observations. The detector was the CCD TI #1 with a pixel size of  $15 \mu\text{m}$ ; a preflash of 20 ADU over the bias was used; a binning by 2 in the direction perpendicular to the dispersion was introduced for the readout. The central wavelength was  $4000 \text{ \AA}$ , with an echelle grating of  $79 \text{ lines mm}^{-1}$ ; all the observations were made through a  $\text{CuSO}_4$  filter. This setting introduced three major complications: first, consecutive orders do not overlap, so that gaps of several angstroms appear in the spectrum; second, there results a field curvature, introducing a variable point-spread function depending on the position in the frame; last, the difficulty of defining properly the center of the calibration lines affected the wavelength calibration, whose rms is about  $0.03 \text{ \AA}$ . Due to the relatively low signal-to-noise ratio of each frame, we had to use slightly modified MIDAS reduction procedures. We extracted the spectrum with an improved routine, kindly provided to us by W. Verschueren (Verschueren & Hensberge 1990) and slightly modified in order to take into account the sky background. The latter was estimated by constructing a sky spectrum which was the average of 6 pixels, i.e., 3 pixels on both sides of each order of the object spectrum; a median filter with a window of 41 pixels was then applied on each order of this spectrum. An estimate of the variance on every pixel of each (object) spectrum was obtained by computing the difference between itself and its two closest neighbors. Comparison with an estimate of the noise on the red wing of the Ly $\alpha$  emission line shows good agreement. The different spectra were then co-added, weighted by the inverses of their corresponding variances.

Only the  $200 \text{ \AA}$  of the B image spectrum for which the signal-to-noise ratio is the best are presented. A Gaussian filter with  $\sigma$  equal to the effective resolution of the instrument was applied. Because of the poor quality of these data, they were not used for the subsequent analysis. However, the reader should compare the absorption-line profiles in this spectrum with the ones in the corresponding spectrum of image A.

### 2.4. The $5000\text{--}6000 \text{ \AA}$ Spectrum

Finally, a spectrum of the A image covering the region from  $5000$  to  $6000 \text{ \AA}$  was taken by P. M. with CASPEC at the ESO 3.6 m telescope in La Silla. The detector was the RCA CCD #3. A binning of  $2 \times 2$  was applied at the reading to reduce the readout noise. The central wavelength was  $5500 \text{ \AA}$ . The wavelength calibration was made using a thorium lamp and leads to an rms of  $0.3 \text{ \AA}$ . Using the calibration lines, the

resolution is measured to be about  $1 \text{ \AA}$ . The reduction procedure followed the standard MIDAS one, except that the order extraction method was similar to the one described for the  $0.33 \text{ \AA}$  spectra. Since the data were not corrected for the instrumental response, we were not able to measure the exact positions of the emission-line peaks.

## 3. LINE LISTS

### 3.1. The $2 \text{ \AA}$ Resolution Spectra

Figures 1 and 2 show the spectra of the UM 673 A and B images and their difference, illustrating their remarkable similarities. Except for a scale factor, we were able to use the same continuum for both  $2 \text{ \AA}$  resolution spectra (hereafter, “ $2 \text{ \AA}$  spectra”) obtained by fitting a fifth-order polynomial. The profiles of the emission lines (O VI, Ly $\alpha$ , N V) have been fitted with Gaussians. The original spectra were then normalized.

Line lists were first made by visual inspection. Positions of the absorption lines were obtained by fitting the lines with Gaussian profiles—a good approximation for the point-spread function—since lines are not resolved at this resolution; obvious blends in these spectra were fitted by multiple Gaussians (up to 3). As we analyzed the two spectra similarly and independently, the higher resolution spectra were not used for any constraints. Formal errors on the equivalent widths  $W$  were obtained using the variance spectra. Lines were accepted if their equivalent widths were at least  $5 \sigma(W)$ . The line lists are presented in Table 2.

As a first result, all the 68 lines which are detected at the  $5 \sigma$  level in image B are present in image A. Because of the difference in signal-to-noise ratios, the opposite is not true. However, a great number of the B lines missing at the  $5 \sigma$  level are present at the  $3 \sigma$  level: they are indicated in parentheses in Table 2. If they are not detected at the  $3 \sigma$  level, we give the value of the  $3 \sigma$  upper limit to the equivalent width of these “missing” lines.

The difference spectrum  $A(2 \text{ \AA}) - B(2 \text{ \AA})$  is very flat, although it exhibits some systematic differences: between  $3240$  and  $3550 \text{ \AA}$ , the mean values measured on  $10 \text{ \AA}$  spectral ranges are of the




FIG. 1.— $2 \text{ \AA}$  resolution spectra of the A and B images of UM 673, and their difference. Wavelengths are vacuum heliocentric. For the presentation, the spectrum of image B has been multiplied by 7.9 and offset by 0.5, and that of image A has been offset by 0.1.






FIG. 2a

FIG. 2.—(a) Extended 2 Å spectra of the A and B images of UM 673 and the difference between the normalized spectra. For comparison the 1 Å, 0.6 Å, and 0.33 Å spectra of UM 673 image A are also shown. Wavelengths are vacuum heliocentric. Lines detected at the  $5\sigma$  level are indicated for the 2 Å, 1 Å, and 0.33 Å spectra, and at the  $4\sigma$  level for the 0.6 Å spectrum. (b) The C IV doublet in the 5000–6000 Å spectrum.




FIG. 2a—Continued

order of 0.1, with a standard deviation of 0.3, while between 4010 and 4200 Å the mean values are of the order of 0.06 and the standard deviation is of the order of 0.12. The flatness of the difference spectrum means that not only has the continuum been reasonably well fitted, but also it is nearly identical in the two spectra (at least in this spectral range). Any features in the difference spectrum are good indicators of variations between corresponding lines in the A and B spectra. They can be due to bad pixels in one or both of the spectra, differences in velocities between the two involved lines, or differences between equivalent widths. First, we define a coincidence as a line which is detected at the  $5\sigma$  level in both A and B spectra. On the other hand, since we are interested only in differences in equivalent widths, we call an anticoincidence a line which is detected at the  $5\sigma$  level in one spectrum, which is not detected at the  $3\sigma$  level in the other, and for which a feature is detected at the  $3\sigma$  level in the difference spectrum. We obtain 68 coincidences and two anticoincidences (at 3987 and 3998 Å, with 3.3 and 4.4 confidence levels respectively). The signal appearing between 4350 and 4400 Å is attributed to some “bad” pixels in the B spectrum and to a difference in velocity between two corresponding lines (cf. §§ 4.1 and 4.3.1).

### 3.2. The 0.33, 0.6, 1 Å Resolution and 5000–6000 Å Spectra

The line lists for the 0.6 and 1 Å spectra have been obtained in exactly the same way. In exceptional cases, fittings of mul-

iple Gaussians were made with up to six components for the 0.6 Å spectrum and up to four for the 0.33 and 1 Å spectra. For the 0.6 Å spectrum, lines detected down to the  $4\sigma$  level are presented.

The positions of the components of the double C IV doublet in the 5000–6000 Å spectral range were obtained by fitting the four lines together. The relative positions and widths of the lines of the same doublet were kept fixed. The results are given in Table 4G.

Figure 2a presents the normalized spectra of the Ly $\alpha$  forest with the identified lines. Figure 2b displays the C IV doublet detected in the 5000–6000 Å spectrum.

## 4. ANALYSIS OF THE LYMAN-ALPHA LINES

### 4.1. Selection of the Lyman-Alpha Lines in the 2 Å Spectra

Among the 2 Å resolution spectral lines (hereafter “2 Å lines”) we discriminated between lines which form blends (called hereafter “blended lines,” indicated by a B in the first column of Table 2) and the lines which remain single (called “single lines,” indicated by an S) when compared with the higher resolution (1, 0.6, or 0.33 Å) spectra.

Heavy-element systems were found at  $z = 1.9406, 1.9417, 1.9436, 1.9441, 2.3556, \text{ and } 2.3566$  (SBS; Paper II; this work),




FIG. 2a—Continued




FIG. 2a—Continued




FIG. 2b

TABLE 2  
LINE LISTS FOR 0.33, 0.6, 1, AND 2 Å RESOLUTION SPECTRA

1	2	B (2Å)				Δ	σ	A (2Å)				A (1Å)								
		λ	Wobs	σ	S/N			λ	Wobs	σ	S/N	λ	σ	Wobs	σ	S/N				
1	M	(3264.02	6.18	1.70	3.6)			3264.66	3.83	0.51	7.60									
2	M		4.22					3279.00	3.42	0.46	7.40									
3			4.35					3287.94	3.90	0.45	8.70									
4			4.09					3298.71	2.16	0.43	5.10									
5												3313.07	0.10	0.35	0.05	7.10				
6	S		2.52					3324.33	1.87	0.36	5.30	3324.33	0.13	1.80	0.11	17.10				
7												3329.25	0.07	0.40	0.05	8.00				
8												3331.43	0.22	0.73	0.12	6.00				
9												3337.69	0.04	0.42	0.07	6.00				
10												3342.93	0.04	0.84	0.07	11.90				
11												3346.63	0.08	0.43	0.09	5.00				
12	S		2.18					3356.08	1.68	0.28	6.00	3355.86	0.01	1.23	0.07	18.30				
13												3361.00	0.04	0.99	0.07	13.50				
14												3364.20	0.06	0.38	0.06	6.90				
15												3367.39	0.11	0.40	0.04	9.90				
16												3369.52	0.04	1.12	0.05	24.00				
17	B	(3372.93	3.65	1.05	3.5)			3374.02	3.02	0.25	12.10	3372.58	0.05	1.94	0.06	33.80				
18												3375.42	0.15	0.85	0.07	12.30				
19												3382.32	0.09	0.69	0.06	11.00				
20												3386.43	0.06	1.82	0.07	24.30				
21												3393.97	0.03	0.65	0.05	12.50				
22												3396.94	0.05	1.06	0.05	20.40				
23	B	L	1.56					3398.93	1.39	0.19	7.30	3398.94	0.05	0.34	0.05	7.30				
24												3402.58	0.06	0.77	0.06	13.20				
25												3407.02	0.02	1.17	0.05	21.50				
26	B	L	(3411.04	1.33	0.41	3.2)		3411.91	0.79	0.15	5.10	3410.18	0.05	1.10	0.06	20.00				
27												3413.58	0.06	0.23	0.04	5.70				
28												3417.29	0.08	0.42	0.03	13.50				
29												3418.74	0.38	1.04	0.08	12.70				
30	S		1.67					3437.61	1.47	0.20	7.30	3436.19	0.05	1.35	0.06	23.30				
31	S	M	3442.79	4.08	0.48	8.50	2.70	12.90	3443.11	4.45	0.20	22.60	3442.32	0.07	4.94	0.06	81.30			
32												3449.79	0.06	0.23	0.05	5.20				
33												3452.97	0.05	0.34	0.03	11.00				
34	B	L	1.61						3455.70	1.05	0.20	5.30	3454.81	0.06	0.85	0.05	15.80			
35	S	L	(3462.66	2.14	0.48	4.5)			3460.59	0.81	0.14	5.60	3459.65	0.04	0.59	0.02	24.90			
36	B	M	1.25						3463.78	2.00	0.26	7.80	3462.65	0.04	2.55	0.06	44.70			
37												3466.64	0.09	0.44	0.03	13.30				
38												3473.74	0.09	0.48	0.03	14.00				
39	B	M	1.36						3476.17	1.43	0.19	7.60	3476.76	0.11	1.34	0.07	19.30			
40												3480.72	0.03	1.11	0.03	35.60				
41	B	M	(3481.49	1.56	0.38	4.1)			3481.45	1.29	0.15	8.80	3482.85	0.04	1.19	0.05	25.80			
42	B	M	0.99						3484.66	1.70	0.20	8.50	3484.70	0.03	0.80	0.03	30.20			
43												3489.08	0.01	0.85	0.04	21.40				
44	S		(3501.75	1.43	0.36	4.0)			3502.13	1.45	0.14	10.60	3501.30	0.06	1.46	0.03	43.40			
45												3505.54	0.51	2.95	0.09	32.90				
46	B		(3506.85	0.93	0.29	3.2)			3506.70	2.56	0.32	8.10	3506.46	0.08	0.32	0.02	18.60			
47												3508.57	0.07	0.28	0.01	19.90				
48												3518.31	0.04	0.17	0.02	7.60				
49	B		1.22						3520.13	0.96	0.17	5.50	3520.73	0.03	0.99	0.06	17.20			
50												3538.77	0.04	0.15	0.02	6.80				
51												3540.73	0.05	0.35	0.04	8.10				
52	S	M	(3547.84	1.69	0.42	4.0)			3548.10	1.64	0.16	10.30	3548.17	0.05	1.91	0.04	48.40			
53	S	M	0.83						3551.73	1.76	0.19	9.10	3551.79	0.05	1.67	0.04	43.90			
54												3556.49	0.06	0.44	0.03	12.70				
55	S	L	(3564.27	1.68	0.37	4.5)			3564.13	1.22	0.12	9.70	3564.50	0.03	1.51	0.04	42.30			
56	S	L	(3568.38	0.86	0.27	3.1)			3567.82	1.80	0.16	11.40	3568.05	0.06	1.71	0.04	44.80			
57	B	M	3574.06	3.63	0.34	10.80	8.80	15.90	3574.24	3.77	0.16	24.10	3574.05	0.15	3.91	0.04	106.60			
58	B	M	3578.79	4.97	0.53	9.40	-3.10	17.10	3578.89	4.25	0.18	23.80	3578.55	0.19	4.38	0.05	95.90			
59												3581.12	0.06	0.66	0.01	54.00				
60												3585.50	0.05	0.35	0.03	11.30				
61	S		3589.70	2.20	0.33	6.80	-1.40	27.70	3589.36	1.90	0.13	14.40	3589.61	0.03	2.10	0.04	57.00			
62												3592.63	0.11	0.58	0.03	20.00				
63	B	L	(3594.62	1.20	0.33	3.6)			3594.89	1.06	0.13	7.90	3594.31	0.08	0.53	0.02	22.60			
64												3596.12	0.05	0.83	0.03	30.50				
65												3598.25	0.08	0.30	0.03	10.30				
66												3601.01	0.04	0.50	0.03	14.40				
67	B	L	(3604.34	1.19	0.29	4.2)			3603.46	0.70	0.13	5.50	3603.15	0.04	0.36	0.03	13.60			
68												3604.77	0.02	0.74	0.03	24.70				
69												3609.45	0.18	0.21	0.03	6.20				
70												3612.49	0.01	0.13	0.02	5.80				
71												3618.87	0.02	0.96	0.03	31.30				
72												3621.08	0.12	0.32	0.05	6.80				
73												3624.81	0.07	0.24	0.03	8.20				
74	S	L	0.86						3633.00	0.62	0.12	5.10	3633.41	0.05	0.76	0.03	22.90			
75												3636.75	0.04	0.51	0.03	16.70				

TABLE 2—Continued

I	2	B ( $\lambda$ )				A ( $\lambda$ )				A ( $\lambda$ )				A ( $0.3\lambda$ )				
		$\lambda$	Wobs	$\sigma$	S/N	$\lambda$	Wobs	$\sigma$	S/N	$\lambda$	Wobs	$\sigma$	S/N	$\lambda$	Wobs	$\sigma$	S/N	
76																		
77	S		0.76			3643.61	0.78	0.11	7.30	3638.69	0.03	0.48	14.50					
78	S	(3647.21	1.08	0.26	4.1)	3647.34	1.15	0.13	8.90	3644.01	0.01	1.16	0.03	37.40				
79	S	(3653.47	0.81	0.26	3.1)	3653.27	0.75	0.12	6.40	3654.05	0.06	1.37	0.04	36.90				
80										3658.55	0.08	1.12	0.02	45.20				
81	B	(3659.94	0.85	0.29	3.0)	3659.19	1.81	0.15	12.10	3662.08	0.15	0.32	0.02	16.50				
82										3665.53	0.02	0.73	0.03	22.50				
83										3672.41	0.05	0.23	0.03	7.70				
84										3675.60	0.07	0.38	0.03	11.80				
85										3679.70	0.05	2.32	0.03	74.20				
86	S	3679.12	1.48	0.23	6.40	3679.06	1.92	0.11	17.60	3683.72	0.08	1.09	0.03	31.60				
87	S	(3683.21	0.89	0.22	4.0)	3683.27	0.75	0.13	6.00	3688.91	0.02	0.69	0.03	23.90				
88										3695.50	0.04	0.27	0.03	9.40				
89	S		0.84			3696.97	0.93	0.12	7.50	3704.17	0.04	0.17	0.03	5.70				
90	L									3706.87	1.44	0.13	11.40					
91										3708.19	0.01	0.85	0.02	36.20				
92	B	M								3713.25	0.99	0.12	8.30					
93	S	(3712.82	1.16	0.25	4.7)	3713.25	0.89	0.12	8.30	3713.71	0.04	1.27	0.03	37.10				
94	S	(3720.88	1.53	0.18	8.70	3720.71	1.45	0.07	20.10	3720.74	0.08	1.58	0.02	89.90				
95	S	3726.36	6.06	0.50	12.20	3726.41	6.43	0.18	35.00	3726.53	0.11	7.03	0.06	124.60				
96	S									3732.95	0.04	0.52	0.03	18.80				
97										3735.50	0.08	0.66	0.03	20.00				
98	S		0.88			3738.75	0.75	0.13	5.70	3738.87	0.03	1.00	0.03	36.00				
99										3741.23	0.04	0.31	0.03	10.60				
100										3744.61	0.05	1.20	0.03	34.90				
101										3747.59	0.05	0.20	0.03	7.40				
102										3750.20	0.06	0.19	0.03	6.60				
103										3755.08	0.19	1.46	0.10	15.00				
104										3756.42	0.04	0.23	0.02	11.70				
105	B	3755.76	1.19	0.24	5.00	3756.11	0.60	0.09	6.90	3759.54	0.05	2.06	0.03	68.60				
106	S	3759.41	1.56	0.23	6.80	3759.47	1.69	0.14	12.30	3763.48	0.05	1.77	0.03	64.00				
107	B	3763.72	2.21	0.26	8.30	3763.41	1.76	0.12	14.30	3765.75	0.08	0.81	0.03	27.50				
108										3770.43	0.06	0.37	0.03	13.30				
109										3774.77	0.06	0.18	0.02	9.80				
110										3777.21	0.18	0.98	0.07	13.50				
111										3783.63	0.02	1.65	0.03	59.50				
112	S	(3782.57	0.44	0.14	3.2)	3783.35	0.72	0.07	10.70	3786.49	0.06	2.39	0.03	93.00				
113	B	3786.28	3.78	0.46	8.20	3786.28	3.33	0.19	17.60	3789.41	0.03	0.79	0.02	34.80				
114										3792.57	0.05	2.09	0.03	69.80				
115	S	3792.53	1.60	0.21	7.50	3792.65	1.91	0.10	18.80	3796.22	0.02	0.54	0.03	19.40				
116										3799.55	0.18	0.54	0.03	18.00				
117										3801.88	0.04	2.22	0.03	74.90				
118	B	3801.87	2.67	0.22	12.20	3801.71	2.23	0.11	21.10	3808.88	0.03	0.26	0.03	9.00				
119										3813.39	0.05	0.86	0.03	27.90				
120	S	(3813.70	0.82	0.17	4.9)	3813.74	1.04	0.11	9.80	3819.32	0.03	0.44	0.03	16.10				
121										3821.65	0.03	0.50	0.02	31.80				
122										3822.90	0.22	1.07	0.06	18.60				
123	B	3822.44	1.87	0.19	9.60	3822.24	2.35	0.14	16.60	3828.85	0.08	0.70	0.09	7.40				
124																		
125																		



TABLE 2—Continued

I	B (2Å)						A (2Å)						A(1Å)						A(0.3Å)																		
	2	λ	Wobs	σ	S/N	Δ	σ	λ	Wobs	σ	S/N	Δ	σ	λ	Wobs	σ	S/N	Δ	σ	λ	Wobs	σ	S/N	Δ	σ	λ	Wobs	σ	S/N								
126	S	M	3832.39	1.84	0.19	9.90	-4.70	9.00	3832.17	2.18	0.10	22.90	3831.51	0.02	1.31	0.02	75.50	3830.96	0.05	2.06	0.10	20.20															
127	B	M	3837.50	0.72	0.13	5.70	13.00	23.70	3836.75	1.22	0.10	11.70	3834.13	0.23	1.42	0.05	28.10	3833.01	0.11	1.57	0.15	10.60															
129	S	S		0.55					3844.57	0.59	0.09	6.20	3843.67	0.04	0.64	0.03	24.10	3844.15	0.06	0.64	0.06	10.50															
130																		3846.64	0.14	0.47	0.05	8.90															
131																		3850.35	0.10	0.42	0.06	6.60															
132	B	B	3856.50	1.45	0.20	7.40	59.00	27.10	3857.20	1.11	0.10	11.20	3855.21	0.08	0.38	0.02	16.70	3855.27	0.05	0.24	0.04	5.90															
133																		3857.62	0.03	0.73	0.08	9.00															
134																		3858.50	0.03	0.25	0.02	10.80															
135																		3867.12	0.03	0.71	0.02	15.40															
136	B	B	3868.72	1.81	0.19	9.60	6.60	8.40	3868.54	1.80	0.10	17.90	3868.92	0.02	0.98	0.02	53.00	3869.02	0.02	0.85	0.05	17.60															
137	S	S	3872.23	0.61	0.14	4.5)			3872.39	0.49	0.08	6.10	3872.02	0.03	0.60	0.02	26.30	3872.27	0.02	0.53	0.05	10.30															
138																		3876.52	0.06	0.43	0.02	19.20															
139	B	B	3877.86	0.81	0.18	4.4)			3878.52	0.91	0.11	8.40	3878.61	0.05	0.63	0.03	24.10	3878.86	0.13	0.62	0.07	8.50															
140																		3878.88	0.14	0.63	0.06	9.80															
141																		3898.60	0.13	1.00	0.11	9.1															
142	S	S	3911.22	1.45	0.21	7.00	0.50	5.40	3911.12	1.56	0.10	13.00	3900.23	0.04	0.26	0.03	8.20	3905.11	0.45	0.37	0.07	5.10															
143																		3906.37	0.04	0.59	0.03	18.00															
144	B	B	3921.39	1.18	0.19	6.10	1.30	28.40	3919.94	0.67	0.12	5.70	3911.16	0.02	1.75	0.03	56.40	3911.21	0.03	1.38	0.06	24.20															
145	S	S	3924.61	1.05	0.17	6.10	-5.80	24.20	3924.29	1.26	0.12	10.70	3921.01	0.08	1.22	0.03	45.00	3920.64	0.03	0.46	0.05	9.80															
146																		3924.34	0.04	1.16	0.06	19.10															
147																		3925.76	0.03	0.21	0.04	5.20															
148																		3927.02	0.07	0.25	0.04	6.30															
149																																					

TABLE 2—Continued

1	2	B (2Å)				Δ	A (2Å)				A(1Å)				A(0.6Å)				A(0.3Å)						
		λ	Wobs	σ	S/N		λ	Wobs	σ	S/N	λ	σ	Wobs	σ	S/N	λ	σ	Wobs	σ	S/N	λ	σ	Wobs	σ	S/N
151	S	3940.05	1.25	0.21	5.80	9.30	7.50	0.10	9.50	3940.19	0.04	1.42	0.03	43.80	3940.19	0.04	1.42	0.03	43.80	3946.40	0.03	0.26	0.05	5.60	
152										3946.71	0.03	0.37	0.03	14.60	3946.71	0.03	0.37	0.03	14.60	3952.09	0.11	0.60	0.07	8.60	
153										3950.33	0.23	0.80	0.04	22.20	3950.33	0.23	0.80	0.04	22.20	3954.88	0.05	1.42	0.09	16.10	
154	S	3954.79	1.28	0.19	6.70	32.30	27.50	0.12	13.50	3952.83	0.12	0.52	0.02	24.80	3952.83	0.12	0.52	0.02	24.80	3955.18	0.07	0.52	0.05	10.70	
155										3955.21	0.03	1.26	0.02	66.00	3955.21	0.03	1.26	0.02	66.00	3998.16	0.02	0.95	0.04	22.90	
156	B	3966.68	1.11	0.22	5.10	10.10	11.70	0.11	10.60	3964.72	0.12	0.32	0.02	16.00	3964.72	0.12	0.32	0.02	16.00	4000.03	0.04	0.25	0.03	8.50	
157										3966.95	0.06	1.08	0.03	37.10	3966.95	0.06	1.08	0.03	37.10	4001.45	0.04	0.21	0.03	7.70	
158										3972.83	0.02	0.25	0.02	12.10	3972.83	0.02	0.25	0.02	12.10	4004.32	0.05	0.24	0.02	10.50	
159										3974.95	0.01	0.51	0.02	24.10	3974.95	0.01	0.51	0.02	24.10	4005.62	0.09	0.73	0.05	13.30	
160										3977.06	0.05	0.22	0.03	8.40	3977.06	0.05	0.22	0.03	8.40	4009.92	0.01	0.25	0.03	8.70	
161	S									3982.53	0.02	0.77	0.03	24.10	3982.53	0.02	0.77	0.03	24.10	4016.85	0.13	0.37	0.04	8.20	
162										3988.17	0.02	1.02	0.03	37.50	3988.17	0.02	1.02	0.03	37.50	4022.97	0.17	0.60	0.06	9.70	
163	S	3991.91	1.15	0.20	5.90	-9.00	3.70	0.09	8.80	3991.82	0.01	1.05	0.02	43.80	3991.82	0.01	1.05	0.02	43.80	4028.09	0.02	0.83	0.07	12.30	
164										3995.11	0.07	0.53	0.03	19.70	3995.11	0.07	0.53	0.03	19.70						
165	B									3998.42	0.03	0.82	0.02	48.80	3998.42	0.03	0.82	0.02	48.80						
166										4000.93	0.23	0.48	0.04	11.80	4000.93	0.23	0.48	0.04	11.80						
167										4005.51	0.05	0.83	0.03	27.10	4005.51	0.05	0.83	0.03	27.10						
168	B									4010.19	0.06	0.17	0.02	8.50	4010.19	0.06	0.17	0.02	8.50						
169																									
170																									
171																									
172	S	(4028.63	0.90	0.20	4.6)			0.09	8.00	4028.09	0.03	0.79	0.03	30.90	4028.09	0.03	0.79	0.03	30.90						
173																									
174																									
175																									
176																									
177																									
178																									
179	B	4040.98	3.58	0.29	12.40	-23.90	31.10	0.15	25.50	4034.68	0.12	0.36	0.03	14.30	4034.68	0.12	0.36	0.03	14.30						
180										4037.85	0.05	1.28	0.02	57.00	4037.85	0.05	1.28	0.02	57.00						
181										4040.05	0.03	1.44	0.02	78.20	4040.05	0.03	1.44	0.02	78.20						
182										4042.35	0.03	1.14	0.02	56.70	4042.35	0.03	1.14	0.02	56.70						
183	B+	4049.27	2.70	0.28	9.80			0.11	22.40	4044.57	0.03	0.68	0.02	28.70	4044.57	0.03	0.68	0.02	28.70						
184										4049.39	0.05	2.75	0.04	76.70	4049.39	0.05	2.75	0.04	76.70						
185	S							0.09	7.90	4052.75	0.25	0.23	0.02	12.00	4052.75	0.25	0.23	0.02	12.00						
186										4058.08	0.02	0.67	0.03	25.50	4058.08	0.02	0.67	0.03	25.50						
187																									
188																									
189	B	4065.49	1.30	0.18	7.10	41.00	65.40	0.10	12.80	4063.05	0.04	0.19	0.02	8.50	4063.05	0.04	0.19	0.02	8.50						
190										4066.24	0.05	1.07	0.03	42.40	4066.24	0.05	1.07	0.03	42.40						
191										4068.50	0.09	0.53	0.02	22.50	4068.50	0.09	0.53	0.02	22.50						
192																									
193																									
194																									
195	B+	4080.62	6.60	0.29	22.90	21.00	4.40	0.12	52.30	4074.28	0.09	0.47	0.03	17.30	4074.28	0.09	0.47	0.03	17.30						
196										4081.14	0.11	6.31	0.03	183.30	4081.14	0.11	6.31	0.03	183.30						
197										4087.27	0.04	0.61	0.03	21.70	4087.27	0.04	0.61	0.03	21.70						
198	B	4092.56	1.23	0.17	7.20	37.90	32.60	0.09	15.00	4092.56	0.03	1.56	0.02	66.70	4092.56	0.03	1.56	0.02	66.70						
199	B	4096.78	3.84	0.39	9.80			0.16	23.40	4095.95	0.10	1.87	0.02	101.90	4095.95	0.10	1.87	0.02	101.90						
200																									

TABLE 2—Continued

1	2	B (2Å)				A (2Å)				A (1Å)				A (0.6Å)				A (0.3Å)									
		λ	Wobs	σ	S/N	Δ	λ	Wobs	σ	S/N	λ	Wobs	σ	S/N	λ	Wobs	σ	S/N	λ	Wobs	σ	S/N					
201																											
202																											
203																											
204																											
205	B	4108.56	2.41	0.22	10.70	0.10	7.90	4108.59	1.94	0.10	20.00	4107.96	0.05	0.95	0.02	41.90	4109.05	0.07	1.09	0.08	13.40	4098.21	0.05	1.73	0.05	35.20	
206	B	4117.70	1.96	0.23	8.52	-1.30	15.40	4117.78	2.30	0.10	23.80	4113.10	0.11	0.29	0.03	51.70	4109.75	0.05	1.04	0.08	13.70	4099.67	0.04	0.40	0.04	9.90	
207	B											4119.96	0.04	1.10	0.02	52.50	4119.82	0.05	1.34	0.04	33.10						
208												4123.40	0.11	1.22	0.05	26.10	4121.68	0.03	0.60	0.02	27.10	4119.88	0.04	0.86	0.05	17.60	
209												4121.53	0.24	0.82	0.03	29.00	4124.91	0.06	1.22	0.10	12.20	4121.69	0.05	0.72	0.03	26.70	
210	B	4122.59	4.13	0.47	8.81	44.80	54.90	4123.17	3.68	0.19	19.50	4124.10	0.42	0.82	0.03	18.60	4122.86	0.04	0.73	0.03	29.20	4123.20	0.06	0.76	0.03	22.50	
211												4126.03	0.69	1.28	0.04	34.90	4126.54	0.05	0.26	0.05	5.40	4125.10	0.08	0.91	0.04	23.50	
212												4130.64	0.02	1.27	0.02	53.00	4130.65	0.04	0.85	0.10	8.70	4126.53	0.09	0.52	0.03	19.00	
213	S	4130.56	1.23	0.18	6.80	-8.00	15.90	4129.81	1.16	0.09	12.30	4136.14	0.02	0.47	0.03	16.90	4136.11	0.10	0.60	0.10	5.90	4130.84	0.01	1.16	0.03	34.00	
214	S											4141.69	0.05	0.33	0.03	12.40	4140.67	0.11	0.26	0.04	6.50	4136.59	0.04	0.63	0.03	18.10	
215																						4142.01	0.03	0.39	0.02	17.30	
216																						4147.36	0.04	0.13	0.02	6.00	
217																											
218	B	4153.01	2.24	0.19	11.90	0.30	27.90	4152.72	2.10	0.10	20.40	4150.15	0.11	0.59	0.03	19.80	4150.15	0.11	0.59	0.03	19.80	4152.87	0.02	1.20	0.03	41.70	
219												4154.34	0.02	1.18	0.02	49.50	4154.11	0.06	0.09	0.02	4.90	4154.29	0.02	0.87	0.02	37.20	
220	S	4155.96	1.89	0.21	9.10	-14.10	24.80	4155.90	1.25	0.09	14.60	4156.24	0.06	1.17	0.05	23.50	4156.24	0.06	1.17	0.05	23.50	4156.55	0.01	1.19	0.03	38.70	
221																											
222																											
223																											
224	S											4166.23	0.11	0.36	0.03	12.60	4170.58	0.09	0.48	0.09	5.40	4166.38	0.09	0.32	0.03	11.00	
225												4170.86	0.04	0.92	0.03	31.20	4170.58	0.12	0.44	0.05	8.10	4170.66	0.04	1.10	0.04	26.90	
226	B	4183.46	1.68	0.21	7.90	15.40	18.00	4182.77	1.57	0.11	14.00	4180.61	0.12	0.44	0.05	8.10	4180.61	0.12	0.44	0.05	8.10	4182.25	0.11	0.63	0.05	11.40	
227												4184.01	0.28	0.94	0.03	32.60	4183.98	0.01	0.47	0.06	8.40	4184.00	0.03	1.06	0.04	30.40	
228	S	(4194.19	0.98	0.21	4.6)			4193.86	0.40	0.07	5.70	4194.06	0.04	0.64	0.02	31.60	4193.74	0.05	0.42	0.10	4.00	4194.00	0.03	1.06	0.04	30.40	
229	S							4196.61	0.65	0.11	6.00	4196.88	0.04	0.50	0.08	5.90	4196.88	0.04	0.50	0.08	5.90	4197.22	0.03	0.41	0.03	14.40	
230												4208.33	0.04	0.45	0.08	5.90	4208.33	0.04	0.45	0.08	5.90	4208.57	0.02	0.55	0.02	24.70	
231	B	4209.42	1.42	0.20	7.30	-0.10	4.50	4209.54	1.33	0.10	12.90	4209.89	0.03	0.91	0.02	11.40	4209.89	0.03	0.91	0.02	11.40	4210.12	0.01	1.14	0.03	43.80	
232												4219.29	0.09	0.22	0.05	4.50	4219.29	0.09	0.22	0.05	4.50						
233												4222.62	0.04	0.42	0.06	7.60	4222.62	0.04	0.42	0.06	7.60	4223.19	0.11	0.21	0.02	10.10	
234												4224.44	0.04	0.28	0.05	5.90	4224.44	0.04	0.28	0.05	5.90	4224.59	0.01	0.23	0.02	11.50	
235												4227.71	0.08	1.17	0.10	11.30	4227.71	0.08	1.17	0.10	11.30	4228.22	0.02	1.43	0.03	46.90	
236	B	4229.34	2.13	0.19	11.10	-9.00	16.80	4229.10	2.62	0.10	26.90	4229.70	0.13	1.05	0.14	7.70	4229.70	0.13	1.05	0.14	7.70	4229.76	0.02	0.63	0.02	38.00	
237																											
238												4232.30	0.07	0.41	0.06	7.00	4232.30	0.07	0.41	0.06	7.00	4232.57	0.02	0.27	0.02	12.90	
239												4234.48	0.07	1.57	0.09	17.30	4234.48	0.07	1.57	0.09	17.30	4234.75	0.03	2.03	0.04	48.10	
240	S	4235.23	3.16	0.19	16.60	16.40	26.70	4235.02	3.44	0.11	32.50	4236.02	0.05	0.84	0.05	16.90	4236.02	0.05	0.84	0.05	16.90	4236.29	0.02	0.81	0.02	38.10	
241												4237.90	0.03	0.23	0.04	5.70	4237.90	0.03	0.23	0.04	5.70						
242												4241.52	0.06	0.16	0.03	5.50	4241.52	0.06	0.16	0.03	5.50						
243																											
244	B	4244.70	1.98	0.20	9.90	-3.40	9.20	4244.67	1.83	0.09	19.80	4244.32	0.02	2.33	0.12	19.70	4244.32	0.02	2.33	0.12	19.70	4243.96	0.04	0.66	0.03	24.40	
245	S	(4250.45	0.86	0.18	4.8)			4250.67	0.96	0.10	10.00	4250.27	0.07	0.98	0.12	7.90	4250.27	0.07	0.98	0.12	7.90	4250.39	0.02	0.89	0.04	23.80	
246	S	4254.54	1.53	0.17	8.90	-2.10	2.20	4254.52	1.55	0.08	19.10	4254.21	0.05	1.59	0.07	21.70	4254.21	0.05	1.59	0.07	21.70	4254.25	0.02	1.78	0.04	40.20	
247	B	4259.48	1.27	0.18	7.20	-11.60	3.00	4259.71	1.68	0.10	16.90	4259.58	0.04	1.08	0.12	8.80	4259.58	0.04	1.08	0.12	8.80	4259.54	0.02	1.22	0.06	19.90	
248												4262.02	0.05	0.33	0.07	4.70	4262.02	0.05	0.33	0.07	4.70						
249												4264.98	0.06	0.26	0.04	6.90	4264.98	0.06	0.26	0.04	6.90						
250												4267.64	0.06	0.06	0.05	7.10	4267.64	0.06	0.06	0.05	7.10						





using mainly the higher resolution spectra and the identification list of Morton, York, & Jenkins (1988). They are presented in Table 4 and are analyzed independently (cf. § 5).

It also appears that the two reported anticoincidences at 3987 and 3998 Å are compatible with the Mg II  $\lambda\lambda 2796$  and 2802 doublet at a redshift  $z = 0.4261$ . A search for other lines at the same redshift leads to the possible identification of Mg I  $\lambda 2852$  at 4068 Å, but no other line was found. If we consider the two anticoincidences as due to Ly $\alpha$  lines, and if we take into account the fact that they are the only anticoincidences, it is a posteriori quite unexpected to find them so close to each other and with their relative positions and strengths similar to those expected for a Mg II doublet. However, this hypothesis cannot be totally excluded: the expected number of Ly $\alpha$  lines which fulfill these conditions within a measurement uncertainty of 0.1 Å (corresponding to an observed  $10^{-4}$  scatter in the determination of the redshift of heavy-element systems) is about  $7.5 \times 10^{-4}$ . On the other hand, we could hope to detect the responsible galaxy at such a redshift. The analysis of a 30 minute *R* image used by Magain et al. (1990) shows a diffuse feature near the quasar: its identification as the responsible object remains in doubt, since it lies closer to B than to A, and since no spectrum had been taken yet. The galaxy D (cf. Paper I) which appears south of UM 673 is not involved, since its redshift has been measured to be 0.17 (Surdej 1990).

In the higher resolution spectra, a line is interpreted as a Ly $\beta$  line if its position relative to an already selected Ly $\alpha$  line, with equivalent width  $W_{Ly\alpha}$ , is similar within the measurement uncertainty to that expected from the wavelength ratio, and if its equivalent width  $W < W_{Ly\alpha} + 3\sigma(W_{Ly\alpha})$ . We have also applied the corresponding criterion to the suspected Ly $\gamma$  lines with respect to the accepted Ly $\beta$  lines.

If any line in the high-resolution spectra (including Ly $\alpha$ ) of one of the heavy-element systems can be interpreted as a sub-component of a blended line (in the 2 Å spectra) or is the corresponding line of a single line (in the 2 Å spectra), this (blended or single) line is marked M in Table 2 (second column) and is not considered as a possible Ly $\alpha$  (cloud) line; this is quite restrictive, as it may happen that the equivalent width of a heavy-element system line in the high-resolution spectra is very small compared with the corresponding blended line. The same rule was applied for the Ly $\beta$  and Ly $\gamma$  lines. However, in this case, the involved 2 Å lines are indicated by an L in Table 2.

Since there remains some doubt on the reality of the anticoincidences as Ly $\alpha$  lines, we shall consider in what follows two distinct cases. Case 1: our sample of Ly $\alpha$  lines contains 47 coincidences i.e., 21 single lines and 26 blended lines, and two anticoincidences. Case 2: we have 46 coincidences, i.e., 20 single lines and 26 blended lines, and no anticoincidences.

## 4.2. Diameter of Lyman-Alpha Clouds

### 4.2.1. Separations between the Light Beams

By definition of the angular-diameter distance, the comoving linear separation  $S(z_d)$  at the redshift of the lens is given by  $S(z_d) = \theta D_{od}$  and also by  $S(z_d) = \theta' D_{sd}$ , where  $\theta'$  is the angle between the light beams at the source position (cf. Fig. 3 for the geometry of this gravitational lens system). Simple geometrical considerations give the proper separation  $S(z_i)$ , for a cloud  $i$  at redshift  $z_i$ , between the light beams:<sup>10</sup>

$$S(z_i) = \theta \frac{D_{od} D_{sc}(z_i)}{D_{sd}} \quad (1)$$

where  $\theta$  is the angular separation between the A and B images of UM 673,  $D_{od}$  is the angular diameter distance between the observer and the deflector,  $D_{sc}(z_i)$  is the angular-diameter distance between the source and the cloud at redshift  $z_i$ , and  $D_{sd}$  is the angular-diameter distance between the source and the deflector.

A general formula to compute these angular-diameter distances is given in Blandford & Kochanek (1987):

$$D_{ij} = \frac{2c}{H_0} \frac{(1 - 2q_0)(G_i - G_j) + (G_i G_j^2 - G_j^2 G_i)}{(2q_0)^2(1 + z_i)(1 + z_j)^2}, \quad (2)$$

where  $G_i = (1 + 2q_0 z_i)^{1/2}$ .

$D_{sc}$  and  $D_{sd}$  are related to  $D_{cs}$  and  $D_{ds}$ , respectively, by

$$D_{sc} = D_{cs} \frac{1 + z_s}{1 + z_c} \quad (3)$$

and

$$D_{sd} = D_{ds} \frac{1 + z_s}{1 + z_d}. \quad (4)$$

For UM 673 the linear separation between the light beams is represented as a function of redshift in Figure 4.

### 4.2.2. Correlation of the Equivalent Widths

Figure 5a plots the rest equivalent width ( $W_B$ ) of Ly $\alpha$  lines in spectrum B versus that ( $W_A$ ) for the corresponding line in spectrum A; for the 21 single lines (*solid-line symbols*) and the 26

<sup>10</sup> Note that, using  $S(z_d) = \theta' D_{ds}$ , instead of  $S(z_d) = \theta' D_{sd}$ , Foltz et al. (1984) have obtained the formula  $S = \theta D_{od} D_{cs}(z_i)/D_{ds}$ , which is then wrong by a factor  $(1 + z_d)/(1 + z_s)$ : the published values for the separation between the light beams of images A and B of Q2345+007 at  $z_a = 1.951$  should then be approximately multiplied by 0.5, 0.7, and 0.9 for  $z_d = 0.5, 1,$  and  $1.5,$  respectively. Basically, the same error has been made Young et al. (1980, 1981): the separation between the light beams for the absorption systems at  $z_a = 1.1249, 1.3911$  in the spectra of Q0957+561 should be approximately multiplied by 0.5 and 0.4, respectively.

NOTE.—Columns are arranged for easy comparison with the corresponding spectra. A “B” in the first column means that the lines in the 2 Å spectra are blended when comparison is made with the higher resolution spectra; otherwise an “S” is indicated. An “M” (“L”) in the second column means that the line—or at least one of the subcomponents in the higher resolution spectra, if the line is blended—belongs to a metallic system (can be a Ly $\beta$  or Ly $\gamma$  line). A “B+” in the first column means that the line is taken as blended most probably because of a bad sky subtraction in the 0.6 Å spectrum. For the 2 Å spectra, central wavelengths ( $\lambda$ ), observed equivalent width ( $W_{obs}$ ), formal error on the equivalent widths ( $\sigma$ ), and signal-to-noise ( $= W_{obs}/\sigma$ ) for the absorption lines are given. Difference in velocities ( $\Delta$ ) and associated formal errors ( $\sigma$ ) in kilometers per second are also given for the single and blended coincidence lines. For the other spectra central wavelengths ( $\lambda$ ) and associated fitting errors ( $\sigma$ ), observed equivalent widths ( $W_{obs}$ ), formal error on equivalent widths ( $\sigma$ ), and signal-to-noise ratios ( $= W_{obs}/\sigma$ ) are given. For the B image, characteristics within parentheses mean that the line is detected at only the  $3\sigma$  level. If only the equivalent width is given, the value is a  $3\sigma$  upper limit. All wavelengths are heliocentric vacuum values. Wavelengths and equivalent widths are in angstroms. (The question mark for the line at 4096 Å means that the cross-correlation function presents two peaks of nearly equal intensity corresponding to  $\Delta = -70.5 \pm 15.8$  km s<sup>-1</sup> and  $\Delta = 40.3 \pm 23.4$  km s<sup>-1</sup>.)




FIG. 3.—Geometry of a gravitational lens system. Symbols are defined in the text.

blended lines (*dashed-line symbols*). In all cases, the correlation is excellent. The dispersion is consistent with that expected from measurement errors alone. In particular, it is reasonable to believe that (1) the number of components among the blended lines is the same in the spectrum of B as in that of A and (2) the individual components of each blended line are also very well correlated in the two spectra. Figure 5b shows  $W_B$  versus  $W_A$  for the B lines detected at the level of  $3\sigma$  to  $5\sigma$ : most of them do follow the general correlation between  $W_A$  and  $W_B$ . This figure presents also the  $3\sigma$  upper limit for the rest equivalent width of the A lines which are “missing” in the B spectrum. Here also, most of them are compatible with the general correlation.

We consider that the remarkable correlation existing between  $W_A$  and  $W_B$  is a proof that both light beams actually cross the same Ly $\alpha$  clouds (assuming their intergalactic nature; SYBT). If small Ly $\alpha$  clouds were clustered, as required, for




FIG. 4.—Separation between the two beams as a function of redshift for the gravitational lens system UM 673.

example, by the models of Tytler or Pettini et al., the two lines of sight would have crossed *different* clouds at the same redshift, since their size ( $0.3$  or  $10^{-5}$  pc) would be very small compared with the separation between the light beams (typically 1 kpc), and no correlation between the measured equivalent widths would be expected.

Thin sheets are still possible, but they would have to have axial ratios of  $10^4$ – $10^8$  and be homogeneous on scales of  $3h_{50}^{-1}$  pc to  $2000h_{50}^{-1}$  pc (the separation between the light beams for the extreme Ly $\alpha$  clouds of our sample).

There is also strong evidence that the equivalent widths of the Ly $\alpha$  lines, as given by Foltz et al. (1984) correlate also in the case of Q2345 + 007, as shown in Figure 6. In consequence, the scale values given above should be increased up to  $22h_{50}^{-1}$  kpc (if the deflector redshift is 1.5), and another order of magnitude should be added to the axial ratios in the case of thin sheets.

4.2.3. Limits on the Diameter of the Lyman-Alpha Clouds: Monte Carlo Simulations

In this section we shall compare the number of anti-coincidences with the total number of lines in our sample in order to put some constraints on the size of the Ly $\alpha$  clouds. We shall then try to use the information given by the equivalent widths of the lines to set constraints on the possible structure of Ly $\alpha$  clouds.

In the following we shall use a simple model: we assume that the clouds are all identical (no evolution) and spherical.

First, we define the ratio  $f$  between the number of anti-coincidences and the sum  $N$  of the number of anti-coincidences and the number of coincidences, for different values of the diameter,  $D$ . We can then compare the observational results with those from Monte Carlo simulations, whose description follows:

a) We use the observed distribution of Ly $\alpha$  clouds along the line of sight, in the sense that we put spherical clouds at each of the redshifts derived for the (single) lines in the observed spectra.

b) Working in the plane perpendicular to the line of sight, we compute  $f$  for each assumed value of  $R_c = D/2$  in the following way (see also Fig. 7): (i) the projection of a spherical cloud on a plane perpendicular to the line of sight is a disk: we consider the center of this disk to be the origin of the coordinates; (ii) for each Ly $\alpha$  line, the separation  $S(z_i)$  between the light beams is computed using equation (1); (iii) two points,  $r_{A,i}$  and  $r_{B,i}$ , separated by  $S(z_i)$  are randomly chosen on a region covering a disk of radius at least equal to  $R_c + S(z_i)$ , such that their distributions are uniform on this disk, and the corresponding impact parameters  $r_{A,i}$  and  $r_{B,i}$  are computed; (iv)  $r_{A,i}$  and  $r_{B,i}$  are then compared with  $R_c$ : (1) if only one of  $r_{A,i}$  or  $r_{B,i}$  is greater than  $R_c$ , the number of anti-coincidences is increased by one; (2) if  $r_{A,i}$  and  $r_{B,i}$  are smaller than  $R_c$ , the number of coincidences is increased by one; (3) if both  $r_{A,i}$  and  $r_{B,i}$  are greater than  $R_c$ , the trial is not counted at all.

We also made similar simulations for an oblate spheroid, given by the equation

$$\frac{x^2 + y^2}{a^2} + \frac{z^2}{b^2} = 1, \tag{5}$$

without any preferential direction for any axes. In particular, if  $b/a$  is very small, we have a simple model for a circular sheet.

For  $N$  lines, this process is repeated a sufficient number of times in order to get accurate results for  $f$  and the associated




FIG. 5a




FIG. 5b




FIG. 5c




FIG. 5d

FIG. 5.—Correlation between rest equivalent widths of coincidence lines in the  $2 \text{ \AA}$  spectra, (a) for the 21 single (solid-line symbols) and the 26 blended Ly $\alpha$  lines (dashed-line symbols); (b) for the  $3 \sigma$  level detected lines in spectrum B and the  $5 \sigma$  level detected lines in spectrum A, and the  $3 \sigma$  upper limit on  $W_B$  for the “missing” lines; (c) the  $z_{\text{abs}} = 1.94$  heavy-element complex, including the upper limit on  $W_B$  for the missing line in spectrum B; (d) the  $z_{\text{abs}} = 2.35$  heavy-element complex, including the upper limit on  $W_B$  for the “missing” lines in spectrum B. All equivalent widths are in angstroms.

errors. The choice of the line sets has two effects: first, generally, if  $N$  increases, the errors get smaller; second, since we use the observed lines to obtain the distribution of the clouds in our simulations, a different set of lines (but with the same value for  $N$ ) will usually produce different results. However, adding more clouds close to the quasar does not affect the results significantly if the separation between the light beams is much smaller than the size of the clouds.

Some results from these simulations are presented in Figure 8 as plots of  $f + 2\sigma$  and  $f - 2\sigma$  versus  $D$ . In turn, for a given

observed value  $f_{\text{obs}}$ , we obtain  $2 \sigma$  lower and upper limits for  $D$ . The whole set of results is discussed below and summarized in Table 3, for different samples and for the two cases.

For case 1, i.e., that the two anticoincidences are Ly $\alpha$  lines, we have the following:

a) Sample 1 is constituted from the isolated lines alone. It forms the most secure sample: its 21 coincidence lines and one anticoincidence line are not affected by blending, and therefore there is a one-to-one correspondence between the lines and




FIG. 6.— $W_B$  vs.  $W_A$  for Q2345+007 images A and B. Rest equivalent widths in angstroms.

their equivalent widths in the high-resolution spectra and the  $2 \text{ \AA}$  spectrum of A. We obtain a  $2 \sigma$  lower limit of  $8h_{50}^{-1}$  kpc and a  $2 \sigma$  upper limit of  $280h_{50}^{-1}$  kpc.

b) Sample 2 contains the 21 isolated and 26 blended coincidence lines, and the one isolated and one blended anticoincidence line. We consider the results ( $12h_{50}^{-1}$  kpc for the  $2 \sigma$  lower limit and  $160h_{50}^{-1}$  kpc for the  $2 \sigma$  upper limit) as the best (the most convincing) values we can obtain with our data: we use the blended lines in the same way as for the single ones. This is equivalent to saying that (i) one subcomponent of a blended line is surely present in the A and in the B spectra, and (ii) nothing is known about the detection of the other ones, for the B spectrum signal-to-noise ratio is not as good as that for the A spectrum.

c) Sample 3 is an attempt to use all the information of the high-resolution spectra. To the 21 single lines, we have added all the subcomponents of the blended lines whose equivalent




FIG. 7.—Schematic diagram related to the Monte Carlo simulations; the symbols are described in the text.




FIG. 8.—Plot of  $f - 2\sigma$  and  $f + 2\sigma$  vs.  $D$  for samples 1, 2, and 3. The horizontal lines mark the observed values of  $f$  for the three samples; their intersections with the curves give the  $2 \sigma$  limits on the cloud sizes.

widths (as measured in the high-resolution spectra) are bigger than twice the formal errors on the equivalent widths of the corresponding B lines. With such a criterion, we use as many lines as possible, while we do not take into account the very low equivalent width lines whose absence is unnoticed because of the noise of the  $2 \text{ \AA}$  spectra. We obtain 62 coincidence lines, and we assume that there are two anticoincidences. The  $2 \sigma$  lower and upper limits are therefore  $21h_{50}^{-1}$  kpc and  $200h_{50}^{-1}$  kpc.

d) Sample 4 is the same as sample 2, but for oblate spheroidal clouds. We find that the results—a  $2 \sigma$  lower and upper limit for  $2a$  of  $20h_{50}^{-1}$  kpc and  $300h_{50}^{-1}$  kpc, respectively—are independent of  $b/a$  if this ratio is less than 0.1, owing to the relatively small number of lines. These values are about twice those obtained for a spherical Ly $\alpha$  cloud.

e) Sample 5 comes from the results of Foltz et al. (1984). From the 19 lines published, we removed lines 8, 11, and 12 because of Note 5 of their table, while line 19 has a larger redshift than that assumed for the quasar and, therefore, cannot be used in our model. We are left with 13 coincidence and two anticoincidence lines. We assume a redshift of 0.5 for the lens. Although we make a better use of the sample, we obtain only a slightly larger value for the  $2 \sigma$  lower limit ( $8h_{50}^{-1}$  kpc instead of  $5h_{50}^{-1}$  kpc) because of their error on the formula giving the separation between the two lines of sight, mentioned in § 4.2.1. However, we can also set a  $2 \sigma$  upper limit of  $104h_{50}^{-1}$  kpc.

f) Sample 6 is the combination of samples 2 and 5. Compared with sample 5, this increases significantly the  $2 \sigma$  lower limit to  $13h_{50}^{-1}$  kpc, while the  $2 \sigma$  upper limit decreases to  $92h_{50}^{-1}$  kpc.

g) Samples 7 and 8 (and samples 9 and 10) are obtained with the same data set as for samples 5 and 6, except that the redshift for the lens in the Q2345+007 system is assumed to be  $z = 1$  ( $z = 1.5$ ). Comparing samples 2 and 8, we see that the Q2345+007 data help to improve the  $2 \sigma$  lower limit to  $24h_{50}^{-1}$  kpc, but not the upper limit (which stays at about  $160h_{50}^{-1}$  kpc). When samples 2 and 10 are compared, we find that the  $2 \sigma$  lower limit is increased to  $54h_{50}^{-1}$  kpc. However, the  $2 \sigma$  upper limit is significantly less restrictive in sample 10 ( $360h_{50}^{-1}$  kpc) than in sample 2 ( $160h_{50}^{-1}$  kpc).



TABLE 3  
 $2\sigma$  LIMITS ON THE SIZE OF LYMAN-ALPHA CLOUDS

CASE 1: ANTICOINCIDENCES ARE $\text{Ly}\alpha$ LINES				CASE 2: ANTICOINCIDENCES ARE A $\text{Mg II}$ DOUBLET			
Sample Number	Description	$f$	Diameter ( $h_{50}^{-1}$ kpc)	Sample Number	Description	$f$	Diameter ( $h_{50}^{-1}$ kpc)
A. Samples from UM 673 Only							
1	21 single lines; 1 anticoincidence	0.045	$8 < D < 280$	13	20 single lines; no anticoincidence	0	...
2	26 blended lines; 21 single lines; 2 anticoincidence	0.041	$12 < D < 160$	14	24 blended lines; 20 single lines; no anticoincidence	0	$23 < D$
3	62 subcomponents; 2 anticoincidence	0.031	$21 < D < 200$	15	61 subcomponents; no anticoincidence	0	$28 < D$
4	As sample 2; spheroidal clouds	0.041	$20 < 2a < 300$	16	As sample 14; spheroidal clouds	0	$40 < 2a$
B. Samples from Q2345+007 Only or with UM 673							
5	13 lines; 2 anticoincidence; $z_D(\text{Q2345+007}) = 0.5$	0.133	$8 < D < 104$	...	...	...	...
6	Sample 2 + sample 5	0.062	$13 < D < 92$	17	Sample 14 + sample 5	0.034	$20 < D < 260$
7	13 lines; 2 anticoincidence; $z_D(\text{Q2345+007}) = 1.0$	0.133	$26 < D < 330$	...	...	...	...
8	Sample 2 + sample 7	0.062	$24 < D < 162$	18	Sample 14 + sample 7	0.034	$35 < D < 480$
9	13 lines; 2 anticoincidence; $z_D(\text{Q2345+007}) = 1.5$	0.133	$62 < D < 770$	...	...	...	...
10	Sample 2 + sample 9	0.062	$54 < D < 360$	19	Sample 14 + sample 9	0.034	$80 < D < 1000$
11	13 lines; 2 anticoincidence; Q2345+007 is not a GL	0.133	$500 < D < 6500$	...	...	...	...
12	Sample 2 + sample 11	0.062	$410 < D < 3400$	20	Sample 14 + sample 11	0.034	$380 < D$

h) Samples 11 and 12 are the same as sample 5 and 6, except that we suppose that Q2345+007 is not a gravitational lens system, but a real quasar pair as proposed by Steidel & Sargent (1990). With the same data, we obtain a  $2\sigma$  lower limit of  $500h_{50}^{-1}$  kpc, instead of  $100h_{50}^{-1}$  kpc as they obtained in their paper, for the reason explained in paragraph *e* above. However, the comparison between samples 2 and 12 shows that the results are incompatible: the  $2\sigma$  upper limit obtained with sample 2 is much smaller ( $160h_{50}^{-1}$  kpc) than the  $2\sigma$  lower limit given in sample 12 ( $410h_{50}^{-1}$  kpc).

We conclude that if the two anticoincidence are indeed  $\text{Ly}\alpha$  lines, the results from UM 673 alone are not compatible with the results obtained under the hypothesis that Q2345+007 is a binary quasar. In this case, the size of  $\text{Ly}\alpha$  clouds is small enough so that two relatively close, nearly parallel light paths, as in the case of a binary quasar, would cross different  $\text{Ly}\alpha$  clouds. Therefore, study of the  $\text{Ly}\alpha$  forest can help to discriminate between the binary quasar and the gravitational lens explanations for close images of high-redshift quasars.

We present also the results for case 2, in which the anticoincidence are in fact due to a  $\text{Mg II}$  doublet, with a  $\text{Mg I}$  line detected at the same  $z$ . The major differences are that the  $2\sigma$  lower limits are larger and that no upper limits can be set. In conclusion, if the two anticoincidence are caused by a  $\text{Mg II}$  doublet—in fact, even if one of the anticoincidence is not a  $\text{Ly}\alpha$  line but a metallic line in a not-yet-identified system—we

are unable to give any meaningful  $2\sigma$  upper limit for the diameter of the  $\text{Ly}\alpha$  clouds, using the results from UM 673 alone. As a consequence, they are compatible with any of the possible hypotheses as to the origin of Q2345+007. In particular, in the case of the binary quasar explanation, we are led to a  $2\sigma$  lower limit for the  $\text{Ly}\alpha$  cloud diameter of the order of  $400h_{50}^{-1}$  kpc.

#### 4.2.4. Structure of the Lyman-Alpha Clouds

The fact that the equivalent widths of absorption lines detected in the two spectra correlate proves that  $\text{Ly}\alpha$  clouds are structured, in the sense that the column density and the velocity dispersion at a given impact parameter set important constraints on these quantities at another one. In the case of UM 673, the column densities and the velocity dispersion must have very similar values for impact parameter differences between  $0h_{50}^{-1}$  and  $2h_{50}^{-1}$  kpc, except if in each case—corresponding to rest equivalent widths ranging from 0.2 to 2 Å—the variation of one of the parameters is well counter-balanced by the appropriate change of the other parameter (that is very unlikely).

On the other hand, nearly any density profile in a cloud produces a variation of the column density with the impact parameter, as recalled by Milgrom (1988). The fact that the equivalent widths of the two anticoincidence are among the lowest in our sample (assuming they are  $\text{Ly}\alpha$  lines) suggests actually that  $\text{Ly}\alpha$  clouds are not of uniform density.

The way the equivalent widths of absorption lines detected in the two spectra correlate gives us a tool to study the structure of the Ly $\alpha$  clouds, as we shall demonstrate in this section by using a simple model: we assume that the Ly $\alpha$  clouds are described by a singular isothermal sphere, whose projected surface density at an impact parameter  $r$  is given by

$$\Sigma(r) = k \frac{R_c}{r}; \quad (6)$$

where  $k$  can be written as  $k = \sigma_{\parallel}^2/2G$  kpc;  $\sigma_{\parallel}$  is the one-component central velocity dispersion; and  $G$  is the gravitational constant.

We then compute the equivalent width of a line using the (simplified) relation

$$W(r) = W_0(1 - e^{-\tau\Sigma(r)}), \quad (7)$$

where  $W_0$  is the equivalent width of a line for which  $r$  is 0 (i.e., the light crosses the center of the cloud);  $\tau$  is a parameter which takes into account the opacity of the cloud. In our simple model,  $W_0$  and the product  $\tau k$  are adjusted in order to match the observed distribution of equivalent widths. With the restriction that it must be larger than the maximum value of  $W$  in the sample, the precise value of  $W_0$  does not much change the following results, since the range of  $r$ -values for which large equivalent widths (between  $W_0$  and the largest value of  $W$ ) are produced is very small compared with  $R_c$ , which is now the value of the impact parameter giving rise to a line whose equivalent width has the lowest value in our sample. This relation is a reasonable approximation for clouds with opacity up to  $\sim 1$ .

Using the distribution in redshifts of the single lines, we can compute the probability density  $P(W_A, W_B)$  of having a line whose equivalent width in spectrum A is  $W_A$ , while the one for its corresponding line in spectrum B is  $W_B$ . The logarithm of  $P(W_A, W_B)$  is presented in Figure 9a for  $R_c = 5h_{50}^{-1}$  kpc and in Figure 9b for  $R_c = 20h_{50}^{-1}$  kpc. The discrete structure of the

probability density is caused by the discrete distribution of the separation between the light beams at the redshift of the Ly $\alpha$  clouds. The inner pattern is produced by the clouds closest to the quasar, while the widest pattern is due to the most remote ones. The probability is very high for the low values of equivalent widths, but with a very small dispersion along the line  $W_A = W_B$ ; the dispersion increases more and more along the line of equal equivalent widths, limited by two asymptotic branches. On the other hand, the dispersion is much larger for  $R_c = 5h_{50}^{-1}$  kpc than for  $R_c = 20h_{50}^{-1}$  kpc; in other words, the allowed region, i.e., the number of possible combinations of  $(W_A, W_B)$ , is much larger for smaller values of  $R_c$ .

Such behavior can easily be explained: the ratio of equivalent widths between a line in spectrum A and the corresponding line in spectrum B is sensitive to the density gradient of the cloud. If a line is produced near the center of a cloud, it is very unlikely that its equivalent width will be very similar to that of the corresponding line in the other spectrum, if the typical separation between the light beams is greater than about  $0.05R_c$  to  $0.1R_c$  (as one can see in comparing Figs. 9a and 9b), since the probability that impact parameters are equal (or have similar values) is much smaller than the probability that they are very different (relative to the cloud size). This produces a dispersion of the points in a graph of  $W_B$  versus  $W_A$ , which is very similar to a noncorrelation if the number of points is small. A strong correlation between equivalent widths of coincidences is then expected only if the typical radius of a cloud is much larger than the typical separation between the light beams. Note that graphs with similar properties can be produced for spherical clouds whose equivalent width decreases more quickly than linearly with  $R_c$ .

The observed correlation between  $W_A$  and  $W_B$  is then once more remarkable. This means the following:

- Both light beams actually cross the same clouds.
- The typical radius of a cloud  $R_c$  is much larger than the typical separation ( $1h_{50}^{-1}$  kpc) between the light beams, since




FIG. 9a




FIG. 9b

FIG. 9.—Logarithm of the probability that a line of equivalent width  $W_A$  occurs in spectrum A when one of  $W_B$  occurs in spectrum B, for an isothermal sphere: (a) with  $R_c = 5h_{50}^{-1}$  kpc; (b) with  $R_c = 20h_{50}^{-1}$  kpc. Contour plots are  $-7.5$  to  $0$ , with steps of  $0.5$ . Contours for lower probabilities are not shown.

the dispersion of the points is still compatible with zero; a maximum-likelihood determination gives a lowest value of about  $50h_{50}^{-1}$  kpc with the single lines, which should be taken only as indicative, because of the very small dispersion of the observed points.

c) As a consequence of item *b*, the available data are not yet sufficient to determine the value of  $R_c$  and the density profile for spherical Ly $\alpha$  clouds.

#### 4.2.5. Clumpiness of the Lyman-Alpha Clouds

The very good correlation observed between the equivalent widths of the lines detected in both spectra means that the two light beams actually cross the same clouds, but does not exclude the possibility that they are clumpy. Let us make some simple assumptions. As an extreme case, we use a model where a Ly $\alpha$  cloud is composed of a very large number of small individual cloudlets in an empty "intercloudlet" medium. First we assume that a given cloudlet contributes an amount  $W_c$  (in mean) to the (total) equivalent width of a Ly $\alpha$  line, i.e., that the cloudlets and the clouds are optically thin. Hence, we may write  $W_A = n_c^A W_c$  and  $W_B = n_c^B W_c$ , where  $n_c^A$  and  $n_c^B$  are the numbers of cloudlets in one cloud on the lines of sight to images A and B, respectively. Second, we can say that, within the same cloud, the light paths to A and B are comparable, so that  $n_c^A$  and  $n_c^B$  should both fall in a range  $[n_c - \sigma_{n_c}, n_c + \sigma_{n_c}]$ , where  $n_c$  is the mean number of cloudlets in the region probed by the light beams. (In this case we interpret the observed correlation between the equivalent widths  $W_A$  and  $W_B$  as due to a very large number of cloudlets.) We can assume that the number of cloudlets encountered along the line of sight to an image follows Poisson statistics, so that  $\sigma_{n_c} = (n_c)^{1/2}$ . Finally, because of the measurement errors, we have:  $\sigma_{\log(W_A/W_B)}^2 > \sigma_{\log(n_c^A/n_c^B)}^2$ , which gives  $n_c > 2/\sigma_{\log(W_A/W_B)}^2$ . In this way, the dispersion observed in Figure 5 could be interpreted as being due partly to the clumpiness of the cloud. Numerically,  $n_c > 550$ , so that we obtain a roughness factor  $\sigma_{n_c}/n_c < 0.04$ , corresponding to statistical Poissonian fluctuations if the model is taken literally. If Ly $\alpha$  clouds are more similar to a clumpy continuous medium, the number of clumps can be less numerous. In conclusion, the clouds must be very smooth on scale ranges from  $3h_{50}^{-1}$  pc to  $2000h_{50}^{-1}$  pc, as an extreme case in terms of differences between the number of clumps encountered along the two lines of sight.

### 4.3. Velocity Differences

Measurements of the velocity difference between two corresponding lines in the spectra of images A and B are also of interest. Dependence of this parameter on the equivalent widths of the lines or on the linear separation and/or redshift will be studied in this section.

#### 4.3.1. Individual Velocity Differences

In order to obtain the most precise values of the velocity differences  $\Delta V = c(\lambda_A - \lambda_B)/\lambda$ , we have used the cross-correlation technique developed by Tonry & Davis (1979) on individual single lines. Each line was first extracted from the spectra, to avoid (or to reduce as much as possible) any contamination of the cross-correlation function by its neighbors. The results are presented in Table 2, together with the formal errors  $\sigma_{\Delta V}$ . The difference in velocities is equal to zero within the error bars, except in very few cases: (a) the nonzero value of  $\Delta V$  for the line at 3726 Å is very probably due to the line shape at zero intensity; (b) nonzero differences are detected at the  $2\sigma$

level for the lines at 3792 Å ( $-9.4 \pm 3.5$  km s $^{-1}$ ), 3991 Å ( $9.0 \pm 3.7$  km s $^{-1}$ ), and 4372 Å ( $21 \pm 7.8$  km s $^{-1}$ ). The velocity differences for the blended lines are also essentially compatible with 0 km s $^{-1}$  within the error bars, except for the line at 4259 Å ( $-11.6 \pm 3.0$  km s $^{-1}$ ).

The histogram of the difference of velocities for the single and blended lines is presented in Figure 10. The standard deviation is found to be 17 km s $^{-1}$ .

No evidence of correlation is found between the equivalent widths and the difference in velocities, which argues against the presence of systematic mass motions linked to density gradients.

#### 4.3.2. Evolution

There is no indication of any dependence of the dispersion of the ratio  $W_A/W_B$  or of the difference in velocities on the combined effect of the redshift and the separation of the light beams.

We present in Figure 11 the graph of  $W_A/W_B$  versus  $z$  as an example.

## 5. ANALYSIS OF THE METALLIC ABSORPTION-LINE SYSTEMS

UM 673 gives new information on the spatial distribution, the column density, and the ionization of the metallic systems, since several metallic absorption-line systems are found or could be expected in our spectra. In order of increasing redshift, they are as follows:

- $z_{\text{abs}} = 0.17$  (redshift of the galaxy D south of UM 673).—No convincing identification could be made; several lines in the bluest part of the 1 Å spectrum could be Mg I  $\lambda 2852$ .
- $z_{\text{abs}} = 0.4261$ .—This possible system is discussed above. Any confirmation of its existence is very important for the study of the Ly $\alpha$  clouds; if it is real, then the distribution of Mg II varies over scales of  $16h_{50}^{-1}$  kpc, similar to that for the 1.483 and 1.491 systems of Q2345+007. This is also to be compared with the small but significant differences found for the C IV lines of the  $z_a = 1.1249$  system of the gravitational lens




FIG. 10.—Histogram of the difference in velocities for all the Ly $\alpha$  lines






FIG. 11.— $|W_A/W_B|$  vs.  $z$  for the single lines

quasar Q0957+561 images A and B (Boksenberg & Sargent 1983), whose line of sight at this redshift are separated by  $5.5h_{50}^{-1}$  kpc.

c)  $z_{\text{abs}} = 0.493$  (redshift of the lensing galaxy).—Ca II H and K lines plus probable Na I D1 and D2, believed to be formed in the lensing galaxy (cf. Paper II). These lines are only detected in component B, closer to the center of the galaxy than A. Our spectra do not detect other associated lines.

d)  $z_{\text{abs}} = 1.9406, 1.9417, 1.9436, 1.9441$ .—Values first reported from SBS. The first two redshift systems, revealed only by C IV lines in SBS are detected in the higher resolution spectra but are blended in the  $2 \text{ \AA}$  spectra. Refer to § 5.1 for more details. Figure 5c presents  $W_B$  versus  $W_A$  for all the  $2 \text{ \AA}$  coincidence lines of this complex.

e) The C IV system at  $z_{\text{abs}} = 2.3245$  (SBS).—This system does not show any line in the spectral range covered by our data, not even Ly $\alpha$ ; in fact the signal-to-noise ratio of our 5000–6000  $\text{\AA}$  spectrum is not sufficient to detect this doublet either.

f)  $z_{\text{abs}} = 2.3556, 2.3566$ .—First reported in Paper II, then in SBS. This complex system is detected in both  $2 \text{ \AA}$  spectra (see § 5.2). Figure 5d presents  $W_B$  versus  $W_A$  for all the  $2 \text{ \AA}$  coincidence lines of this complex.

The possible  $z_{\text{abs}} = 1.8987$  system (Paper II) is not confirmed by our new data.

Tables 4A–4G give a summary of the properties of the  $z = 1.94$  and  $z = 2.35$  complexes, described also below. The best values for the study of the heavy-element systems are of course found in the higher resolution spectra, while an attempt at comparison of the two lines of sight can be made with the  $2 \text{ \AA}$  spectra. In general this is difficult since most of the lines are blended with Ly $\alpha$  or metal lines. Hopefully, we know already that the equivalent widths of the Ly $\alpha$  lines are nearly equal in the A and B spectra.

### 5.1. The Complex System at $z_{\text{abs}} = 1.94$

Nine lines are detected at the  $5 \sigma$  level in the  $2 \text{ \AA}$  spectra of both components A and B. The four subcomponents are revealed mainly by their C IV doublets, which are responsible for the complicated structure of the involved lines. For example, C IV  $\lambda 1548$  of the  $z = 1.9441$  system falls on the wing of C IV  $\lambda 1550$  of the  $z = 1.9406$  system. The O I  $\lambda 1302$  line is probably detected for all the subcomponents. The C II  $\lambda 1334$

line is found only in the 1.9406 and 1.9417 systems. The Si IV doublet is only possibly detected in the 1.9417 system, but is unambiguous in the 1.9406 one. Other metallic lines (Al II  $\lambda 1670$ , Si II  $\lambda 1260$ , and possibly Si II  $\lambda 1193$ , Si III  $\lambda 1206$ ) are found only in the latter system. Except perhaps for the Si II lines, there is no significant difference between the equivalent widths in the A and B  $2 \text{ \AA}$  spectra. Velocity differences are also equal to zero within the error bars.

We may conclude that there is no dramatic change in the column density of the region which gives rise to this system on scales of  $2.1h_{50}^{-1}$  kpc. It is worthwhile to remember that there are differences on larger scales ( $10h_{50}^{-1}$  kpc to  $91h_{50}^{-1}$  kpc, depending on the redshift of the lensing galaxy) for the 1.483 and 1.491 systems of Q2345+007 (Tyson et al. 1986).

### 5.2. The Saturated Lyman-Alpha System at

$$z_{\text{abs}} = 2.3556, 2.3566$$

Seven lines are detected in both spectra, and six lines are detected only in the spectrum of image A for this complex. The identification of the O VI lines is far from certain; we mention it because of the expected equivalent width ratio and of the poorer wavelength calibration in the bluest part of the  $1 \text{ \AA}$  spectrum. If they are confirmed, then the two light beams probe regions with different ionization states. We note that Ly $\gamma$  is detected in the  $2 \text{ \AA}$  spectrum of A and is also present at the  $3 \sigma$  level in the spectrum of B. The existence of two components separated by  $300 \text{ km s}^{-1}$  is ascertained by the double structure of the Si II  $\lambda 1260$ , Si III  $\lambda 1206$ , Si IV, and C IV lines. The N I  $\lambda \lambda 1134$  and  $1200$  and the N II  $\lambda 1083$  lines are detected only in the  $z = 2.3566$  system. There is no significant difference between the equivalent widths of the lines of this complex between the  $2 \text{ \AA}$  spectrum of images A and B. We attribute the significant difference of velocities between the Ly $\alpha$  lines in both  $2 \text{ \AA}$  spectra as due to the shape of the line at zero intensity: if it were real, such a difference would have been expected also for the Ly $\beta$  lines. Otherwise, the difference in velocities between corresponding lines in the two spectra is comparable with zero within the error bars.

## 6. CONCLUSIONS

The size of Ly $\alpha$  clouds is a basic parameter required to understand their nature and eventually their origin. An upper limit of  $800h_{50}^{-1}$  kpc has been fixed by the observations of relatively close pairs of quasars (Shaver & Robertson 1983). However, gravitational lenses offer to probe the Ly $\alpha$  clouds on much smaller scales. A first lower limit was obtained with PG 1115+080 (Weymann & Foltz 1983), then improved with the lens system Q2345+007 (Foltz et al. 1984). Unfortunately, because of the faintness of the images and because the deflector has not yet been identified, it is not possible to exploit the latter case fully. However, it is interesting to note that a correlation between the rest equivalent widths of the coincidences does exist in this case too. UM 673 is more useful, in spite of a smaller separation between the light beams.

The results of our analysis of the spectra of the gravitational lens system UM 673 images A and B are given below.

### 6.1. For the Lyman-Alpha Cloud Lines and the Metallic System Lines

a) All the 68 lines detected at  $5 \sigma$  in the spectrum of the fainter component appear also in the spectrum of the brighter one. Because of the difference in the signal-to-noise ratio, the opposite is not true; however, a number of “missing” lines at



TABLE 4A  
HEAVY-ELEMENT SYSTEM AT  $z_{\text{abs}} = 1.9406$

	B (2 Å)			$\Delta V$	A (2 Å)			A (1 Å)			A (0.6 Å)			A (0.33 Å)		
	$\lambda$	$W_{\text{res}}$	$z$		$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$
B*	3506.85	0.32 ± 0.10	1.9388)		3506.70	0.87 ± 0.11	1.9387	3508.57	0.095 ± 0.003	1.9402						?Si II $\lambda$ 1193
S	3547.84	0.57 ± 0.14	1.9406)		3548.10	0.56 ± 0.05	1.9408	3548.17	0.65 ± 0.01	1.9409						Si III $\lambda$ 1206
B	3574.06	1.23 ± 0.12	1.9400	+ 8.80 ± 15.9	3574.24	1.28 ± 0.05	1.9401	3574.05	1.33 ± 0.01	1.9400						Lya
B		0.29			3706.87	0.49 ± 0.04	1.9410	3706.47	0.27 ± 0.01	1.9406						Si II $\lambda$ 1260
B	3924.61	0.36 ± 0.06	1.9408	- 5.80 ± 24.2	3924.29	0.43 ± 0.04	1.9406	3924.55	0.55 ± 0.01	1.9408						O I $\lambda$ 1302
B	4096.78	1.31 ± 0.13	1.9394	<sup>b</sup>	4096.69	1.29 ± 0.05	1.9393	4098.30	0.72 ± 0.01	1.9405						C II $\lambda$ 1334
B	4122.59	1.40 ± 0.16	1.9389	+ 44.8 ± 54.9	4123.17	1.25 ± 0.06	1.9393	4124.10	0.28 ± 0.01	1.9400						Si IV $\lambda$ 1393
B	4552.97	0.57 ± 0.04	1.9408	- 12.4 ± 7.0	4552.52	0.74 ± 0.02	1.9405									Si IV $\lambda$ 1402
B	4559.71	0.70 ± 0.05	1.9403	- 6.50 ± 29.2	4559.33	0.76 ± 0.02	1.9400									C IV $\lambda$ 1548
$\bar{z}$			1.9400				1.9400			1.9404						Al II $\lambda$ 1670

<sup>a</sup> See explanation of first column of Table 2, given in the Note to Table 2.

<sup>b</sup> The cross-correlation function presents two peaks of nearly equal intensity corresponding to  $\Delta V = -70.5 \pm 15.8 \text{ km s}^{-1}$  and  $40.3 \pm 23.4 \text{ km s}^{-1}$ .

TABLE 4B  
HEAVY-ELEMENT SYSTEM AT  $z_{\text{abs}} = 1.9417$

	B (2 Å)			$\Delta V$	A (2 Å)			A (1 Å)			A (0.6 Å)			A (0.33 Å)		
	$\lambda$	$W_{\text{res}}$	$z$		$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$
B	3578.79	1.67 ± 0.18	1.9439	- 3.1 ± 17.1	3578.89	1.44 ± 0.06	1.9440	3578.55	1.49 ± 0.02	1.9437						Lya
B	3832.39	0.62 ± 0.06	1.9431	+ 4.70 ± 9.00	3832.17	0.74 ± 0.03	1.9429	3831.51	0.445 ± 0.007	1.9424						?O I $\lambda$ 1302
B	3924.61	0.36 ± 0.06	1.9408	- 5.80 ± 24.2	3924.29	0.43 ± 0.04	1.9406	3924.55	0.55 ± 0.01	1.9408						C II $\lambda$ 1334
B	4096.78	1.31 ± 0.13	1.9394	<sup>a</sup>	4096.69	1.29 ± 0.05	1.9393									? Si IV $\lambda$ 1393
B	4122.59	1.40 ± 0.16	1.9389	+ 44.8 ± 54.9	4123.17	1.25 ± 0.06	1.9393	4126.03	0.43 ± 0.01	1.9413						? Si IV $\lambda$ 1402
B	4552.97	0.57 ± 0.04	1.9408	- 12.4 ± 7.7	4552.52	0.74 ± 0.02	1.9405									C IV $\lambda$ 1548
B	4559.71	0.70 ± 0.05	1.9403	- 6.50 ± 29.2	4559.33	0.76 ± 0.02	1.9400									C IV $\lambda$ 1550
$\bar{z}$			1.9410				1.9409			1.9415 <sup>b</sup>						

<sup>a</sup> The cross-correlation function presents two peaks of nearly equal intensity corresponding to  $\Delta V = -70.5 \pm 15.8 \text{ km s}^{-1}$  and  $40.3 \pm 23.4 \text{ km s}^{-1}$ .

<sup>b</sup> The Lya line was not used to compute the mean redshift.

TABLE 4C  
HEAVY-ELEMENT SYSTEM AT  $z_{\text{abs}} = 1.9436$

	B (2 Å)			$\Delta V$	A (2 Å)			A (1 Å)			A (0.6 Å)			A (0.33 Å)		
	$\lambda$	$W_{\text{res}}$	$z$		$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$
B	3578.79	1.67 ± 0.18	1.9439	- 3.1 ± 17.1	3578.89	1.44 ± 0.06	1.9440	3578.55	1.49 ± 0.02	1.9437						Lya
B	3837.50	0.24 ± 0.04	1.9470	+ 13.00 ± 23.70	3836.75	0.41 ± 0.03	1.9464	3834.13	0.48 ± 0.02	1.9444						? O I $\lambda$ 1302
B	4559.71	0.70 ± 0.05	1.9451	- 6.50 ± 29.2	4559.33	0.76 ± 0.02	1.9449									C IV $\lambda$ 1548
B	4565.23	0.31 ± 0.05	1.9438	+ 13.90 ± 16.40	4565.48	0.27 ± 0.02	1.9440									C IV $\lambda$ 1550
$\bar{z}$			1.9449				1.9448			1.9440						

TABLE 4D  
HEAVY-ELEMENT SYSTEM AT  $z_{\text{abs}} = 1.9441$

B (2 Å)				A (2 Å)				A (1 Å)				A (0.6 Å)				A (0.33 Å)			
$\lambda$	$W_{\text{res}}$	$z$	$\Delta V$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	ION
S	0.83			3551.73	0.60 ± 0.06	1.9438	3551.79	0.57 ± 0.01	1.9439										Si III $\lambda$ 1206
B	1.67 ± 0.18	1.9439	-3.1 ± 17.1	3578.89	1.44 ± 0.06	1.9440	3578.55	1.49 ± 0.02	1.9437										Ly $\alpha$
S	3837.50	0.24 ± 0.04	+13.0 ± 23.70	3836.75	0.41 ± 0.03	1.9464	3834.13	0.48 ± 0.02	1.9444										? O I $\lambda$ 1302
B	4559.71	0.70 ± 0.05	-6.50 ± 29.2	4559.33	0.76 ± 0.02	1.9449	4558.23	0.15 ± 0.01	1.9442	4558.55	0.180 ± 0.003	1.9444	4565.61	0.19 ± 0.02	1.9441	4565.96	0.051 ± 0.003	1.9443	C IV $\lambda$ 1548
B	4565.23	0.31 ± 0.05	+13.90 ± 16.40	4565.48	0.27 ± 0.02	1.9440													C IV $\lambda$ 1550
$\bar{z}$		1.9449		1.9446		1.9440									1.9441				1.9441

TABLE 4E  
HEAVY-ELEMENT SYSTEM AT  $z_{\text{abs}} = 2.3556$

B (2 Å)				A (2 Å)				A (1 Å)				A (0.6 Å)				A (0.33 Å)			
$\lambda$	$W_{\text{res}}$	$z$	$\Delta V$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	ION
(3264.02	1.84 ± 0.51	2.3562)		3264.66	1.14 ± 0.15	2.3568													Ly $\gamma$
	1.26			3279.00	1.02 ± 0.14	2.3561													C III $\lambda$ 977
S	3442.79	1.21 ± 0.14	+2.70 ± 12.9	3443.11	1.33 ± 0.06	2.3568	3442.32	1.47 ± 0.02	2.3560										Ly $\beta$
B	0.37			3463.78	0.60 ± 0.08	2.3566	3462.65	0.76 ± 0.02	2.3555										? O VI $\lambda$ 1031
B	(3481.49	0.46 ± 0.11	2.3553)	3481.45	0.38 ± 0.06	2.3552	3482.85	0.35 ± 0.04	2.3566										? O VI $\lambda$ 1037
B	0.19			4005.40	0.22 ± 0.03	2.3566													? Si III $\lambda$ 1193
B+	4049.27	0.80 ± 0.08	+2.10 ± 27.20	4049.65	0.75 ± 0.03	2.3565	4048.05	0.25 ± 0.01	2.3552	4048.05	0.25 ± 0.01	2.3552	4079.22	0.71 ± 0.04	2.3555	4079.22	0.71 ± 0.04	2.3555	Si III $\lambda$ 1206
B+	4080.62	1.97 ± 0.09	+21.0 ± 4.4	4080.46	1.92 ± 0.04	2.3565	4081.14	1.88 ± 0.01	2.3571										Ly $\alpha$
B	4229.34	0.63 ± 0.06	-9.00 ± 16.80	4229.10	0.78 ± 0.03	2.3553	4229.70	0.31 ± 0.04	2.3556	4229.70	0.31 ± 0.04	2.3556	4229.70	0.188 ± 0.006	2.3558	4229.76	0.188 ± 0.006	2.3558	Si II $\lambda$ 1260
B	4478.70	0.40 ± 0.04	-5.00 ± 7.30	4478.97	0.48 ± 0.02	2.3562	4477.94	0.22 ± 0.01	2.3554	4477.94	0.22 ± 0.01	2.3554	4478.25	0.224 ± 0.006	2.3557	4478.25	0.224 ± 0.006	2.3557	? C II $\lambda$ 1334
B	4677.40	0.50 ± 0.07	-6.05 ± 9.75	4677.56	0.50 ± 0.04	2.3561	4676.78	0.28 ± 0.01	2.3555	4676.78	0.28 ± 0.01	2.3555	4676.79	0.22 ± 0.01	2.3555	4676.79	0.22 ± 0.01	2.3555	Si IV $\lambda$ 1393
B	4707.58	0.32 ± 0.06	-2.00 ± 12.60	4707.40	0.26 ± 0.03	2.3558	4706.78	0.17 ± 0.01	2.3553	4706.78	0.17 ± 0.01	2.3553							Si IV $\lambda$ 1402
$\bar{z}$		2.3559		2.3560		2.3559 <sup>a</sup>			2.3554 <sup>a</sup>						2.3556				2.3556

<sup>a</sup> The Ly $\alpha$  line was not used to compute the mean redshift.

TABLE 4F  
HEAVY-ELEMENT SYSTEM AT  $z_{\text{abs}} = 2.3566$

	B (2 Å)			$\Delta V$	A (2 Å)			A (1 Å)			A (0.6 Å)			A (0.33 Å)			ION
	$\lambda$	$W_{\text{res}}$	$z$		$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	$\lambda$	$W_{\text{res}}$	$z$	
S	(3264.02	1.84 ± 0.51	2.3562)		3264.66	1.14 ± 0.15	2.3568										Ly $\gamma$
		1.26			3279.00	1.02 ± 0.14	2.3561										C III $\lambda$ 977
B	3442.79	1.21 ± 0.14	2.3565	+ 2.70 ± 12.9	3443.11	1.33 ± 0.06	2.3568	3442.32	1.47 ± 0.02	2.3560							Ly $\beta$
B	(3481.49	0.46 ± 0.11	2.3553)		3463.78	0.60 ± 0.08	2.3566	3462.65	0.76 ± 0.02	2.3555							? O VI $\lambda$ 1031
		0.37			3481.45	0.38 ± 0.06	2.3552	3482.85	0.35 ± 0.04	2.3566							? O VI $\lambda$ 1037
								3636.75	0.14 ± 0.01	2.3566							N II $\lambda$ 1083
								3808.88	0.08 ± 0.01	2.3569							N I $\lambda$ 1134
B	(4028.63	0.27 ± 0.06	2.3572)		4005.40	0.22 ± 0.03	2.3566	3995.11	0.18 ± 0.01	2.3561							? Si II $\lambda$ 1190
		0.19			4028.70	0.21 ± 0.03	2.3572	4028.09	0.23 ± 0.01	2.3567							? Si II $\lambda$ 1193
B+	4049.27	0.80 ± 0.08	2.3562	+ 2.10 ± 27.20	4049.65	0.75 ± 0.03	2.3565	4049.39	0.82 ± 0.01	2.2563	4027.89	0.31 ± 0.04	2.3566	4005.62	0.22 ± 0.01	2.3568	? Si II $\lambda$ 1200
B+	4080.62	1.97 ± 0.09	2.3567	+ 21.0 ± 4.4	4080.46	1.92 ± 0.04	2.3565	4081.14	1.88 ± 0.01	2.3571	4049.79	0.46 ± 0.02	2.3566	4028.09	0.25 ± 0.02	2.3567	N I $\lambda$ 1200
B	4229.34	0.63 ± 0.06	2.3555	- 9.00 ± 16.80	4229.10	0.78 ± 0.03	2.3553				4082.55	0.54 ± 0.04	2.3583	4949.64	0.78 ± 0.02	2.3565	Si III $\lambda$ 1206
B	4478.70	0.40 ± 0.04	2.3560	- 5.00 ± 7.30	4478.97	0.48 ± 0.02	2.3562							4081.16	1.72 ± 0.02	2.3571	Ly $\alpha$
B	4677.40	0.50 ± 0.07	2.3560	- 6.05 ± 9.75	4677.56	0.50 ± 0.04	2.3561				4479.44	0.15 ± 0.01	2.3566	4230.96	0.110 ± 0.06	2.3568	Si II $\lambda$ 1260
B	4707.58	0.32 ± 0.06	2.3560	- 2.00 ± 12.60	4707.40	0.26 ± 0.03	2.3558				4678.38	0.18 ± 0.01	2.3567	4479.74	0.125 ± 0.003	2.3568	? C II $\lambda$ 1334
											4708.54	0.14 ± 0.01	2.3566	4678.37	0.22 ± 0.01	2.3567	Si IV $\lambda$ 1393
$\bar{z}$																	Si IV $\lambda$ 1402
											2.3564						
																	2.3566 <sup>a</sup>

<sup>a</sup> The Ly $\alpha$  line was not used to compute the mean redshift.

TABLE 4G  
C IV DOUBLET OF THE  $z = 2.35$  COMPLEX

$\lambda$	$W_{\text{res}}$	$z$	Ion
5194.97	$0.16 \pm 0.01$	2.3555	C IV $\lambda 1548$
5203.60	$0.13 \pm 0.01$	2.2555	C IV $\lambda 1550$
5196.76	$0.35 \pm 0.02$	2.3566	C IV $\lambda 1548$
5205.39	$0.50 \pm 0.02$	2.3566	C IV $\lambda 1550$

the  $5\sigma$  level are detected at the  $3\sigma$  level. Following our definition (§ 3.1), we present evidence for two anticoincident lines.

b) The rest equivalent widths of the lines in spectrum B show a clear correlation with the corresponding lines in spectrum A.

c) The difference in velocities between the corresponding lines is zero within the measurement errors, except for six line pairs, among which two are probably due to the shape of the lines at zero intensity, three have a significance level of less than 2.7 (with a velocity difference less than  $21 \pm 7.8 \text{ km s}^{-1}$ ), and one has a significance level of 3.9 (with a velocity difference of  $11.6 \pm 3.0 \text{ km s}^{-1}$ ).

### 6.2. For the Lyman-Alpha Cloud Lines

We have used Monte Carlo simulations to obtain a best value for the lower limit of the diameter of Ly $\alpha$  clouds, assuming that the clouds are either spherical or oblate spheroids. We find that the diameter of spherical Ly $\alpha$  clouds must be larger than  $12h_{50}^{-1}$  kpc at a  $2\sigma$  confidence level. If the two anticoincidences are Ly $\alpha$  lines, then we obtain a  $2\sigma$  upper limit of  $160h_{50}^{-1}$  kpc. For oblate spheroids with an axis ratio less than 0.1, these values are doubled.

We have further used the equivalent width information as a tool to explore the structure of the clouds, assuming they are isothermal spheres. The correlation between equivalent widths is so good that we were only able to confirm that in this case the size of the Ly $\alpha$  clouds is at least 10 times the typical separation between the light beams. A maximum-likelihood calculation yields a lower limit of  $50h_{50}^{-1}$  kpc.

If the Ly $\alpha$  clouds are clumpy we estimate an upper limit to the roughness factor of 0.04. Hence, they must be homogeneous on scales from  $3h_{50}^{-1}$  pc to  $2000h_{50}^{-1}$  pc, given by the separation between the light beams for the most remote and the closest Ly $\alpha$  clouds. The velocity gradient across the clouds must be very small, but it is not significantly different from that observed for the heavy-element systems on comparable scales.

### 6.3. For the Metallic Line Systems

The complex metallic line systems do not show any significant variations on scales from  $0.7h_{50}^{-1}$  to  $2.2h_{50}^{-1}$  kpc, at  $z = 2.3566$  and  $z = 1.9406$ , respectively. In particular, the system at  $z = 2.356$  appears to be double at  $0.6 \text{ \AA}$  resolution; and it appears that the subcomponents themselves are quite similar. If the halos of galaxies do produce the metallic line systems, then their density and their ionization do not vary significantly over these scales, except perhaps for the Si II lines in the  $z = 1.9406$  system. If the two anticoincidences reported in the Ly $\alpha$  forest are due to a Mg II doublet, then the distribution of Mg II varies over scales of  $16h_{50}^{-1}$  kpc, comparable to that for the  $z = 1.483$  and  $1.491$  systems of Q2345+007.

### 6.4. For the Study of Gravitational Systems

If the actual size of the Ly $\alpha$  clouds is within the range given above, comparing the Ly $\alpha$  lines of two close quasar images should discriminate between the quasar pair and the gravitational lens hypothesis. In particular, if the two anticoincidences are actually Ly $\alpha$  lines, our results confirm the gravitational lens hypothesis for Q2345+007.

A. S. thanks B. Carswell and specially E. Gosset for discussion, the  $\Phi.V.$  for support, and M. Faucherre for his help during the edition of the main tables. This research was supported in part by the National Science Foundation under grants AST 87-00741 and 90-01181 (U.S.A.) and in part by contract ARC 90/94-140, "Action de recherche concertée de la Communauté Française" (Belgium). A. S. is grateful to ESO for the studentship under which this work was carried out.

### REFERENCES

- Banse, K., Grosbøl, P., Ponz, D., Ounnas, C., & Warmels, R. 1988, in Proc. Ninth Santa Cruz Summer Workshop in Astronomy, ed. L. B. Robinson (New York: Springer-Verlag), 431
- Blandford, R. D., & Kochanek, C. D. 1987, in Proc. Fourth Jerusalem Winter School for Theoretical Physics, Dark Matter in the Universe, ed. J. Bahcall, T. Piran, & S. Weinberg (Singapore: World-Scientific), 133
- Boksenberg, A., & Sargent, W. L. W. 1983, in Proc. 24th Liège Astrophys. Colloq., Quasars and Gravitational Lenses, ed. J. P. Swings (Liège: Institut d'Astrophysique), 500
- Bond, J. R., Szalay, A. S., & Silk, J. 1988, ApJ, 324, 627
- Carswell, R. F., Lanzetta, K. M., Parnell, H. C., & Webb, J. K. 1991, ApJ, 371, 36
- Foltz, C. B., Weymann, R. J., Röser, H.-J., & Chaffee, F. H., Jr. 1984, ApJ, 281, L1
- Hogan, C. B. 1987, ApJ, 316, L59
- Ikeuchi, S., & Ostriker, J. P. 1986, ApJ, 301, 522
- Lake, G. 1988, ApJ, 327, 99
- Magain, P., Remy, M., Surdej, J., Swings, J.-P., & Smette, A. 1990, in Proc. Workshop on Gravitational Lensing (Toulouse), ed. Y. Mellier, B. Fort, & G. Soucail (Berlin: Springer-Verlag), 88
- Milgrom, M. 1988, A&A, 202, L9
- Morton, D. C., York, D. G., & Jenkins, E. B. 1988, ApJS, 68, 449
- Pettini, M., Hunstead, R. W., Smith, L. J., & Mar, D. P. 1990, MNRAS, 246, 545
- Rees, M. J. 1970, ApJ, 160, L29
- . 1986, MNRAS, 218, 25
- Refsdal, S. 1964, MNRAS, 128, 307
- Sargent, W. L. W., Boksenberg, A., & Steidel, C. C. 1988, ApJS, 68, 539 (SBS)
- Sargent, W. L. W., Young, P., Boksenberg, M., & Tytler, D. 1980, ApJ, 42, 41 (SYBT)
- Shaver, P. A., & Robertson, J. G. 1983, ApJ, 268, L57
- Smette, A., Surdej, J., Shaver, P. A., Foltz, C. B., Chaffee, F. H., Jr., & Magain, P. 1990, in Proc. Workshop on Gravitational Lensing (Toulouse), ed. Y. Mellier, B. Fort, & G. Soucail (Berlin: Springer-Verlag), 122
- Steidel, C. C., & Sargent, W. L. W. 1990, AJ, 99, 1693
- Surdej, J. 1990, in Proc. Workshop on Gravitational Lensing (Toulouse), ed. Y. Mellier, B. Fort, & G. Soucail (Berlin: Springer-Verlag), 57
- Surdej, J., et al. 1987, Nature, 329, 695 (Paper I)
- . 1988, A&A, 198, 49 (Paper II)
- Tonry, J., & Davis, M. 1979, ApJ, 84, 1511
- Tyson, J. A., Seitzer, P., Weymann, R. J., & Foltz, C. 1986, AJ, 91, 1274
- Tytler, D. 1987, ApJ, 321, 49
- Verschueren, W., & Hensberge, H. 1990, A&A, 240, 216
- Weymann, R. J., & Foltz, C. B. 1983, ApJ, 272, L1
- Young, P., Gunn, J. E., Kristian, J., Oke, J. B., & Westphal, J. A. 1980, ApJ, 241, 507
- Young, P., Sargent, W. L. W., Boksenberg, A., & Oke, J. B. 1981, ApJ, 249, 415