

NOVEL STEALTHY Gd(III)-DOTA/POLYMER CONJUGATES FOR MAGNETIC RESONANCE IMAGING (MRI)

Mathurin Grogna^a, Stéphanie Bémelmans^a, Christian Vanasschen^b, Jean-François Desreux^b, André Luxen^c, Rudi Cloots^d, Christine Jérôme^a, Christophe Detrembleur^a


^aCenter for Education and Research on Macromolecules, University of Liège, Sart-Tilman, B6, B-4000 Liège

^bCoordination and Radiochemistry lab, University of Liège, Sart-Tilman, B16, B-4000 Liège, Belgium


^cCentre de Recherches du Cyclotron, University of Liège, Sart-Tilman, B30, B-4000 Liège, Belgium

^dLaboratoire de Chimie Inorganique Structurale, Université of Liège, Sart-Tilman, B6a, B-4000 Liège, Belgium

Magnetic resonance imaging (MRI) is a routine diagnostic tool in modern clinical medicine. MRI has many advantages as a diagnostic imaging modality. It is noninvasive, delivers no radiation, and has excellent (submillimeter) spatial resolution. Some Gadolinium(III) complexes are commonly used to enhance the contrast between adjacent tissues when the resolution/sensitivity of MRI is too low. Because free Gd^{3+} is very toxic in doses required for MRI, Gd(III) is chelated by poly(amino-carboxylate) such as diethylenetriamine pentaacetic acid (DTPA; scheme 1) or 1,4,7,10-Tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA; scheme 2)).



Scheme 1



Scheme 2

Although $DTPA/Gd^{3+}$ and $DOTA/Gd^{3+}$ are water soluble, they have a very short circulation lifetime in blood, a low molecular weight and a short rotational time that make the contrast poor. To enhance the contrast, the Gd^{3+} /complex doses have to be increased.

In order to increase the sensitivity of the technique, while not increasing the concentration of the contrast agent, we were investigating different strategies to improve (i) the circulation lifetime in blood, (ii) the relaxation rate of Gd(III) (and consequently, the contrasting efficiency) and (iii) the targeting of the contrast agent. This presentation aims at reporting how a multifunctional (co)polymer can be designed and exploited for improving the contrasting ability and bioavailability of gadolinium-based complexes.