A preliminary timing analysis of two intermediate polars: UU Col and Swift J0939.7-3224

Nikita RAWAT^{1,2,*}, J. C. PANDEY¹, Arti JOSHI², Srinivas M RAO¹ and Michaël DE BECKER³

¹ Aryabhatta Research Institute of observational sciencES (ARIES), Nainital 263001, India

² Deen Dayal Upadhyaya Gorakhpur University, Gorakhpur 273009, India

³ Pontificia Universidad Católica de Chile, Av. Vicuña Mackenna 4860, 782-0436 Macul, Santiago, Chile

⁴ Space Sciences, Technologies and Astrophysics Research (STAR) Institute, University of Liège, Quartier Agora, 19c, Allée du 6 Aôut, B5c, B-4000 Sart Tilman, Belgium

* Corresponding author: rawatnikita221@gmail.com

Abstract

We present the preliminary timing analysis of confirmed intermediate polar UU Col and possible intermediate polar Swift J0939.7-3224 in the optical band with the help of long-term, high-cadence continuous photometry from *Transiting Exoplanet Survey Satellite (TESS)*. For UU Col, we revise previously reported orbital and spin periods as 3.464 ± 0.005 h and 863.74 ± 0.08 s, respectively. Using the second harmonic of the beat frequency, the beat period is estimated as ~928 s. These findings indicate that UU Col is a disc-fed dominated disc-overflow accretor. For J0939, we establish the spin period as 2671.8 ± 0.8 s and refine the provisionally suggested orbital period as 8.49 ± 0.03 h. The absence of beat frequency in J0939 signifies that it might be a pure disc-fed accretor; however, an X-ray study of this source will help to understand its true nature.

Keywords: Cataclysmic Variable, Intermediate polars (UU Col, Swift J0939.7-3224), Accretion flow

1. Introduction

The modest magnetic field strength (B ~1-10 MG) subclass of magnetic cataclysmic variables (MCVs) is known as intermediate polars (IPs). These are semi-detached binaries where the magnetic field influences the accretion process that takes place in the system, as the matter is transferred from the companion (typically, a Roche-lobe filling late-type star) to the primary (white dwarf; WD). A full review of IPs is available in Warner (1983); Patterson (1994); Hellier (1995). For IPs, the spin period of WD (P_{ω}) is less than the orbital period (P_{Ω}) of the binary system (generally, P_{ω}/P_{Ω}~0.1). A truncated accretion disc can form due to a moderately strong magnetic field in these systems. Therefore, accretion occurs via disc or stream or a combination of both. The mass accretion rate, the magnetic field strength of

WD, and binary orbital separation are the key factors that determine which of the three accretion scenarios: disc-fed, stream-fed, and disc-overflow occurs in these systems. Because of the intricate interactions between the spin and orbital modulations, IPs are characterized by the existence of various periodicities in the X-ray and optical power spectra, which decide the governing accretion scenario. The sole presence of the spin frequency (ω) in the power spectrum depicts disc-fed accretion (Kim and Beuermann, 1995; Norton et al., 1996), while for a pure stream-fed accretion, there is strong power at the lower orbital sideband of the spin frequency, i.e., beat ($\omega - \Omega$) frequency (Hameury et al., 1986; Hellier, 1991; Wynn and King, 1992; Norton, 1993). Although ω frequency can also be present in the case of a stream-fed accretion if the magnetic poles are asymmetric in terms of polecap luminosities and size, the presence of $2\omega \cdot \Omega$ frequency decides between a pure stream-fed and dic-fed accretion (Wynn and King, 1992). On the other hand, for a disc-overflow accretion, both frequencies (ω and $\omega - \Omega$) should be simultaneously present in the power spectrum (Lubow, 1989; Armitage and Livio, 1996; Hellier, 1991, 1993). Disc-overflow accretion can further be described into three categories: disc-fed dominated disc-overflow accretor, stream-fed dominated discoverflow accretor, and disc-overflow with equal dominance depending on whether the dominating frequency is ω , $\omega - \Omega$ or equal dominance of both, respectively. IPs are fascinating candidates for understanding the physics of magnetically controlled accretion. Therefore, we selected two sources, namely UU Col and Swift J0939.7-3224, from the IP catalogue of Koji Mukai (https://asd.gsfc.nasa.gov/Koji.Mukai/iphome/catalog/alpha.html) with a motivation to ensure the IP nature and to inspect the nature of the accretion flow.

1.1. UU Col

UU Col (discovery name: RX J0512.2-3241) was identified as a soft X-ray intermediate polar by Burwitz et al. (1996). From B and V band CCD photometric data, they derived two periods of 3.45 ± 0.03 h and 863.5 ± 0.7 s, suggested to be P_{Ω} and P_{ω} , respectively. Burwitz et al. (1996) also found a very weak signal at 928.1 ± 0.9 s, which was proposed to be $P_{\omega-\Omega}$. From X-ray data, de Martino et al. (2006) found the dominant frequency corresponding to the period of 863.3 ± 1.4 s (P_{ω}) and very weak variability at $P_{\omega-\Omega} \sim 935$ s. The orbital period in X-ray data was found to be 3.55 ± 0.56 h. Katajainen et al. (2010) found that the polarization is modulated at the spin period of the WD in UU Col with amplitudes between +0.4% and -0.4% and +0.6% and -0.7% in the B band and I band, respectively. These values suggest that the polarized variability is most likely due to the cyclotron emission from two poles.

1.2. Swift J0939.7-3224

Swift J0939.7-3224 (hereafter J0939) was identified by Halpern and Thorstensen (2015) from *Swift*-XRT and *Swift*-UVOT images as a candidate MCV. From radial-velocity periodogram analysis and time-series photometric analysis, Halpern and Thorstensen (2015) found two probable periods in their data sets: 8.51 ± 0.02 h and 2670 ± 7 s, which were provisionally suggested to be P_Ω and P_ω, respectively. Apart from this, we are not aware of any other published results on this source.

2. Observations and Data

We have utilized *TESS* data of both targets for the present study. The *TESS* instrument consists of four wide-field CCD cameras, each with a field-of-view of $24^{\circ} \times 24^{\circ}$ so that all cameras can image a region of the sky measuring $24^{\circ} \times 96^{\circ}$. *TESS* observations are broken up into sectors, each lasting about 27.4 days and conduct its downlink of data while at perigee. The *TESS* bandpass extends from 600 to 1000 nm with an effective wavelength of traditional Cousins I-band, i.e., 786.5 nm (see Ricker et al., 2015, for details). We downloaded the data from Mikulski Archive for Space Telescopes (MAST) data archive (https://mast.stsci.edu/portal/Mashup/Clients/Mast/Portal.html) with unique identification numbers 'TIC 77841332' and 'TIC 25611385' for UU Col and J0939, respectively. The *TESS* observations of UU Col and J0939 were conducted at a cadence of 2 min from November 19, 2020, to December 16, 2020 (sector 32) and February 9, 2021, to March 6, 2021 (sector 35), respectively.




Figure 1: *TESS* light curve of UU Col spanning \sim 26 d.

3. Timing Analysis

3.1. UU Col

Figure 1 depicts the *TESS* light curve of UU Col, which, when closely examined, reveals short-term variations superimposed on long-term variability. In order to identify the periodicities in the data, we used Lomb-Scargle (LS) periodogram analysis (Lomb, 1976; Scargle, 1982)

in which an error in the peak is obtained by computing the half-size of a single frequency bin centred on the peak and then converting it to period units. The power spectrum obtained from *TESS* data is shown in the top panel of Figure 2, where we have marked the position of all dominant frequencies. The significance of these frequencies was obtained by computing the false alarm probability (FAP; Horne and Baliunas, 1986) and the dashed line implies a 90 % significance level, indicating that we are 90 % confident that the present frequencies are intrinsic in origin and not due to noise. The power spectral analysis revealed the presence of Ω , ω , 2ω , and $2(\omega - \Omega)$ frequencies in the data and the corresponding periods are provided in Table 1. The Ω frequency was found to be the dominant one, followed by ω and $2(\omega - \Omega)$.




Figure 2: Power spectra of UU Col and J0939, where all dominant frequencies are marked for clear visual inspection.

Table 1: Periods corresponding to dominant peaks in the LS power spectra of UU Col and J0939.

Object	\mathbf{P}_{Ω} (h)	$\mathbf{P}_{2\Omega}$ (h)	$\mathbf{P}_{\boldsymbol{\omega}}$ (s)	$\mathbf{P}_{2\omega}$ (s)	$\mathbf{P}_{2(\boldsymbol{\omega}-\boldsymbol{\Omega})}$ (s)
UU Col	3.464 ± 0.005		863.74 ± 0.08	431.87 ± 0.02	464.00 ± 0.02
J0939	8.49 ± 0.03	4.253 ± 0.008	2671.8 ± 0.8		

3.2. J0939

In Figure 3, we have shown the *TESS* light curve of J0939 and a clear variability pattern is observed. As discussed earlier, we have performed the LS periodogram analysis and the bottom panel of Figure 2 shows the corresponding power spectrum. Similar to UU Col, FAP was calculated and the dashed horizontal line denotes a 90 % significance level. The frequencies identified in the power spectrum are Ω , 2Ω , and ω . Due to the better time-cadence and longer observation duration in *TESS*, we obtained more precise values of periods than available in the

literature (see Table 1). Using these values of P_{Ω} and P_{ω} , the inferred value of $P_{\omega-\Omega}$ period comes out to be ~2927 s which corresponds to the frequency of ~29.5 c/d. We do not see any peak near ~29.5 c/d in our power spectrum, similar to the power spectrum of Halpern and Thorstensen (2015). Further, the Ω frequency dominates the power spectrum over ω frequency.




Figure 3: TESS light curve of J0939 spanning \sim 24 d.

4. Discussion and Conclusions

Using optical data from *TESS*, we have performed a preliminary timing analysis of two IPs, UU Col and J0939. For UU Col, we refine previously determined orbital and spin periods, whereas, for J0939, we confirm that ~2672 s is indeed the spin period of the WD. With $P_{\omega}/P_{\Omega} \sim 0.1$, both targets fall in the category of intermediate rotators. The power spectrum plays a crucial role in determining the mode of accretion in these systems based on the presence of spin, beat, orbital, and other sideband frequencies and their amplitudes. UU Col was identified as a disc-overflow accreting IP by de Martino et al. (2006) since they found a very weak signal at $P_{\omega-\Omega}$ in their dataset. Contrary to Burwitz et al. (1996) and de Martino et al. (2006), we have identified a clear peak at $2(\omega - \Omega)$ frequency in the power spectrum of UU Col, from which the inferred value of $P_{\omega-\Omega}$ comes out to be ~928 s. This value is more precise than those obtained by Burwitz et al. (1996) and de Martino et al. (2006). The presence of $2(\omega - \Omega)$ frequency in the power spectrum indicates that a part of the accreting material also

flows through the stream along with the majority of accretion taking place via a disc. It also suggests that the field distribution might have an up-down symmetry. Therefore, we conclude that UU Col is a disc-fed dominated disc-overflow accretor (see Table 2 for different accretion scenarios based on frequencies present in the power spectrum). Furthermore, the presence of the spin period in the TESS power spectrum of J0939 confirms that it is certainly an IP. The presence of ω frequency and absence of $\omega - \Omega$ indicates that J0939 is a pure disc-fed accretor (see Table 2), however, Ferrario and Wickramasinghe (1999) have demonstrated the significance of stream extensions for the presence of ω frequency in the power spectrum. As the azimuthal extension of the accretion stream in the orbital plane increases, power shifts from $\omega - \Omega$ to ω for stream-fed accretion. Moreover, Wynn and King (1992) have shown that stream-fed accretion can also produce a modulation at the spin frequency, in addition to that at the beat frequency, if there is an asymmetry between two magnetic poles. Hence, the presence of $2\omega - \Omega$ frequency in X-ray bands provides a clear identification between disc-fed and stream-fed systems and is only present in disc-less systems (Wynn and King, 1992). Therefore, X-ray observations of J0939 are needed to ascertain its nature further and see any change in governing accretion mechanism in the future. A variable disc-overflow accretion has been observed in many IPs, for example, TX Col (Norton et al., 1997; Rawat et al., 2021; Littlefield et al., 2021), FO Agr (Burwitz et al., 1996; Littlefield et al., 2020), V2400 Oph (Joshi et al., 2019), V902 Mon and Swift J0746.3-1608 (Rawat et al., 2022). In light of this, comprehensive time-resolved timing analysis of these targets is still required in the future.

Accretion mechanism	Frequencies present	Remarks	
Disc-fed accretion	ω and/or harmonics	Present in UU Col and J0939	
Stream-fed accretion	$\omega - \Omega$ and/or harmonics	Present in UU Col	
Disc-overflow accretion	ω , $\omega - \Omega$, and/or their har-	Present in UU Col	
	monics		
Disc-overflow with disc-fed	ω , $\omega - \Omega$, and/or their har-	Present in UU Col	
dominance	monics and ω is the domi-		
	nant frequency in comparison		
	with $\omega - \Omega$		

Table 2: Accretion mechanisms and frequencies present in the optical power spectrum of UU

 Col and J0939.

Acknowledgments

We thank the anonymous referee for reading our paper. NR and SMR acknowledge the Belgo-Indian Network for Astronomy and Astrophysics (BINA) consortium for the local support to attend the 3rd BINA workshop. This work includes data collected with the *TESS* mission, obtained from the MAST data archive at the Space Telescope Science Institute (STScI). Funding for the *TESS* mission is provided by the NASA Explorer Program. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 526555.

Further Information

ORCID identifiers of the authors

0000-0002-4633-6832 (Nikita RAWAT) 0000-0002-4331-1867 (J. C. PANDEY) 0000-0002-1303-6534 (Michaël DE BECKER)

Author contributions

The corresponding author worked on the data and the core writing, while other authors contributed to the discussion, reviewing, and editing of the content.

Conflicts of interest

The authors declare no conflict of interest.

References

- Armitage, P. J. and Livio, M. (1996) Accretion Disks in Interacting Binaries: Simulations of the Stream-Disk Impact. ApJ, 470, 1024. https://doi.org/10.1086/177928.
- Burwitz, V., Reinsch, K., Beuermann, K. and Thomas, H. C. (1996) A new soft intermediate polar: RX J0512.2-3241 in Columba. A&A, 310, L25–L28.
- de Martino, D., Matt, G., Mukai, K., Bonnet-Bidaud, J. M., Burwitz, V., Gänsicke, B. T., Haberl, F. and Mouchet, M. (2006) The X-ray properties of the magnetic cataclysmic variable UU Columbae. A&A, 454(1), 287–294. https://doi.org/10.1051/0004-6361:20065078.
- Ferrario, L. and Wickramasinghe, D. T. (1999) The power of intermediate polars. MNRAS, 309(2), 517–527. https://doi.org/10.1046/j.1365-8711.1999.02860.x.
- Halpern, J. P. and Thorstensen, J. R. (2015) Optical Studies of 13 Hard X-Ray Selected Cataclysmic Binaries from the Swift-BAT Survey. AJ, 150(6), 170. https://doi.org/10.1088/ 0004-6256/150/6/170.
- Hameury, J. M., King, A. R. and Lasota, J. P. (1986) Accretion flows in the non-synchronous magnetic cataclysmic variables. MNRAS, 218, 695–710. https://doi.org/10.1093/mnras/218. 4.695.
- Hellier, C. (1991) Do observations reveal accretion discs in intermediate polars ? MNRAS, 251, 693. https://doi.org/10.1093/mnras/251.4.693.

- Hellier, C. (1993) Disc-overflow accretion in the intermediate polar FO Aquarii. MNRAS, 265, L35–L39. https://doi.org/10.1093/mnras/265.1.L35.
- Hellier, C. (1995) The Accretion Geometry of Intermediate Polars. In Magnetic Cataclysmic Variables, edited by Buckley, D. A. H. and Warner, B., vol. 85 of Astronomical Society of the Pacific Conference Series, p. 185.
- Horne, J. H. and Baliunas, S. L. (1986) A Prescription for Period Analysis of Unevenly Sampled Time Series. ApJ, 302, 757. https://doi.org/10.1086/164037.
- Joshi, A., Pandey, J. C. and Singh, H. P. (2019) X-Ray Observations of an Intermediate Polar V2400 Oph. AJ, 158(1), 11. https://doi.org/10.3847/1538-3881/ab1ea6.
- Katajainen, S., Butters, O., Norton, A. J., Lehto, H. J., Piirola, V. and Berdyugin, A. (2010) Discovery of Polarized Emission from Two Soft X-ray-emitting Intermediate Polars: UU Col and NY Lup. ApJ, 724(1), 165–170. https://doi.org/10.1088/0004-637X/724/1/165.
- Kim, Y. and Beuermann, K. (1995) Spin-modulated radiation of intermediate polars. I. X-ray spectrum and light curves. A&A, 298, 165.
- Littlefield, C., Garnavich, P., Kennedy, M. R., Patterson, J., Kemp, J., Stiller, R. A., Hambsch, F.-J., Heras, T. A., Myers, G., Stone, G., Sjöberg, G., Dvorak, S., Nelson, P., Popov, V., Bonnardeau, M., Vanmunster, T., de Miguel, E., Alton, K. B., Harris, B., Cook, L. M., Graham, K. A., Brincat, S. M., Lane, D. J., Foster, J., Pickard, R., Sabo, R., Vietje, B., Lemay, D., Briol, J., Krumm, N., Dadighat, M., Goff, W., Solomon, R., Padovan, S., Bolt, G., Kardasis, E., Debackère, A., Thrush, J., Stein, W., Walter, B., Coulter, D., Tsehmeystrenko, V., Gout, J.-F., Lewin, P., Galdies, C., Fernandez, D. C., Walker, G., Boardman, J., James and Pellett, E. (2020) The Rise and Fall of the King: The Correlation between FO Aquarii's Low States and the White Dwarf's Spin-down. ApJ, 896(2), 116. https://doi.org/10.3847/1538-4357/ab9197.
- Littlefield, C., Scaringi, S., Garnavich, P., Szkody, P., Kennedy, M. R., Iłkiewicz, K. and Mason, P. A. (2021) Quasi-periodic Oscillations in the TESS Light Curve of TX Col, a Diskless Intermediate Polar on the Precipice of Forming an Accretion Disk. AJ, 162(2), 49. https://doi.org/10.3847/1538-3881/ac062b.
- Lomb, N. R. (1976) Least-Squares Frequency Analysis of Unequally Spaced Data. Ap&SS, 39(2), 447–462. https://doi.org/10.1007/BF00648343.
- Lubow, S. H. (1989) On the Dynamics of Mass Transfer over an Accretion Disk. ApJ, 340, 1064. https://doi.org/10.1086/167458.
- Norton, A. J. (1993) Simulation of the X-ray light curves of intermediate polars. MNRAS, 265, 316–328. https://doi.org/10.1093/mnras/265.2.316.
- Norton, A. J., Beardmore, A. P. and Taylor, P. (1996) On the interpretation of intermediate polar X-ray power spectra. MNRAS, 280(3), 937–952. https://doi.org/10.1093/mnras/280.3.937.

Norton, A. J., Hellier, C., Beardmore, A. P., Wheatley, P. J., Osborne, J. P. and Taylor, P. (1997) Stream-fed and disc-fed accretion in TX Columbae. MNRAS, 289(2), 362–370. https://doi.org/10.1093/mnras/289.2.362.

Patterson, J. (1994) The DQ Herculis Stars. PASP, 106, 209. https://doi.org/10.1086/133375.

- Rawat, N., Pandey, J. C. and Joshi, A. (2021) TESS Observations of TX Col: Rapidly Varying Accretion Flow. ApJ, 912(1), 78. https://doi.org/10.3847/1538-4357/abedae.
- Rawat, N., Pandey, J. C., Joshi, A. and Yadava, U. (2022) A step towards unveiling the nature of three cataclysmic variables: LS Cam, V902 Mon, and SWIFT J0746.3-1608. MNRAS, 512(4), 6054–6066. https://doi.org/10.1093/mnras/stac844.
- Ricker, G. R., Winn, J. N., Vanderspek, R., Latham, D. W., Bakos, G. Á., Bean, J. L., Berta-Thompson, Z. K., Brown, T. M., Buchhave, L., Butler, N. R., Butler, R. P., Chaplin, W. J., Charbonneau, D., Christensen-Dalsgaard, J., Clampin, M., Deming, D., Doty, J., De Lee, N., Dressing, C., Dunham, E. W., Endl, M., Fressin, F., Ge, J., Henning, T., Holman, M. J., Howard, A. W., Ida, S., Jenkins, J. M., Jernigan, G., Johnson, J. A., Kaltenegger, L., Kawai, N., Kjeldsen, H., Laughlin, G., Levine, A. M., Lin, D., Lissauer, J. J., MacQueen, P., Marcy, G., McCullough, P. R., Morton, T. D., Narita, N., Paegert, M., Palle, E., Pepe, F., Pepper, J., Quirrenbach, A., Rinehart, S. A., Sasselov, D., Sato, B., Seager, S., Sozzetti, A., Stassun, K. G., Sullivan, P., Szentgyorgyi, A., Torres, G., Udry, S. and Villasenor, J. (2015) Transiting Exoplanet Survey Satellite (TESS). Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003. https://doi.org/10.1117/1.JATIS.1.1.014003.
- Scargle, J. D. (1982) Studies in astronomical time series analysis. II. Statistical aspects of spectral analysis of unevenly spaced data. ApJ, 263, 835–853. https://doi.org/10.1086/160554.
- Warner, B. (1983) The intermediate polars. In IAU Colloq. 72: Cataclysmic Variables and Related Objects, edited by Livio, M. and Shaviv, G., vol. 101 of Astrophysics and Space Science Library, pp. 155–171. https://doi.org/10.1007/978-94-009-7118-9_20.
- Wynn, G. A. and King, A. R. (1992) Theoretical X-ray power spectra of intermediate polars. MNRAS, 255, 83–91. https://doi.org/10.1093/mnras/255.1.83.