

INTRINSIC MODIFICATION OF TENDON STRUCTURE AFTER CONCENTRIC OR ECCENTRIC TRAINING

e-Poster: 616

Congress: EFOST 2010

Type: efost2010

Topic: Basic Research

Authors: J.-F. Kaux¹, P. Drion¹, B. Besançon¹, V. Libertiaux¹, C. Jean-Louis¹, B. Forthomme¹, A. Colige¹, C. Le Goff¹, R. Franzen¹, J.-O. Defraigne¹, S. Cescotto¹, M. Rickert², J.-M. Crielaard¹; ¹Liège/BE, ²Heidelberg/DE

Any information contained in this pdf file is automatically generated from digital material submitted to e-Poster by third parties in the form of scientific presentations. References to any names, marks, products, or services of third parties or hypertext links to third-party sites or information are provided solely as a convenience to you and do not in any way constitute or imply EFOST's endorsement, sponsorship or recommendation of the third party, information, product, or service. EFOST is not responsible for the content of these pages and does not make any representations regarding the content or accuracy of material in this file.

As per copyright regulations, any unauthorised use of the material or parts thereof as well as commercial reproduction or multiple distribution by any traditional or electronically based reproduction/publication method is strictly prohibited.

You agree to defend, indemnify, and hold EFOST harmless from and against any and all claims, damages, costs, and expenses, including attorneys' fees, arising from or related to your use of these pages.

Please note: Links to movies, ppt slideshows and any other multimedia files are not available in the pdf version of presentations.

www.efost2010.com

1. Body

Introduction: It is well known that eccentric training is a successful way of treating chronic tendinopathy. Although clinical results are very positive, beneficial morphological and histological effects have not yet been elucidated. The aim of our experiment was to determine if there exist any intrinsic modifications in a tendon trained in concentric or eccentric modes, in a rat model, using an original method of measurement (cryo-jaws).

Methods: 18 rats were divided into 3 groups: 6 for the control group, without physical restraint; 12 for a training of 1 hour, 3 times a week, for 5 weeks, at a speed of 17m/min (1km/h), on a inclined treadmill: 6 rats running uphill at +15° for the concentric effort (group C) and 6 rats running downhill at -15° for the eccentric effort (group E) (Video 1). After this training period, the Achilles, patellar and tricipital tendons of both limbs were surgically removed in all 18 rats. Tendons taken from five rats of each group were subjected to a tensile test up to rupture using a "cryo" jaw (Video 2). Tendons of the remaining rat of each group were subjected to a histological study.

Results: The results (Table 1) showed significant changes in group E only: (1) an increase of the force required to rupture the patellar and tricipital tendons; (2) an improvement of the ratio between the force necessary to rupture the tricipital tendon and the body mass of the rats; (3) an increase of the surface area of the section of the tricipital tendon. No significant change was observed as far as constraint was concerned between groups. Histologically, we saw, in the group E, more peripheral blood vessels and a greater proportion of collagen.

Conclusion: This study showed that the mechanical properties of tendon tissue are enhanced by eccentric training. Tendons become stronger, the amount of collagen increases and there is probably more interaction between collagen fibers (mechanotransduction).

Keywords: tendinopathy, eccentric rehabilitation, mechanotransduction.